

**Factors affecting dispersal in a butterfly -  
A comparison between core and edge populations**

Inauguraldissertation

zur

Erlangung des akademischen Grades eines  
Doktors der Naturwissenschaften (Dr. rer. nat.)

der

Mathematisch-Naturwissenschaftlichen Fakultät

der

Universität Greifswald

vorgelegt von

Elisabeth Reim

Greifswald, Mai 2018

Dekan: Prof. Dr. Werner Weitschies

1. Gutachter: Prof. Dr. Klaus Fischer

2. Gutachter: Prof. Dr. Robby Stoks

Tag der Promotion: 03.09.2018

# TABLE OF CONTENT

ABSTRACT .....	5
<b>1 INTRODUCTION .....</b>	<b>11</b>
<b>2 SYNOPSIS .....</b>	<b>25</b>
2.1 Influence of sex and weather conditions on movement paths .....	27
2.2 Factors affecting emigration propensity and flight performance .....	29
2.3 Consequences of larval and adult food stress for flight performance and exploratory behavior .....	31
2.4 Comparison between core and edge populations .....	33
<b>3 REFERENCES .....</b>	<b>37</b>
<b>4 PUBLICATION LIST .....</b>	<b>51</b>
4.1 Movement patterns differ between sexes and depend on weather conditions in the butterfly <i>Lycaena tityrus</i> .....	55
4.2 Emigration propensity and flight performance are decoupled in a butterfly .....	79
4.3 Nutritional stress reduces flight performance and exploratory behavior in a butterfly .....	137
4.4 Successful despite poor flight performance: range expansion is associated with enhanced exploratory behavior and fast development .....	157
EIGENSTÄNDIGKEITSERKLÄRUNG .....	223
CURRICULUM VITAE .....	227
ACKNOWLEDGEMENTS .....	231



# **ABSTRACT**



## Abstract

In the current era of anthropogenic climate change is the long-term survival of all organisms dependent on their ability to respond to changing environmental conditions either by (1) phenotypic plasticity, which allows species to tolerate novel conditions, (2) genetic adaptation, or (3) dispersal to more suitable habitats. The third option, dispersal, allows individuals to escape unfavorable conditions, the colonization of new areas (resulting in range shifts), and affects patterns of local adaptation. It is a complex process serving different functions and involving a variety of underlying mechanisms, but its multi-causality though has been fully appreciated in recent years only. Thus, the aim of this doctoral thesis was to disentangle the relative importance of the multiple factors relevant to dispersal in the copper butterfly *Lycaena tityrus*, including the individual condition (e.g. morphology, physiology, behavior) and the environmental context (e.g. habitat quality, weather). *L. tityrus* is a currently northward expanding species, which makes it particularly interesting to investigate traits underlying dispersal. In the first experiment, the influence of weather and sex on movement patterns under natural conditions was investigated. Using the Metatron, a unique experimental platform consisting of interconnected habitat patches, the second experiment aimed to examine the influence of environmental factors (resources, sun) on emigration propensity in experimental metapopulations. Human-induced global change (e.g. climate change, agricultural intensification) poses a substantial challenge to many herbivores due to a reduced availability or quality of feeding resources. Therefore, in the third experiment, the impact of larval and adult food stress on traits related to dispersal ability was investigated. Additionally, the effect of different ambient temperatures was tested. In the fourth experiment, core (Germany) and recently established edge (Estonia) populations were compared in order to explore variation in dispersal ability and life history traits indicative of local adaptation. Dispersal is often related to flight performance, and morphological and physiological traits, which was investigated in experiments 2-4. Butterflies were additionally subjected to behavioral experiments testing for the individual's exploratory behavior (experiments 3 and 4).

Males and females differed substantially in morphology, with males showing traits typically associated with a better flight performance, which most likely result from

selection on males for an increased flight ability to succeed in aerial combats with rivaling males and competition for females. This pattern could be verified by mobility measures under natural conditions and flight performance tests. Interestingly, although females showed traits associated with diminished flight performance, they had a higher emigration propensity than males (though in a context dependent manner). Reasons might be the capability of single mated females to found new populations, to spread their eggs over a wide range or to escape male harassment. Conditions indicative of poor habitat quality such as shade and a lack of resources promoted emigration propensity. The environmental context also affected condition and flight performance. The presence of resources increased the butterflies' condition and flight performance. Larval and adult food stress in turn diminished flight performance, despite some reallocation of somatic resources in favor of dispersal-related traits. These detrimental effects seem to be mainly caused by reductions in body mass and storage reserves. A similar pattern was found for exploratory behavior. Furthermore, higher temperatures increased flight performance and mobility in the field, demonstrating the strong dependence of flight, and thus likely dispersal, on environmental conditions. Flight performance and exploratory behavior were positively correlated, probably indicating the existence of a dispersal syndrome. The population comparison revealed several differences between edge and core populations indicative of local adaptation and an enhanced dispersal ability in edge populations. For instance, edge populations were characterized by shorter development times, smaller size, and a higher sensitivity to high temperatures, which seem to reflect adaptations to the cooler Estonian climate and a shorter vegetation period. Moreover, Estonian individuals had an enhanced exploratory behavior, which can be advantageous in all steps of the dispersal process and may have facilitated the current range expansion.

In summary, these findings may have important implications for dispersal in natural environments, which should be considered when trying to forecast future species distributions. First, dispersal in this butterfly seems to be a highly plastic, context-dependent trait triggered largely by habitat quality rather than by individual condition. This suggests that dispersal in *L. tityrus* is not random, but an active process. Second, fast development and an enhanced exploratory behavior seem to facilitate the current range expansion. But third, while deteriorating habitat conditions are expected to promote dispersal, they may at the same time impair flight ability (as well as exploratory

behavior) and thereby likely dispersal rates. For a complete understanding of a complex process such as dispersal, further research is required.



# **INTRODUCTION**



# 1 Introduction

## **Anthropogenic climate change and options for organisms to respond**

Human-made climate change entails global warming as well as temperature anomalies and extreme weather events. These changes in turn have a high impact via effects on the water cycle, which can result in drought periods or heavy rainfall and floods (Rahmstorf & Coumou, 2011; Hansen *et al.*, 2012). The consequences for ecosystems and biodiversity will be enormous. Global warming is already causing substantial changes in abundance and phenology of organisms (Parmesan & Yohe, 2003; Parmesan, 2006; Chown *et al.*, 2010). Next to the above climatic changes, many other human-induced impacts on the environment can be observed, including land conversion resulting in habitat loss and fragmentation, the usage of environmental toxins and fertilizers, burning of fossil fuels or species overharvesting (Hansen *et al.*, 2012). Thus, long-term survival of all organisms is dependent on their ability to respond to these changing environmental conditions either by (1) phenotypic plasticity, which allows species to tolerate novel conditions, (2) genetic adaptation, or (3) dispersal to more suitable habitats, often resulting in range shifts (Warren *et al.*, 2001; Davis *et al.*, 2005; Hickling *et al.*, 2006; Berg *et al.*, 2010; Hofmann & Todgham, 2010).

The first option, phenotypic plasticity, describes the ability of a genotype to produce different phenotypes depending on environmental conditions. Thus, it represents direct environmental effects on the phenotype (Bradshaw, 1965; Fischer & Karl, 2010). These short term responses allow for maximum flexibility and might be crucial for species survival in the current era of climate change (Franks & Hoffmann, 2012; Kellermann *et al.*, 2012). But plastic responses are also costly, the most common cost is the maintenance of the sensory and regulatory machinery needed for plasticity, which frequently require energy and material expenses (DeWitt *et al.*, 1998; Pigliucci, 2005). Second, in the long term, environmental variation may induce evolutionary (genetic) adaptation, which is considered to be less costly, but often requires stable environments to evolve (DeWitt *et al.*, 1998). Further, even rapid evolutionary changes might be too slow to keep up with quickly changing environmental conditions (Williams *et al.*, 2008; Merilä, 2012). This leads us towards the third option, dispersal, allowing organisms to move away from unsuitable areas in order to find a better habitat to settle

(e.g. Matthysen, 2012). Hence, understanding the ultimate and proximate causes of dispersal will help to improve population management and predicting consequences of environmental changes on species persistence and distributions (Cote *et al.*, 2010a).

In this thesis, the focus will be mainly on the last option, dispersal, but nevertheless is the survival and maintenance of species often determined by a combination of all three options. Four main studies were conducted, focusing on factors underlying and facilitating dispersal. As study organism the butterfly *Lycaena tityrus* was used as butterflies in general are excellent model organisms in spatial ecology, especially in dispersal studies (see below; Settele *et al.*, 2009; Stevens *et al.*, 2010a; Ducatez *et al.*, 2012).

## **Dispersal**

Dispersal is of crucial importance in driving a species' capacity to respond to human-induced environmental changes like unpredictable climatic variation or habitat loss and fragmentation (Cote & Clobert, 2007; Travis *et al.*, 2013). It is often defined as any movements potentially leading to gene flow (Ronce, 2007) and a complex process serving different functions and involving a variety of underlying mechanisms (Bowler & Benton, 2005; Bonte *et al.*, 2012; Matthysen, 2012). Dispersal involves three consecutive phases, which are departure from the current patch, movement between the patches (transient phase), and settlement in a new patch (Bowler & Benton, 2005; Clobert *et al.*, 2009). The evolutionary causes of dispersal include the avoidance of kin competition, inbreeding, resource competition and environmental stochasticity (Bowler & Benton, 2005; Ronce, 2007; Clobert *et al.*, 2009). Dispersal can also incur costs, for instance energetic, time and opportunity costs (reviewed in Bonte *et al.*, 2012). Note that causes and costs might differ between the three stages of the dispersal process (Bowler & Benton, 2005; Clobert *et al.*, 2009; Bonte *et al.*, 2012). The dispersal of individuals influences their individual fitness, but has also consequences for population dynamics and genetics through metacommunity regulation, gene flow or speciation, as well as for species' distributions (Hanski & Gilpin, 1997; Hanski, 1999; Roff & Fairbairn, 2001; Clobert *et al.*, 2012; Baguette *et al.*, 2013). This might be of special importance for the rescue of declining populations or the recolonization of empty habitat patches

after local extinctions (Hanski & Gilpin, 1997; Hanski, 1999; Parmesan, 2006). Moreover, dispersal is a key process in the colonization of new areas to track shifting climate niches and thereby affecting patterns of local adaptation (Doebeli & Dieckmann, 2003; Hickling *et al.*, 2006; Ronce, 2007).

## **Quantifying dispersal and dispersal proxies**

The importance of predicting effects of dispersal on population dynamics seems to be clear, but practical problems associated with the study of dispersal often hinder accurate conclusions. Thus, theoretical studies play a particularly relevant role (Bowler & Benton, 2005). However, these model assumptions often lack a great deal of realism (Travis & French, 2000) and do not incorporate the condition-dependence of dispersal strategies (e.g. Hanski, 1994; Murrell *et al.*, 2002). Other studies found that models frequently underestimate the rate of long-distance dispersal, probably because these events are difficult to document and consequently also the rate of spread is hard to predict (Hastings *et al.*, 2005; Clark *et al.*, 2017). Furthermore, populations at range margins are often influenced by strong ecological and evolutionary forces, which are not or less experienced by populations within the range core (Lindstrom *et al.*, 2013). The difficulty of quantifying dispersal also depends on the model species. In insects, dispersal is relatively easy to measure in wing-dimorphic species, which often produce dispersive morphs characterized by wings and flight muscles (Roff & Fairbairn, 2007). In contrast, in wing-monomorphic insects (like butterflies), which comprise the vast majority of insect species, quantifying dispersal is more challenging as dispersal can be a continuously varying trait (Saastamoinen *et al.*, 2010, 2012) and more subtle changes in adult flight morphology are observed (Hill *et al.*, 1999, 2011). Thus, a number of proxies such as flight performance, wing shape and size or thorax mass and ratio (see below) are commonly used. Although the link between proxies and dispersal in the field has been frequently demonstrated (Hill *et al.*, 1999; Thomas *et al.*, 2001; Niitepõld *et al.*, 2009; Stevens *et al.*, 2010), these proxies might sometimes potentially measure components of flight that also share different functions (Saastamoinen *et al.*, 2012). Consequently, generalizations should be concluded carefully, and the potentially important role of proxies to examine dispersal evolution should not be neglected.

## **Multi-causality of dispersal and dispersal syndromes**

Traits potentially affecting dispersal may include morphology, ecology, physiology, behavior and overall condition (Bowler & Benton, 2005; Bonte & Saastamoinen, 2012; Stevens *et al.*, 2013; Therry *et al.*, 2014). Considering the various functions and mechanisms, the dispersal process is likely driven by a complex interaction between several factors rather than a single trait (Bowler & Benton, 2005; Matthysen, 2012; Legrand *et al.*, 2015). Furthermore, growing evidence indicates that dispersal is not a fixed but a highly plastic trait (Ronce, 2007; Clobert *et al.*, 2009; Clobert & Chaine, 2012). Organisms seem to be able to respond to environmental cues / conditions during all steps of the dispersal process (Bowler & Benton, 2005; Ronce, 2007). In a recent study of Legrand *et al.* (2015) dispersal propensity was linked to a wide range of factors, including direct (morphology and performance), interactive (sex ratio) and indirect / interactive effects (sex, habitat quality, weather) clearly demonstrating the importance of comprehensive approaches.

### **Morphology and physiology**

In most insects and many other animals, dispersal is related to flight performance. Flight performance in turn is influenced by morphological differences (Arnold, 1983; Gilchrist, 1990; Berwaerts *et al.*, 2002), which could also influence dispersal and colonization success, for instance in butterflies (Hill *et al.*, 1999; Sekar, 2012). Traits that have been associated with flight ability include wing size, wing aspect ratio, body size, wing loading, thorax mass, thorax-abdomen ratio, storage reserves, and flight metabolic rate (Betts & Wootton, 1988; Berwaerts *et al.*, 2002; Niitepõld *et al.*, 2009). For instance, high body mass may reflect an overall good condition, which can positively influence dispersal (Jenkins *et al.*, 2007). However, evidence exists for both larger and smaller individuals dispersing into new areas (reviewed in Bowler & Benton, 2005). A high body mass in combination with small wings leads to a high wing loading, which is likely detrimental for flight performance due to increased energetic costs (Wickman, 2009; Saastamoinen *et al.*, 2012). Almbro & Kullberg (2012) could demonstrate that a high wing loading reduced flight speed in male and led to a lower take-off angle in female *Pieris napi* butterflies. Nevertheless, high wing loadings have also been shown to be associated with fast flight (Betts & Wootton, 1988). Wing aspect

ratio, a trait that is related to the shape of the wing, is known to influence aerodynamic aspects of flight (Dudley, 2000; Hassall, 2015). High ratios have been shown to increase flight ability and acceleration capacity (Betts & Wootton, 1988; Berwaerts *et al.*, 2002; Berwaerts & Van Dyck, 2004). However, respective dispersal patterns are not consistent, and high as well as low ratios have been found to be associated with dispersing individuals (Hill *et al.*, 1999, 2011; Hassall, 2015). An important physiological trait associated with flight is fat, which represents the storage reserves of an organism. Fat is necessary to fuel flight activity and might thus also be a determinant of dispersal events (Zera *et al.*, 1998; Karl & Fischer, 2008). Note here that flight-related morphology / physiology could also show environmentally-induced plasticity, which may in turn feedback on dispersal ability (see below, environmental conditions). Many of the traits mentioned above are sex specific and likely result from differential selective pressures and thus causally underlie sexual differences in flight performance or dispersal strategies (Gilchrist, 1990; Van Dyck & Wiklund, 2002; Merckx & Van Dyck, 2005; Berwaerts *et al.*, 2006). Sex-biased dispersal has been shown in both directions, with males or females emigrating more frequently (Baguette *et al.*, 1998; Legrand *et al.*, 2015).

### **Behavioral and social factors**

Factors affecting dispersal may also include behavioral traits such as exploration, aggression, and sociability (Dingemanse *et al.*, 2003; Duckworth & Badyaev, 2007; Cote *et al.*, 2010b; Ducatez *et al.*, 2012). For instance, inter-individual differences in exploration behavior have been linked to variation in dispersal propensity (reviewed in Cote *et al.*, 2010a) with explorative individuals dispersing more likely (Fraser *et al.*, 2001; Dingemanse *et al.*, 2003; Cote *et al.*, 2010a). Pronounced exploration skills could be advantageous in all steps of the dispersal process (Ronce, 2007; Cote *et al.*, 2010b), in making the decision to leave a habitat patch, facilitating the movement itself, and increasing the likelihood to find and successfully settle in a suitable habitat (Ducatez *et al.*, 2012). Conspecific density has also been suggested to influence dispersal activity (Dethier & MacArthur, 1964). Interestingly, both negatively and positively density-dependent dispersal in butterflies have been reported (Enfjäll & Leimar, 2005). Furthermore, conspecific density itself could affect the habitat quality in an area, for instance if the density exceeds the local carrying capacity and emigration

rates consequently increase (Dethier & MacArthur, 1964). However, as already highlighted above, single traits are unlikely to represent phenotypic differences. These differences often imply several traits, which could potentially result in syndromes.

### **Dispersal syndromes**

Dispersal is an evolutionary and ecologically important process, but the motivation as well as the ability to disperse may differ strongly among populations and between individuals within populations (Van Dyck & Baguette, 2005; Baguette & Van Dyck, 2007; Ducatez *et al.*, 2014; Bestion *et al.*, 2015). In general, dispersing individuals do not seem to comprise a random subset of a given population. Rather, dispersers often possess several correlated life-history traits forming a so-called dispersal syndrome discriminating between dispersing and non-dispersing conspecifics (Bonte & Saastamoinen, 2012; Stevens *et al.*, 2013). Dispersal syndromes have been shown for a wide range of taxa (Fjerdingstad *et al.*, 2007; Moles & Westoby, 2014; Bestion *et al.*, 2015) and there is growing evidence for these syndromes in butterflies (Cote *et al.*, 2010a; Ducatez *et al.*, 2012; Saastamoinen *et al.*, 2012; Legrand *et al.*, 2015, 2016). In a recent study on *Pieris brassicae*, dispersing and resident individuals showed distinct behavioral and morphological attributes with good flight performers having longer wings and showing higher emigration rates (Legrand *et al.*, 2015). Further, flight performance in *P. brassicae* was also shown to be correlated with exploration skills (Ducatez *et al.*, 2012).

An individual's dispersal propensity also results from the interaction between the individual phenotype and local conditions (Clobert *et al.*, 2009). It seems that environmental conditions may have a considerable influence on traits related to dispersal (Saastamoinen *et al.*, 2012). Thus, dispersal syndromes are expected to show variation depending on environmental conditions (Ducatez *et al.*, 2014). Recent studies demonstrated that the environmental context might even prevail over phenotypic traits or social factors (e.g. Legrand *et al.*, 2015).

### **Environmental conditions**

Dispersal is a crucial process to deal with environmental variation (Cote & Clobert, 2007; Travis *et al.*, 2013). Furthermore, organisms are expected to base their dispersal

decisions on informative cues, like environmental circumstances (Dall *et al.*, 2005; Clobert *et al.*, 2009), which has rarely been tested so far (but see Cote & Clobert, 2007; Baguette *et al.*, 2011). Conditions that have been found to influence dispersal include habitat quality (e.g. resource availability) and weather conditions (e.g. temperature, wind; Legrand *et al.*, 2015; Kuussaari *et al.*, 2016).

Temperature is one of the most important ecological factors, affecting species survival, performance and determining their distributions (Overgaard & Sørensen, 2008; Sunday *et al.*, 2011). Consequently, current increases in Earth's mean temperatures are expected to strongly affect ecosystems and biodiversity (Hansen *et al.*, 2012; Stanton *et al.*, 2015). Especially ectotherms are vulnerable to global temperature changes because their basic physiological functions such as locomotion, growth, and reproduction are strongly determined by ambient temperatures (Deutsch *et al.*, 2008). Hence, also insect flight is a temperature-dependent process, which may in turn directly influence dispersal rates (Delattre *et al.*, 2013; Kuussaari *et al.*, 2016). In general, butterflies require high muscle temperatures between 28-38 °C for flight activity, which is achieved through basking in the sun (Dennis, 1993; Watt, 1995; Wickman, 2009). Shady conditions will therefore substantially reduce overall flight activity. Indeed, dispersal rate was shown to decrease with increasing cloudiness (Cormont *et al.*, 2011; Kuussaari *et al.*, 2016). Wind speed and rainfall could also negatively influence dispersal (Kuussaari *et al.*, 2016).

Habitat quality can be influenced by the degree of fragmentation, which is expected to impose strong selective pressures on dispersal rates (Heino & Hanski, 2001; Schtickzelle *et al.*, 2006). Host plant abundance and abundance of nectar-providing flowers are also known to affect dispersal probabilities (Öckinger & Smith, 2008). For instance, predicted changes in temperature and precipitation and / or exploiting land-use practices could negatively influence the growth and phenology of plants, which may in turn negatively affect food availability or quality for herbivores (Parmesan & Yohe, 2003; Parmesan, 2006; Pleasants & Oberhauser, 2013; Lebeau *et al.*, 2016). The resulting food stress might negatively affect flight, which is highly energy demanding, due to widespread effects on body size, storage reserves and physiology (Candy *et al.*, 1997; Suarez, 2000).

Dispersal and associated traits are often affected by environmental conditions experienced immediately before and during dispersal (Bonte & Saastamoinen, 2012). But also environmental conditions experienced during development and even in previous generations may affect dispersal-related traits or adult dispersal behavior (Karlsson & Van Dyck, 2005; Bonte *et al.*, 2008; Bonte & Saastamoinen, 2012). For instance, suboptimal conditions (e.g. during development) may decrease the investment into costly dispersal traits (Bonte *et al.*, 2008). On the other hand, such conditions may increase dispersal ability or at least the willingness to do so, as an adaptive response for dealing with detrimental conditions in later life (e.g. Monaghan, 2008). This has been demonstrated through alterations in body composition or behavior (Bonte *et al.*, 2008; Saastamoinen *et al.*, 2010; van den Heuvel *et al.*, 2013).

Following these chapters, the complexity and multi-causality of dispersal seem apparently clear, highlighting the importance of comprehensive approaches, which are still rare. Thus, the aim of this work was to disentangle the relative importance of multiple factors relevant to dispersal from the individual phenotype through to environmental conditions. Moreover, to predict whether a species can shift to new areas requires additionally an understanding of the colonization process at the expanding edge of the species range.

## **Range shifts and local adaptation**

The distribution of a species is determined by a suit of ecological and evolutionary processes (Gaston, 2003) and has evolved to maximize fitness in a particular ecological niche (MacArthur, 1972; Wingfield *et al.*, 2015). But recent environmental changes have impacted species habitats. At the southern range margins, we find environmental conditions shifting to warmer, unsuitable conditions, and corresponding range retractions or climate driven extinctions have become widespread (Parmesan *et al.*, 1999; Franco *et al.*, 2006; Thomas *et al.*, 2006). In contrast, at the northern margins conditions that were once too cold for species have now become suitable (Parmesan *et al.*, 1999; Parmesan, 2006; Bestion *et al.*, 2015). Considering rising temperatures, species could either adapt to new thermal conditions or they could track their shifting climate niche (see above, Parmesan, 2006). Several studies demonstrated poleward range shifts of species due to climate change (e.g. Parmesan *et al.*, 1999; Hickling *et*

*al.*, 2006). Thomas (2010), for instance, estimated numbers of about 68% of all species. A study on nonmigratory European butterflies with data being obtained from northern and southern range margins could show that about 63% have shifted their distribution ranges to the north (by 35-240 km, Parmesan *et al.*, 1999). For instance, *L. tityrus* was frequently observed in the Montseny region of central Catalonia in the 1920s, but sightings from the 1990s are only from the Pyrenees, which is located 50 km to the north (Parmesan *et al.*, 1999). At the same time, *L. tityrus* entered Estonia for the first time and quickly established several successful breeding populations. Recent populations already stem from north-eastern Estonia (Õunap & Tartes, 2014).

A sufficient dispersal ability is only one aspect of range expansions. Once a suitable habitat has been reached by dispersing individuals, the ability to establish populations beyond the current range is of crucial importance. In 1967, Janzen stated a climatic variability hypothesis declaring that individuals at higher latitudes should have broader ranges of thermal tolerance. There is as well evidence for an increased flexibility in physiological traits at higher latitudes to cope with often more fluctuating environmental conditions (Janzen, 1967; Naya *et al.*, 2012). The variation of conditions along geographical gradients seems to require an 'optimization' of phenotypic values and consequently local adaptation (Hoffmann *et al.*, 2002; Castañeda *et al.*, 2005). Hence, phenotypic differences between recently established edge and long established core populations are expected (Hanski *et al.*, 2004; Calosi *et al.*, 2008; Lindstrom *et al.*, 2013), which might arise from differences in selection pressures but could also result from differences in the genetic structure of populations (Volis *et al.*, 1998; Purves, 2009; Mägi *et al.*, 2011). For instance, a positive relationship between thermal tolerance and latitude has been reported for different taxa (Calosi *et al.*, 2008; Naya *et al.*, 2012). Next to this, the process of range expansion itself could entail a strong selection pressure (Van Petegem *et al.*, 2016) Given that recent populations are founded by the most dispersive individuals (cf. Hill *et al.*, 1998; Hanski *et al.*, 2002, 2004) and dispersal-related traits are heritable (Roff & Fairbairn, 2001), more dispersive genotypes should accumulate at the expansion front (Phillips *et al.*, 2010; Shine *et al.*, 2011). Indeed, increased dispersal abilities at the range front have been demonstrated (e.g. Phillips *et al.*, 2006; Mitikka & Hanski, 2010; Van Petegem *et al.*, 2016).

Depending on dispersal ability, establishing capacity and thermal tolerance some species may benefit while others may suffer from current climate change (Pimm, 2001; Thomas *et al.*, 2001; Klockmann *et al.*, 2016). Many temperate zone ectotherms actually benefit from rising temperatures (Pimm, 2001; Deutsch *et al.*, 2008; Bestion *et al.*, 2015), so does the study organism, which is spreading and shifting its range northward.

## Study organism

*Lycaena tityrus* (Poda, 1761), the sooty copper, is a butterfly of the Lycaenidae family (Fig. 1; Tolman & Lewington 1998). It is a widespread temperate-zone butterfly, ranging from Western Europe to central Asia. *L. tityrus* does not occur in Great Britain, Scandinavia and Siberia (Ebert & Rennwald, 1991). The species is bivoltine with two discrete generations per year in most parts of its range, although populations with one or three generations per year occur. Flight periods last from the end of April until June and from the beginning of July until September (Kolligs, 2014). Overwintering takes place as half-grown larva. The principal larval host-plant is *Rumex acetosa* L. (Polygonaceae), but several congeneric *Rumex* species are utilized as well (Ebert & Rennwald, 1991; Tolman & Lewington, 1998; Settele *et al.*, 2008). Adults are nectar feeders, using a broad array of species including several Asteraceae (Ebert & Rennwald, 1991). *L. tityrus* inhabits different habitat types, including unimproved grasslands, meadows, fenlands and forest edges (Ebert & Rennwald, 1991; Settele *et al.*, 2008). This species is currently expanding its range towards higher latitudes (see above) and altitudes, which is assumed to be largely driven by anthropogenic climate change (Brunzel *et al.*, 2008; Settele *et al.*, 2008; Klockmann *et al.*, 2016). For experiment 1 butterflies were individually tracked in the field. For the three following experiments (experiments 2-4) freshly eclosed, mated females from bivoltine populations were caught. Butterflies from the core of the range were sampled from three German populations and edge populations were sampled in north-eastern Estonia (Fig. 2). All females were transferred to climate chambers at Greifswald University for egg laying.



Figure 1 Top: *Lycaena tityrus* larvae (left side) and pupae (right side, <http://www.bkmakro.de>)  
Middle: *L. tityrus* lateral view (left side: male © Lothar Feisel, right side: female © Jan Sevcik)  
Below: *L. tityrus* top view (left side: male © Jens Philipp, right side: female © Robert Flogaus-Faust)




Figure 2      Sampling areas of *Lycaena tityrus* (map created with google maps)  
Blue: German populations (Waren, Guest, Ueckermünde; f.l.t.r.)  
Red: Estonian populations (Kasispea, Karepa, Mahu; f.l.t.r.)

# **SYNOPSIS**



## 2 Synopsis


### 2.1 Influence of sex and weather on movement paths

Habitat loss and fragmentation as well as climate change are important threats to biodiversity conservation (e.g. Forester & Machlis, 1996; Sanderson *et al.*, 2002). The concomitantly deteriorating habitat quality often forces individuals to disperse to new habitats (e.g. Hanski, 1998). Thus, dispersal is important for dealing with environmental variation (Cote & Clobert, 2007; Travis *et al.*, 2013). However, dispersal ability may differ within and among populations (Van Dyck & Baguette, 2005; Baguette & Van Dyck, 2007; Ducatez *et al.*, 2014; Bestion *et al.*, 2015) and is dependent on environmental conditions such as prevailing weather conditions (Legrand *et al.*, 2015; Kuussaari *et al.*, 2016). Moreover, males and females may differ in their dispersal behavior (e.g. Fischer *et al.*, 1999).

Against this background, the study investigated sexual differences in and environmental effects on movement patterns in the copper butterfly *Lycaena tityrus*. Therefore, flight paths of both sexes were individually tracked (max. 15 min) within a habitat patch and weather conditions were recorded in parallel. As GPS devices Android smartphones in combination with a tracking application were used. Additionally, it was examined whether such devices are suitable for investigating insect behavior. The spatially restricted scale only allowed to examine movement behavior rather than dispersal per se, but addressing such questions could still provide valuable knowledge to assess dispersal abilities of populations or species (Niitepõld *et al.*, 2009). During tracking, several parameters (e.g. sex, number of stops) were recorded and the tracking app was used to extract data (e.g. total track length, average movement speed). To assess whether a high activity may facilitate dispersal, the linear distance between the starting and end point of each track was measured. Weather data were obtained from the weather station closest to the study area.

The study revealed sex-specific differences in behavioral patterns of *L. tityrus*. Males spent more time flying and had longer track lengths and linear distances than females. These differences might result from males' higher activity in general (see also Kingsolver, 1983), presumably owing to mate location and territorial disputes (Fig. 3a;

Ebert & Rennwald, 1991; Fischer & Fiedler, 2000). In addition, males alighted more often on flowers than females, probably to fuel their high flight activity (Fig. 3b). Females showed a higher affinity to host-plants for egg-laying (Fig. 3c). They seem to generally spend less time flying and allocate more of their time to oviposition (Kingsolver, 1983; Fischer *et al.*, 1999). Effects of environmental factors were less pronounced, but at least support the notion that detrimental weather conditions may negatively affect flight activity and therefore dispersal in insects (Cormont *et al.*, 2011; Kuussaari *et al.*, 2016).



**Figure 3** Sexual differences in *Lycaena tityrus* in track length (a) and the number of stops on flowers (b) and *Rumex* host-plants (c).


In summary, these findings provide a striking example of sex-related differences in animal behavior. Furthermore, customary smartphones can apparently comprise suitable means to reveal biologically significant behavioral patterns. More problematic than the technical device used seems to be the challenge of following individual butterflies for long enough in the field. Thus, the results seem to be not appropriate to estimate rates of dispersal and dispersal distances, such that any conclusions regarding insect dispersal need to be drawn with the greatest care.

## 2.2 Factors affecting emigration propensity and flight performance in a butterfly

Dispersal is a complex process serving different functions and involving a variety of underlying mechanisms (Bowler & Benton, 2005; Bonte *et al.*, 2012; Matthysen, 2012), but its multi-causality though has been fully appreciated in recent years only (reviewed by Matthysen, 2012). Currently, there is a lack of studies aiming to disentangle the relative importance of multiple factors relevant to dispersal, including the individual condition (e.g. morphology, behavior) and the environmental context (e.g. habitat quality, weather; but see e.g. Legrand *et al.*, 2015; Turlure *et al.*, 2016). Also, growing evidence indicates that dispersal is not a fixed but a highly plastic trait (e.g. Ronce, 2007) and organisms might be able to collect information on their current environment influencing their dispersal decision (Clobert *et al.*, 2009). However, only a few studies have focused on such cues triggering dispersal decisions (but see Cote & Clobert, 2007; Baguette *et al.*, 2011). These deficiencies are worrying given the crucial ecological and evolutionary importance of dispersal (Cote & Clobert, 2007; Clobert *et al.*, 2012; Travis *et al.*, 2013).


Against this background, a multifaceted approach was used to uncover the relative importance of several potentially dispersal-relevant factors in the butterfly *Lycaena tityrus*, ranging from the individual condition through to environmental variation. Using the Metatron, a unique experimental platform designed to study dispersal in experimental metapopulations (Legrand *et al.*, 2012), it was examined to what extent emigration propensity is affected by the environmental context. Therefore, two-patch experimental meta-populations were used where butterflies were released into departure cages with different environmental conditions (varied availability of sun and / or resources), in which they could either stay or which they could leave. Furthermore, flight performance was measured to examine relations between movement patterns under semi-natural conditions and flight performance under laboratory conditions (Ducatez *et al.*, 2012; Legrand *et al.*, 2015). In addition, individual variation in several morphological (e.g. wing size and shape, thorax mass and ratio, fat content) parameters was assessed that may be directly or indirectly associated with movement ability (Legrand *et al.*, 2016).

The results show that males and females differed substantially in morphology. Males showed traits typically associated with a better flight performance, which most likely result from selection on males for an increased flight ability in order to succeed in aerial combats with rivaling males and in competition for females (Saastamoinen *et al.*, 2012). This could be verified in a flight performance test (Fig. 4). The environmental context affected condition and flight performance. The presence of resources in the departure cage increased the butterflies flight performance (Fig. 4).



**Figure 4** Flight performance (means + 1 SE) of *Lycaena tityrus* in relation to the presence or absence of resources in the departure cage. Males: filled bars; females: open bars

Interestingly, individual condition did not generally influence emigration propensity. Although females showed traits associated with diminished flight performance, they had a higher emigration propensity than males (when resources were present; Fig. 5). Reasons might be the capability of single mated females, in contrast to males, to found new populations or to escape male harassment (Hill *et al.*, 1999; Hopper, 1999; Trochet *et al.*, 2013). This indicates that flight performance and the first step of dispersal, i.e. the actual decision whether to disperse or not, do not necessarily equate. Moreover, conditions indicative of poor habitat quality such as shade and a lack of resources promoted emigration propensity (Fig. 5).



**Figure 5** Emigration propensity (%) in relation to the presence or absence of resources in the departure cage in *Lycaena tityrus*. Males: filled bars; females: open bars

This suggests that dispersal in *L. tityrus* is not random but an active process, following an accurate assessment of patch quality. Thus, dispersal in this butterfly is a highly plastic, context-dependent trait triggered largely by habitat quality rather than by individual condition (Saastamoinen *et al.*, 2012; Legrand *et al.*, 2015). Climate change is expected to frequently reduce habitat quality, but this will at the same time likely increase emigration propensity and thereby potentially facilitate range shifts in flying insects (Travis *et al.*, 2013). This should be considered when trying to forecast future species distributions.

### 2.3 Consequences of larval and adult food stress for flight performance and exploratory behavior


Dispersal sets the potential of individuals to spread, for instance away from local unfavorable conditions, to avoid competition or to more suitable habitats (Matthysen, 2012). Anthropogenic global change, including agricultural intensification (fertilization, pesticide use and high mowing frequencies) and climate change (changes in

temperature and precipitation), poses a substantial challenge to many herbivores due to a reduced availability or quality of feeding resources (Parmesan & Yohe, 2003; Pleasants & Oberhauser, 2013; Lebeau *et al.*, 2016). Resulting food stress is expected to detrimentally affect performance, amongst others in dispersal-related traits as flight is a highly energy demanding process (Candy *et al.*, 1997; Suarez, 2000). Larval food stress may additionally shape the adult's physiological state due to a reallocation of resources among competing body parts and functions such as reproduction, storage, and flight-related traits (McNamara & Houston, 1996; Boggs, 2009). Further, conditions experienced during development may affect the adult phenotype as a predictive adaptive response in order to be better prepared for dealing with detrimental conditions in later life (e.g. Monaghan, 2008). This has already been demonstrated for dispersal ability through alterations in body composition or behavior (Saastamoinen *et al.*, 2010; van den Heuvel *et al.*, 2013).

Therefore, this study investigated the impact of larval and adult food stress on traits related to dispersal ability in the butterfly *Lycaena tityrus*. In the larval food stress treatment, individuals experienced a period of food deprivation of 30 h to mimic a situation in which a larva has to leave a deteriorated host plant and search for a new one. To implement adult food stress, butterflies were exposed to a period of food deprivation for 2 days (i.e. the period from eclosion to testing), during which they were provided with water only. Following the treatments, all butterflies were first subjected to a tunnel test, testing for exploratory behavior, a trait that may reflect behavioral differences in butterflies (Ducatez *et al.*, 2012, 2014). Afterwards, butterflies were subjected to a flight performance test (Ducatez *et al.*, 2012; Legrand *et al.*, 2015). Both tests were conducted at different ambient temperatures. Additionally, several morphological and physiological traits were investigated (see 2.2).

The study revealed that inadequate nutrition during development and in the adult stage diminishes flight performance, despite some reallocation of somatic resources in favor of dispersal-related traits (Fig. 6a). Similar results were found for exploratory behavior (Fig. 6b). These detrimental effects seem to be mainly caused by reductions in body mass and storage reserves, which may have important implications for dispersal in natural environments. While deteriorating habitat conditions are expected to promote dispersal as an adaptive response (Legrand *et al.*, 2015), they may at the same time

impair flight ability and thereby likely dispersal rates. Higher temperatures increased flight performance, demonstrating the strong dependence of flight performance and presumably also dispersal on environmental conditions (Fig. 6a). As shown before (see 2.2), the sexes differed in morphology.



**Figure 6** Flight endurance in a vortex (a, means + 1 SE) in *Lycaena tityrus* in relation to adult feeding treatment and test temperature. Food: filled bars; no food: open bars. Flight distance covered in a tunnel (b, means + 1 SE) in *Lycaena tityrus* males and females in relation to larval feeding treatment. Males: filled bars; females: open bars

Exploratory behavior was found to be (moderately) repeatable at the individual level. Furthermore, flight performance and exploratory behavior were positively correlated, probably indicating the existence of a dispersal syndrome (e.g. for *Pieris brassicae* in Ducatez *et al.*, 2012). Thus, the individuals performing repeatedly better than others in different types of flight tests might be the ones more likely to disperse (Delattre *et al.*, 2013).

## 2.4 Comparison between core and edge populations


Human-induced environmental changes, like rising temperatures or habitat degradation, force species to respond. Next to phenotypic plasticity, which allows species to tolerate novel conditions, and genetic adaptation, dispersal and concomitant

range shift is one possible option (Parmesan, 2006; Berg *et al.*, 2010; Hofmann & Todgham, 2010). Dispersal allows individuals to escape unfavorable conditions, the colonization of new areas, and affects patterns of local adaptation (Doebeli & Dieckmann, 2003; Hickling *et al.*, 2006; Baguette *et al.*, 2013). Evidence suggests that dispersing individuals may differ in a set of phenotypic traits from conspecifics, indicating the existence of a dispersal syndrome (Bonte & Saastamoinen, 2012; Stevens *et al.*, 2013). Assuming that recent populations are founded by the most dispersive individuals (cf. Hill *et al.*, 1998; Hanski *et al.*, 2002, 2004) and dispersal-related traits are heritable (Roff & Fairbairn, 2001), more dispersive genotypes should accumulate in recently established populations (Shine *et al.*, 2011). Besides a good dispersal capacity, the ability to establish populations beyond the former range, once suitable habitats have been reached, is of crucial importance. Different environmental condition at range margins might require local adaptation (Hoffmann *et al.*, 2002; Castañeda *et al.*, 2005). Thus, phenotypic differences between recently established edge and core populations are expected, especially in traits related to dispersal, due to differences in selection pressures and genetic structure (Volis *et al.*, 1998; Purves, 2009; Mägi *et al.*, 2011).

Against this background, the study compared replicated core (Germany) and recently established edge (Estonia) populations of the currently northward expanding butterfly *Lycaena tityrus* (Brunzel *et al.*, 2008; Settele *et al.*, 2008; Klockmann *et al.*, 2016), which has colonized north-eastern Estonia very recently. The aim was to explore variation in dispersal ability and life history traits indicative of local adaptation. Larvae of both countries were reared at control (26°C, 60% relative humidity, L18:D6 photoperiod) and cool (18°C, 60% relative humidity, L18:D6 photoperiod) conditions. One day-old butterflies were first subjected to a behavioral experiment, testing for the individual's exploratory behavior towards a food source in an unknown experimental setup. On the following day butterflies were tested for flight performance (Ducatez *et al.*, 2012; Legrand *et al.*, 2015). Additionally, morphological and physiological parameters were investigated (see above, Berwaerts *et al.*, 2002).

The results show that sexes differed in developmental, morphological, and physiological traits (e.g. Gilchrist, 1990; Berwaerts *et al.*, 2006). In general, males showed traits associated with increased flight ability, and enhanced exploratory

behavior, which might be helpful to quickly establish a territory (Fischer & Fiedler, 2000; Saastamoinen *et al.*, 2012). Despite the very recent colonization, several differences between edge and core populations indicative of local adaptation were found. Edge populations were characterized by shorter development times and smaller size, a higher sensitivity to high temperature and an enhanced exploratory behavior (Fig. 7).



**Figure 7** Exploratory behavior (animals finding a food source within 90 minutes in %) in *Lycaena tityrus* males (filled bars) and females (open bars) in relation to origin (country) and developmental temperature.

These differences seem to partly reflect adaptations to the cooler Estonian climate and the concomitantly short vegetation period (or a loss of adaptation to warmer conditions). Enhanced exploratory behavior can be advantageous in all steps of the dispersal process and may have facilitated the current range expansion. It is currently unclear whether the above differences reflect clinal variation or an accumulation of cold-adapted and / or more dispersive genotypes in north-eastern edge populations. A contribution of parental effects can also not be excluded (Wolf *et al.*, 1998; Mattila & Hanski, 2014). In summary, this study provides knowledge which might be useful for a better understanding of species responses to environmental change, in particular insights into the factors associated with a current range expansion.



# REFERENCES



### 3 References

- Almbro, M. & Kullberg, C. (2012) Weight loading and reproductive status affect the flight performance of *Pieris napi* butterflies. *Journal of Insect Behavior*, **25**, 441–452.
- Arnold, S.J. (1983) Morphology, performance and fitness. *American Zoologist*, **23**, 347–361.
- Baguette, M., Blanchet, S., Legrand, D., Stevens, V.M. & Turlure, C. (2013) Individual dispersal, landscape connectivity and ecological networks. *Biological Reviews*, **88**, 310–326.
- Baguette, M., Clobert, J. & Schtickzelle, N. (2011) Metapopulation dynamics of the bog fritillary butterfly: experimental changes in habitat quality induced negative density-dependent dispersal. *Ecography*, **34**, 170–176.
- Baguette, M. & Dyck, H. Van. (2007) Landscape connectivity and animal behavior: functional grain as a key determinant for dispersal. *Landscape Ecology*, **22**, 1117–1129.
- Baguette, M., Vansteenwegen, C., Convi, I. & Nève, G. (1998) Sex-biased density-dependent migration in a metapopulation of the butterfly *Proclossiuna eunomi*. *Acta Oecologica*, **19**, 17–24.
- Berg, M.P., Toby Kiers, E., Driessen, G., Heijden, M. van der, Kooi, B.W., Kuenen, F., *et al.* (2010) Adapt or disperse: understanding species persistence in a changing world. *Global Change Biology*, **16**, 587–598.
- Berwaerts, K., Aerts, P. & Dyck, H. Van. (2006) On the sex-specific mechanisms of butterfly flight: flight performance relative to flight morphology, wing kinematics, and sex in *Pararge aegeria*. *Biological Journal of the Linnean Society*, **89**, 675–687.
- Berwaerts, K. & Dyck, H. Van. (2004) Take-off performance under optimal and suboptimal thermal conditions in the butterfly *Pararge aegeria*. *Oecologia*, **141**, 536–545.
- Berwaerts, K., Dyck, H. Van & Aerts, P. (2002) Does flight morphology relate to flight performance? An experimental test with the butterfly *Pararge aegeria*. *Functional Ecology*, **16**, 484–491.
- Bestion, E., Clobert, J. & Cote, J. (2015) Dispersal response to climate change: scaling down to intraspecific variation. *Ecology Letters*, **18**, 1226–1233.

- Betts, C.R. & Wootton, R.J. (1988) Wing shape and flight behaviour in butterflies (Lepidoptera: Papilionoidea and Hesperioidea): a preliminary analysis. *Journal of Experimental Biology*, **138**, 271–288.
- Boggs, C.L. (2009) Understanding insect life histories and senescence through a resource allocation lens. *Functional Ecology*, **23**, 27–37.
- Bonte, D., Dyck, H. Van, Bullock, J.M., Coulon, A., Delgado, M., Gibbs, M., *et al.* (2012) Costs of dispersal. *Biological Reviews*, **87**, 290–312.
- Bonte, D. & Saastamoinen, M. (2012) Dispersal syndromes in butterflies and spiders. In *Dispersal ecology and evolution* (ed. by Clobert, J., Baguette, M., Benton, T.G. & Bullock, J.M.). Oxford University Press, Oxford, pp. 161–170.
- Bonte, D., Travis, J.M.J., Clercq, N. De, Zwertvaegher, I. & Lens, L. (2008) Thermal conditions during juvenile development affect adult dispersal in a spider. *Proceedings of the National Academy of Sciences of the United States of America*, **105**, 17000–17005.
- Bowler, D.E. & Benton, T.G. (2005) Causes and consequences of animal dispersal strategies: relating individual behaviour to spatial dynamics. *Biological Reviews*, **80**, 205–225.
- Bradshaw, A.D. (1965) Evolutionary significance of phenotypic plasticity in plants. *Advances in Genetics*, **13**, 115–155.
- Brunzel, S., Bussmann, M. & Obergruber, H. (2008) Deutliche Veränderungen von Tagfalterzönosen als Folge von Ausbreitungsprozessen. Erste Ergebnisse eines Monitorings über 17 Jahre. *Natur und Landschaft*, **6**, 280–287.
- Calosi, P., Bilton, D.T. & Spicer, J.I. (2008) Thermal tolerance, acclimatory capacity and vulnerability to global climate change. *Biology Letters*, **4**, 99–102.
- Candy, D.J., Becker, A. & Wegener, G. (1997) Coordination and integration of metabolism in insect flight. *Comparative Biochemistry and Physiology - B Biochemistry and Molecular Biology*, **117**, 497–512.
- Castañeda, L.E., Lardies, M.A. & Bozinovic, F. (2005) Interpopulational variation in recovery time from chill coma along a geographic gradient: a study in the common woodlouse, *Porcellio laevis*. *Journal of Insect Physiology*, **51**, 1346–1351.
- Chown, S.L., Hoffmann, A.A., Kristensen, T.N., Angilletta, M.J., Stenseth, N.C. & Pertoldi, C. (2010) Adapting to climate change: a perspective from evolutionary physiology. *Climate Research*, **43**, 3–15.

- Clark, J.S., Lewis, M., McLachlan, J.S. & HilleRisLambers, J. (2017) Estimating population spread: what can we forecast and how well? *Ecology*, **84**, 1979–1988.
- Clobert, J., Baguette, M., Benton, T.G. & Bullock, J.M. (2012) *Dispersal ecology and evolution*. Oxford University Press, Oxford.
- Clobert, J. & Chaine, A.S. (2012) Dispersal. In *Behavioural responses to a changing world* (ed. by Candolin, U. & Wong, B.B.M.). Oxford University Press, Oxford, pp. 63–79.
- Clobert, J., Galliard, J.F. Le, Cote, J., Meylan, S. & Massot, M. (2009) Informed dispersal, heterogeneity in animal dispersal syndromes and the dynamics of spatially structured populations. *Ecology Letters*, **12**, 197–209.
- Cormont, A., Malinowska, A.H., Kostenko, O., Radchuk, V., Hemerik, L., WallisDeVries, M.F., *et al.* (2011) Effect of local weather on butterfly flight behaviour, movement, and colonization: significance for dispersal under climate change. *Biodiversity and Conservation*, **20**, 483–503.
- Cote, J. & Clobert, J. (2007) Social information and emigration: lessons from immigrants. *Ecology Letters*, **10**, 411–417.
- Cote, J., Clobert, J., Brodin, T., Fogarty, S. & Sih, A. (2010a) Personality-dependent dispersal: characterization, ontogeny and consequences for spatially structured populations. *Philosophical transactions of the Royal Society B*, **365**, 4065–4076.
- Cote, J., Fogarty, S., Weinersmith, K., Brodin, T. & Sih, A. (2010b) Personality traits and dispersal tendency in the invasive mosquitofish (*Gambusia affinis*). *Proceedings of the Royal Society B: Biological Sciences*, **277**, 1571–1579.
- Dall, S.R.X., Giraldeau, L.A., Olsson, O., McNamara, J.M. & Stephens, D.W. (2005) Information and its use by animals in evolutionary ecology. *Trends in Ecology and Evolution*, **20**, 187–193.
- Davis, M.B., Shaw, R.G. & Etterson, J.R. (2005) Evolutionary responses to changing climate. *Ecology*, **86**, 1704–1714.
- Delattre, T., Baguette, M., Burel, F., Stevens, V.M., Quénol, H. & Vernon, P. (2013) Interactive effects of landscape and weather on dispersal. *Oikos*, **122**, 1576–1585.
- Dennis, R.L.H. (1993) *Butterflies and climate change*. Manchester University Press, Manchester.
- Dethier, V.G. & MacArthur, R.H. (1964) A field's capacity to support a butterfly

- population. *Nature*, **201**, 728–729.
- Deutsch, C.A., Tewksbury, J.J., Huey, R.B., Sheldon, K.S., Ghalambor, C.K., Haak, D.C., *et al.* (2008) Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of the National Academy of Sciences of the United States of America*, **105**, 6668–6672.
- DeWitt, T.J., Sih, A. & Wilson, D.S. (1998) Cost and limits of phenotypic plasticity. *Trends in Ecology & Evolution*, **13**, 77–81.
- Dingemanse, N.J., Both, C., Noordwijk, A.J. van, Rutten, A.L. & Drent, P.J. (2003) Natal dispersal and personalities in great tits (*Parus major*). *Proceedings of the Royal Society B: Biological Sciences*, **270**, 741–747.
- Doebeli, M. & Dieckmann, U. (2003) Speciation along environmental gradients. *Nature*, **421**, 259–264.
- Ducatez, S., Humeau, A., Congretel, M., Fréville, H. & Baguette, M. (2014) Butterfly species differing in mobility show different structures of dispersal-related syndromes in the same fragmented landscape. *Ecography*, **37**, 378–389.
- Ducatez, S., Legrand, D., Chaput-Bardy, A., Stevens, V.M., Fréville, H. & Baguette, M. (2012) Inter-individual variation in movement: is there a mobility syndrome in the large white butterfly *Pieris brassicae*? *Ecological Entomology*, **37**, 377–385.
- Duckworth, R.A. & Badyaev, A. V. (2007) Coupling of dispersal and aggression facilitates the rapid range expansion of a passerine bird. *Proceedings of the National Academy of Sciences*, **104**, 15017–15022.
- Dudley, R. (2000) *The biomechanics of insect flight*. Princeton University Press, Princeton.
- Ebert, G. & Rennwald, E. (1991) *Die Schmetterlinge Baden-Württembergs*. Vol. 2. Ulmer, Stuttgart.
- Enfjäll, K. & Leimar, O. (2005) Density-dependent dispersal in the Glanville fritillary, *Melitaea cinxia*. *Oikos*, **108**, 465–472.
- Fischer, K., Beinlich, B. & Plachter, H. (1999) Population structure, mobility and habitat preferences of the violet copper *Lycaena helle* (Lepidoptera: Lycaenidae) in Western Germany: implications for conservation. *Journal of Insect Conservation*, **52**, 43–52.
- Fischer, K. & Fiedler, K. (2000) Sex-related differences in reaction norms in the butterfly *Lycaena tityrus* (Lepidoptera: Lycaenidae). *Oikos*, **90**, 372–380.
- Fischer, K. & Karl, I. (2010) Exploring plastic and genetic responses to temperature

- variation using copper butterflies. *Climate Research*, **43**, 17–30.
- Fjerdingstad, E.J., Schtickzelle, N., Manhes, P., Gutierrez, A. & Clobert, J. (2007) Evolution of dispersal and life history strategies – *Tetrahymena ciliates*. *BMC Evolutionary Biology*, **7**, 133.
- Forester, D.J. & Machlis, G.E. (1996) Modeling human factors that affect the loss of biodiversity. *Conservation Biology*, **10**, 1253–1263.
- Franco, A.M.A., Hill, J.K., Kitschke, C., Collingham, Y.C., Roy, D.B., Fox, R., *et al.* (2006) Impacts of climate warming and habitat loss on extinctions at species' low-latitude range boundaries. *Global Change Biology*, **12**, 1545–1553.
- Franks, S.J. & Hoffmann, A.A. (2012) Genetics of climate change adaptation. *Annual Review of Genetics*, **46**, 185–208.
- Fraser, D.F., Gilliam, J.F., Daley, M.J., Le, A.N. & Skalski, G.T. (2001) Explaining leptokurtic movement distributions: intrapopulation variation in boldness and exploration. *The American Naturalist*, **158**, 124–135.
- Gaston, K.J. (2003) *The structure and dynamics of geographic ranges*. Oxford University Press, Oxford.
- Gilchrist, G.W. (1990) The consequences of sexual dimorphism in body size for butterfly flight and thermoregulation. *Functional Ecology*, **4**, 475–487.
- Hansen, J., Sato, M. & Ruedy, R. (2012) Perception of climate change. *Proceedings of the National Academy of Sciences*, **109**, E2415–E2423.
- Hanski, I. (1994) A practical model of metapopulation dynamics. *Journal of Animal Ecology*, **63**, 151–162.
- Hanski, I. (1998) Metapopulation dynamics. *Nature*, **396**, 41–49.
- Hanski, I. (1999) Habitat connectivity, habitat continuity, and metapopulations in dynamic landscapes. *Oikos*, **87**, 209–219.
- Hanski, I., Breuker, C.J., Schops, K., Setchfield, R. & Nieminen, M. (2002) Population history and life history influence the migration rate of female Glanville fritillary butterflies. *Oikos*, **98**, 87–97.
- Hanski, I., Erälahti, C., Kankare, M., Ovaskainen, O. & Sirén, H. (2004) Variation in migration propensity among individuals maintained by landscape structure. *Ecology Letters*, **7**, 958–966.
- Hanski, I. & Gilpin, M.E. (1997) *Metapopulation biology: ecology, genetics, and evolution*. Academic Press, New York.
- Hassall, C. (2015) Strong geographical variation in wing aspect ratio of a damselfly,

- Calopteryx maculata* (Odonata: Zygoptera). *PeerJ*, **3**, e1219.
- Hastings, A., Cuddington, K., Davies, K.F., Dugaw, C.J., Elmendorf, S., Freestone, A., *et al.* (2005) The spatial spread of invasions: new developments in theory and evidence. *Ecology Letters*, **8**, 91–101.
- Heino, M. & Hanski, I. (2001) Evolution of migration rate in a spatially realistic metapopulation model. *The American Naturalist*, **157**, 495–511.
- Heuvel, J. van den, Saastamoinen, M., Brakefield, P.M., Kirkwood, T.B.L., Zwaan, B.J. & Shanley, D.P. (2013) The predictive adaptive response: modeling the life-history evolution of the butterfly *Bicyclus anynana* in seasonal environments. *The American naturalist*, **181**, E28-E42.
- Hickling, R., Roy, D.B., Hill, J.K., Fox, R. & Thomas, C.D. (2006) The distributions of a wide range of taxonomic groups are expanding polewards. *Global Change Biology*, **12**, 450–455.
- Hill, J.K., Griffiths, H.M. & Thomas, C.D. (2011) Climate change and evolutionary adaptations at species' range margins. *Annual review of entomology*, **56**, 143–159.
- Hill, J.K., Thomas, C.D. & Blakeley, D.S. (1999) Evolution of flight morphology in a butterfly that has recently expanded its geographic range. *Oecologia*, **121**, 165–170.
- Hill, J.K., Thomas, C.D. & Lewis, O.T. (1998) Flight morphology in fragmented populations of a rare British butterfly, *Hesperia comma*. *Biological Conservation*, **87**, 277–283.
- Hoffmann, A.A., Anderson, A. & Hallas, R. (2002) Opposing clines for high and low temperature resistance in *Drosophila melanogaster*. *Ecology letters*, **5**, 614–618.
- Hofmann, G.E. & Todgham, A.E. (2010) Living in the now: physiological mechanisms to tolerate a rapidly changing environment. *Annual Review of Physiology*, **72**, 127–145.
- Hopper, K.R. (1999) Risk-spreading and bet-hedging in insect population biology. *Annual Review of Entomology*, **44**, 535–560.
- Janzen, D.H. (1967) Why mountain passes are higher in the tropics. *The American Naturalist*, **101**, 233-249.
- Jenkins, D.G., Brescacin, C.R., Duxbury, C.V., Elliott, J.A., Evans, J.A., Grablow, K.R., *et al.* (2007) Does size matter for dispersal distance? *Global Ecology and*

- Biogeography*, **16**, 415–425.
- Karl, I. & Fischer, K. (2008) Why get big in the cold? Towards a solution to a life-history puzzle. *Oecologia*, **155**, 215–225.
- Karlsson, B. & Dyck, H. Van. (2005) Does habitat fragmentation affect temperature-related life-history traits? A laboratory test with a woodland butterfly. *Proceedings of the Royal Society B: Biological Sciences*, **272**, 1257–1263.
- Kellermann, V., Overgaard, J., Hoffmann, A.A., Flojgaard, C., Svenning, J.-C. & Loeschcke, V. (2012) Upper thermal limits of *Drosophila* are linked to species distributions and strongly constrained phylogenetically. *Proceedings of the National Academy of Sciences*, **109**, 16228–16233.
- Kingsolver, J.G. (1983) Ecological significance of flight activity in *Colias* butterflies: implications for reproductive strategy and population structure. *Ecology*, **64**, 546–551.
- Klockmann, M., Karajoli, F., Kuczyk, J., Reimer, S. & Fischer, K. (2016) Fitness implications of simulated climate change in three species of copper butterflies (Lepidoptera: Lycaenidae). *Biological Journal of the Linnean Society*, DOI: 10.1111/bij.12846.
- Kolligs, D. (2014) *Schmetterlinge Norddeutschlands*. Wachholtz Verlag, Neumünster/Hamburg.
- Kuussaari, M., Rytteri, S., Heikkinen, R.K., Helio, J. & Bagh, P. von. (2016) Weather explains high annual variation in butterfly dispersal. *Proceedings of the Royal Society B*, **283**, 20160413.
- Lebeau, J., Wesselingh, R.A. & Dyck, H. Van. (2016) Floral resource limitation severely reduces butterfly survival, condition and flight activity in simplified agricultural landscapes. *Oecologia*, **180**, 421–427.
- Legrand, D., Guillaume, O., Baguette, M., Cote, J., Trochet, A., Calvez, O., *et al.* (2012) The Metatron: an experimental system to study dispersal and metaecosystems for terrestrial organisms. *Nature Methods*, **9**, 828–833.
- Legrand, D., Larranaga, N., Bertrand, R., Ducatez, S., Calvez, O., Stevens, V.M., *et al.* (2016) Evolution of a butterfly dispersal syndrome. *Proceedings of the Royal Society B: Biological Sciences*, **283**, 20161533.
- Legrand, D., Trochet, A., Moulherat, S., Calvez, O., Stevens, V.M., Ducatez, S., *et al.* (2015) Ranking the ecological causes of dispersal in a butterfly. *Ecography*, **38**, 822–831.

- Lindstrom, T., Brown, G.P., Sisson, S.A., Phillips, B.L. & Shine, R. (2013) Rapid shifts in dispersal behavior on an expanding range edge. *Proceedings of the National Academy of Sciences*, **110**, 13452–13456.
- MacArthur, R.H. (1972) *Geographical ecology: patterns in the distribution of species*. Princeton University Press, Princeton.
- Mägi, M., Semchenko, M., Kalamees, R. & Zobel, K. (2011) Limited phenotypic plasticity in range-edge populations: a comparison of co-occurring populations of two *Agrimonia* species with different geographical distributions. *Plant Biology*, **13**, 177–184.
- Matthysen, E. (2012) Multicausality of dispersal: a review. In *Dispersal ecology and evolution* (ed. by Clobert, J. et al.). Oxford University Press, Oxford, pp. 1–18.
- Mattila, A.L.K. & Hanski, I. (2014) Heritability of flight and resting metabolic rates in the Glanville fritillary butterfly. *Journal of Evolutionary Biology*, **27**, 1733–1743.
- McNamara, J.M. & Houston, A.I. (1996) State-dependent life histories. *Nature*, **380**, 215–221.
- Merckx, T. & Dyck, H. Van. (2005) Mate location behaviour of the butterfly *Pararge aegeria* in woodland and fragmented landscapes. *Animal Behaviour*, **70**, 411–416.
- Merilä, J. (2012) Evolution in response to climate change: in pursuit of the missing evidence. *BioEssays*, **34**, 811–818.
- Mitikka, V. & Hanski, I. (2010) Pgi genotype influences flight metabolism at the expanding range margin of the European map butterfly. *Annales Zoologici Fennici*, **47**, 1–14.
- Moles, A.T. & Westoby, M. (2014) Seed size and plant strategy across the whole life cycle. *Oikos*, **113**, 91–105.
- Monaghan, P. (2008) Early growth conditions, phenotypic development and environmental change. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, **363**, 1635–1645.
- Murrell, D.J., Travis, J.M.J. & Dytham, C. (2002) The evolution of dispersal distance in spatially-structured populations. *Oikos*, **97**, 229–236.
- Naya, D.E., Spangenberg, L., Naya, H. & Bozinovic, F. (2012) Latitudinal patterns in rodent metabolic flexibility. *The American Naturalist*, **179**, E172–E179.
- Niitepõld, K., Smith, A.D., Osborne, J.L., Reynolds, D.R., Carreck, N.L., Martin, A.P., et al. (2009) Flight metabolic rate and Pgi genotype influence butterfly dispersal

- rate in the field. *Ecology*, **90**, 2223–2232.
- Öckinger, E. & Smith, H.G. (2008) Do corridors promote dispersal in grassland butterflies and other insects? *Landscape Ecology*, **23**, 27–40.
- Õunap, E. & Tartes, U. (2014) *Eesti päevaliblikad [Estonian butterflies]*. Varrak, Tallinn.
- Overgaard, J. & Sørensen, J.G. (2008) Rapid thermal adaptation during field temperature variations in *Drosophila melanogaster*. *Cryobiology*, **56**, 159–162.
- Parmesan, C. (2006) Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology, Evolution, and Systematics*, **37**, 637–669.
- Parmesan, C., Ryrholm, N., Stefanescu, C., Hill, J.K., Thomas, C.D., Descimon, H., et al. (1999) Poleward shifts in geographical ranges of butterfly species associated with regional warming. *Nature*, **399**, 579–583.
- Parmesan, C. & Yohe, G. (2003) A globally coherent fingerprint of climate change impacts across natural systems. *Nature*, **421**, 37–42.
- Phillips, B.L., Brown, G.P., Webb, J.K. & Shine, R. (2006) Invasion and the evolution of speed in toads. *Nature*, **439**, 803.
- Phillips, B.L., Brown, G.P. & Shine, R. (2010) Life-history evolution in range-shifting populations. *Ecology*, **91**, 1617–1627.
- Pigliucci, M. (2005) Evolution of phenotypic plasticity: where are we going now? *Trends in Ecology and Evolution*, **20**, 481–486.
- Pimm, S.L. (2001) Entrepreneurial insects. *Nature*, **411**, 531–532.
- Pleasants, J.M. & Oberhauser, K.S. (2013) Milkweed loss in agricultural fields because of herbicide use: effect on the monarch butterfly population. *Insect Conservation and Diversity*, **6**, 135–144.
- Purves, D.W. (2009) The demography of range boundaries versus range cores in eastern US tree species. *Proceedings of the Royal Society B: Biological Sciences*, **276**, 1477–1484.
- Rahmstorf, S. & Coumou, D. (2011) Increase of extreme events in a warming world. *Proceedings of the National Academy of Sciences*, **108**, 17905–17909.
- Roff, D.A. & Fairbairn, D.J. (2001) The genetic basis of dispersal and migration, and its consequences for the evolution of correlated traits. In *Dispersal* (ed. by Clobert, J., Danchin, E., Dhondt, A.A. & Nichols, J.D.). Oxford University Press, Oxford, pp. 191–202.
- Roff, D.A. & Fairbairn, D.J. (2007) The evolution and genetics of migration in insects. *BioScience*, **57**, 155–164.

- Ronce, O. (2007) How does it feel to be like a rolling stone? Ten questions about dispersal evolution. *Annual Review of Ecology, Evolution, and Systematics*, **38**, 231–253.
- Saastamoinen, M., Brakefield, P.M. & Ovaskainen, O. (2012) Environmentally induced dispersal-related life-history syndrome in the tropical butterfly, *Bicyclus anynana*. *Journal of Evolutionary Biology*, **25**, 2264–2275.
- Saastamoinen, M., Sterren, D. van der, Vastenhout, N., Zwaan, B.J. & Brakefield, P.M. (2010) Predictive adaptive responses: condition-dependent impact of adult nutrition and flight in the tropical butterfly *Bicyclus anynana*. *The American naturalist*, **176**, 686–698.
- Sanderson, E.W., Jaiteh, M., Levy, M.A., Redford, K.H., Wannebo, A. V. & Woolmer, G. (2002) The human footprint and the last of the wild. *BioScience*, **52**, 891–904.
- Schtickzelle, N., Mennechez, G. & Baguette, M. (2006) Dispersal depression with habitat fragmentation in the bog fritillary butterfly. *Ecology*, **87**, 1057–1065.
- Sekar, S. (2012) A meta-analysis of the traits affecting dispersal ability in butterflies: can wingspan be used as a proxy? *Journal of Animal Ecology*, **81**, 174–184.
- Settele, J., Kudrna, O., Harpke, A., Kühn, I. & Swaay, Van, C. (2008) *Climatic risk atlas of European butterflies. Biodiversity and ecosystem risk assessment 1*. Pensoft, Sofia.
- Settele, J., Shreeve, T.G., Konvička, M. & Van Dyck, H. (2009) *Ecology of butterflies in Europe*. Cambridge University Press, Cambridge.
- Shine, R., Brown, G.P. & Phillips, B.L. (2011) An evolutionary process that assembles phenotypes through space rather than through time. *Proceedings of the National Academy of Sciences*, **108**, 5708–5711.
- Stanton, J.C., Shoemaker, K.T., Pearson, R.G. & Akçakaya, H.R. (2015) Warning times for species extinctions due to climate change. *Global Change Biology*, **21**, 1066–1077.
- Stevens, V.M., Trochet, A., Blanchet, S., Moulherat, S., Clobert, J. & Baguette, M. (2013) Dispersal syndromes and the use of life-histories to predict dispersal. *Evolutionary Applications*, **6**, 630–642.
- Stevens, V.M., Turlure, C. & Baguette, M. (2010) A meta-analysis of dispersal in butterflies. *Biological Reviews*, **85**, 625–462.
- Suarez, R.K. (2000) Energy metabolism during insect flight: biochemical design and

- physiological performance. *Physiological and Biochemical Zoology*, **73**, 765–771.
- Sunday, J.M., Bates, A.E. & Dulvy, N.K. (2011) Global analysis of thermal tolerance and latitude in ectotherms. *Proceedings of the Royal Society B: Biological Sciences*, **278**, 1823–1830.
- Therry, L., Zawal, A., Bonte, D. & Stoks, R. (2014) What factors shape female phenotypes of a poleward-moving damselfly at the edge of its range? *Biological Journal of the Linnean Society*, **112**, 556–568.
- Thomas, C.D. (2010) Climate, climate change and range boundaries. *Diversity and Distributions*, **16**, 488–495.
- Thomas, C.D., Bodsworth, E.J., Wilson, R.J., Simmons, A.D., Davies, Z.G., Musche, M., *et al.* (2001) Ecological and evolutionary processes at expanding range margins. *Nature*, **411**, 577–581.
- Thomas, C.D., Franco, A.M.A. & Hill, J.K. (2006) Range retractions and extinction in the face of climate warming. *Trends in Ecology and Evolution*, **21**, 415–416.
- Tolman, T. & Lewington, R. (1998) *Die Tagfalter Europas und Nordwestafrikas*. Franckh-Kosmos, Stuttgart.
- Travis, J.M.J., Delgado, M., Bocedi, G., Baguette, M., Bartoń, K., Bonte, D., *et al.* (2013) Dispersal and species' responses to climate change. *Oikos*, **122**, 1532–1540.
- Travis, J.M.J. & French, D.R. (2000) Dispersal functions and spatial models: expanding our dispersal toolbox. *Ecology Letters*, **3**, 163–165.
- Trochet, A., Legrand, D., Larranaga, N., Ducatez, S., Calvez, O., Cote, J., *et al.* (2013) Population sex ratio and dispersal in experimental, two-patch metapopulations of butterflies. *Journal of Animal Ecology*, **82**, 946–955.
- Turlure, C., Schtickzelle, N., Dyck, H. Van, Seymoure, B. & Rutowski, R. (2016) Flight morphology, compound eye structure and dispersal in the bog and the cranberry fritillary butterflies: an inter- and intraspecific comparison. *PLOS ONE*, **11**, e0158073.
- Van Dyck, H. & Baguette, M. (2005) Dispersal behaviour in fragmented landscapes: routine or special movements? *Basic and Applied Ecology*, **6**, 535–545.
- Van Dyck, H. & Wiklund, C. (2002) Seasonal butterfly design: morphological plasticity among three developmental pathways relative to sex, flight and thermoregulation. *Journal of Evolutionary Biology*, **15**, 216–225.

- Van Petegem, K.H.P., Boeye, J., Stoks, R. & Bonte, D. (2016) Spatial selection and local adaptation jointly shape life-history evolution during range expansion. *The American Naturalist*, **188**, 485–498.
- Volis, S., Mendlinger, S., Olsvig-Whittaker, L., Safriel, U.N. & Orlovsky, N. (1998) Phenotypic variation and stress resistance in core and peripheral populations of *Hordeum spontaneum*. *Biodiversity and Conservation*, **7**, 799–813.
- Warren, M.S., Hill, J.K., Thomas, J.A., Asher, J., Fox, R., Huntley, B., *et al.* (2001) Rapid responses of British butterflies to opposing forces of climate and habitat change. *Nature*, **414**, 65–69.
- Watt, W.B. (1995) Adaptive significance of pigment polymorphisms in *Colias* butterflies. I. Variation of melanin pigment in relation to thermoregulation. *Evolution*, **22**, 437–458.
- Wickman, P.O. (2009) Thermoregulation and habitat use in butterflies. In *Ecology of butterflies in Europe* (ed. by Settele, J., Shreeve, T.G., Konvicka, M. & Dyck, H. Van). Cambridge University Press, Cambridge, pp. 55–61.
- Williams, S.E., Shoo, L.P., Isaac, J.L., Hoffmann, A.A. & Langham, G. (2008) Towards an integrated framework for assessing the vulnerability of species to climate change. *PLoS Biology*, **6**, e325.
- Wingfield, J.C., Krause, J.S., Perez, J.H., Chmura, H.E., Németh, Z., Word, K.R., *et al.* (2015) A mechanistic approach to understanding range shifts in a changing world: what makes a pioneer? *General and Comparative Endocrinology*, **222**, 44–53.
- Wolf, J.B., Brodie, E.D., Cheverud, J.M., Moore, A.J. & Wade, M.J. (1998) Evolutionary consequences of indirect genetic effects. *Trends in Ecology and Evolution*, **13**, 64–69.
- Zera, A.J., Potts, J. & Kobus, K. (1998) The physiology of life-history trade-offs: experimental analysis of a hormonally induced life-history trade-off in *Gryllus assimilis*. *The American Naturalist*, **152**, 7–23.

# **PUBLICATION LIST**



## 4 Publication list

- **Reim, E.**, Arnstedt, I., Barwisch, I., Baumgarten, M., Bock, S., Eberspach, J., Ellerbrok, J., Gebremeskel, M., Küpper, S., Guth, L., Lassen, A., Letro, L., Meth, R., Möller, M., Närmann, F., Neunaber, I., Seliger, A., v. Stein, W., Vallinga, C., Vögele, P., Wagner, M., Fischer, K. (2018) Movement patterns differ between sexes and depend on weather conditions in the butterfly *Lycaena tityrus*. Journal of Insect Behavior, 31(3), 309-320.

Contribution to publication:

**All authors** conceived the ideas and designed methodology; IA, IB, MB, SB, JE, JE, MG, SK, LG, AL, LL, RM, MM, FN, IN, AS, WS, CV, PV and MW collected the data; **ER** analyzed the data; IA, IB, MB, SB, JE, JE, MG, SK, LG, AL, LL, RM, MM, FN, IN, AS, WS, CV, PV and MW wrote the first draft of the manuscript. **ER** and KF revised and edited the manuscript.

- **Reim, E.**, Baguette, M., Günter, F., Fischer, K. Emigration propensity and flight performance are decoupled in a butterfly. In revision.

Contribution to publication:

**ER** and KF conceived the ideas and designed methodology; **ER** and FG collected the data; **ER** analyzed the data with input from MB and KF; **ER** led and KF and MB contributed to the writing of the manuscript.

- **Reim, E.**, Eichhorn, D., Roy, J.D., Steinhoff, P.O.M., Fischer, K. (2018) Nutritional stress reduces flight performance and exploratory behavior in a butterfly. Insect Science, DOI: 10.1111/1744-7917.12596.

Contribution to publication:

**ER** and KF conceived the ideas and designed methodology; **ER**, DE, JR and PS collected the data; **ER** analyzed the data with input from KF; **ER** led and KF contributed to the writing of the manuscript.

- **Reim, E.**, Blesinger, S., Förster, L., Fischer, K. Successful despite poor flight performance: range expansion is associated with enhanced exploratory behavior and fast development. *Journal of Evolutionary Biology*, in press.

Contribution to publication:

**ER** and KF conceived the ideas and designed methodology; **ER**, SB and LF collected the data; **ER** analyzed the data with input from KF; **ER** led and KF contributed to the writing of the manuscript.

#### 4.1 Movement patterns differ between sexes and depend on weather conditions in the butterfly *Lycaena tityrus*

Published as:

Reim, E., Arnstedt, I., Barwisch, I., Baumgarten, M., Bock, S., Eberspach, J., Ellerbrok, J., Gebremeskel, M., Küpper, S., Guth, L., Lassen, A., Letro, L., Meth, R., Möller, M., Närmann, F., Neunaber, I., Seliger, A., v. Stein, W., Vallinga, C., Vögele, P., Wagner, M., Fischer, K. (2018) Movement patterns differ between sexes and depend on weather conditions in the butterfly *Lycaena tityrus*. *Journal of Insect Behavior*, 31(3), 309-320.



## Movement Patterns Differ between Sexes and Depend on Weather Conditions in the Butterfly *Lycaena tityrus*

Elisabeth Reim<sup>1</sup>  · Ingo Arnstedt<sup>1</sup> · Isabel Barwisch<sup>1</sup> · Max Baumgarten<sup>1</sup> · Sascha Bock<sup>1</sup> · Julia Eberspach<sup>1</sup> · Julia Ellerbrok<sup>1</sup> · Mulugeta Gebremeskel<sup>1</sup> · Simon Küpper<sup>1</sup> · Lukas Guth<sup>1</sup> · Aninha Lassen<sup>1</sup> · Letro Letro<sup>1</sup> · Rebecca Meth<sup>1</sup> · Maria Möller<sup>1</sup> · Felix Närmann<sup>1</sup> · Inga Neunaber<sup>1</sup> · Alexander Seliger<sup>1</sup> · Wilderich v. Stein<sup>1</sup> · Carolin Vallinga<sup>1</sup> · Philipp Vögele<sup>1</sup> · Martin Wagner<sup>1</sup> · Klaus Fischer<sup>1</sup>

Revised: 10 April 2018 / Accepted: 11 April 2018

© Springer Science+Business Media, LLC, part of Springer Nature 2018

**Abstract** Habitat loss and anthropogenic climate change are important threats to biodiversity conservation. Owing to the concomitantly deteriorating habitat quality, individuals are often forced to disperse to new habitats, rendering dispersal an ecologically important process. However, dispersal ability may differ within and among populations, and is further dependent on environmental conditions. We therefore studied sexual differences in and environmental effects on movement patterns in the sooty copper butterfly *Lycaena tityrus*. As predicted, males were more active and covered longer distances than females, presumably owing to mate location and territorial disputes. Males alighted more often on flowers than females, probably to fuel their high flight activity, while females showed a high affinity to host-plants for egg-laying. Our findings provide a striking example of sex-related differences in animal behavior, as revealed by the use of customary smartphones, which apparently can comprise suitable means to reveal biologically significant behavioural patterns. More problematic than the technical device used seems to be the challenge of following individual butterflies for long enough in the field, such that any extrapolations to dispersal seem difficult.

**Keywords** Animal behavior · dispersal · environmental context · flight activity · mate location · sexual differences

---

**Electronic supplementary material** The online version of this article (<https://doi.org/10.1007/s10905-018-9679-8>) contains supplementary material, which is available to authorized users.

---

✉ Elisabeth Reim  
elisabeth.reim@uni-greifswald.de

<sup>1</sup> Zoological Institute and Museum, Greifswald University, Soldmannstraße 14, D-17489 Greifswald, Germany

## Introduction

Human impact, driving habitat loss and fragmentation as well as climate change, is all pervasive on Earth (e.g. Forester and Machlist 1996; Sanderson et al. 2002). Owing to the concomitantly deteriorating habitat quality, individuals are often forced to disperse to new habitats (e.g. Hanski 1998), although the likelihood to reach these is often low due to reduced habitat availability and fragmentation (Fernández-Chacón et al. 2014). Dispersal is therefore important for dealing with environmental variation (Cote and Clobert 2007; Travis et al. 2013), as it may allow individuals to track their shifting climate niche (Warren et al. 2001; Hickling et al. 2006) and to maintain metapopulation connectivity (Hanski and Gilpin 1997; Baguette et al. 2013).

Despite its concomitant ecological importance, the motivation as well as the ability to disperse may differ strongly among and within populations (Van Dyck and Baguette 2005; Baguette and Van Dyck 2007; Ducatez et al. 2014; Bestion et al. 2015). Factors that may affect dispersal ability include morphology, physiology, and overall condition (Bowler and Benton 2005; Bonte et al. 2012; Therry et al. 2014). Moreover, sexes may differ in dispersal behavior. In insects, females are often the more dispersive sex (e.g. Fischer et al. 1999) as (1) single-mated females are, unlike males, able to found a new population (Hill et al. 1999; Hanski et al. 2002, 2004), (2) females may need to escape from male harassment (Hovestadt and Nieminen 2009; Trochet et al. 2013) or as (3) they pursue a risk spreading strategy by distributing their eggs over a wider area (Hopper 1999). In addition to such intrinsic factors, dispersal is also affected by external factors such as prevailing weather conditions (Legrand et al. 2015). In general, dispersal increases with rising temperature and solar radiation but decreases with increasing cloud cover, rainfall, and wind speed (Kuussaari et al. 2016).

Against this background, we here investigate (within-patch) movement patterns in the temperate-zone butterfly *Lycaena tityrus*. Focussing on movement patterns rather than dispersal per se was due to the fact that dispersal is very difficult if not impossible to observe directly in flying insects (Riley and Smith 2002; Cant et al. 2005). We thus argue that knowledge on movement patterns is an important first step for a better understanding of dispersal, as movement may eventually result in dispersal. For instance, flight track length has been shown to be associated with flight metabolic rate and PGI genotype, parameters that are in turn known to be related to dispersal (Niitepõld et al. 2009). Butterflies are suitable models to study movement for several reasons. First, they exhibit large variation in dispersal ability and their natural history is well known (Stevens et al. 2010). Second, they are sensitive indicators of habitat change and therefore ecosystem quality (Murphy et al. 1990; van Swaay 1990; Erhardt and Thomas 1991). Third, as pollinators they play an important role in ecosystems by providing pollination services, the lack of which may cause a loss of plant diversity (Potts et al. 2010). Fourth, as herbivores they may be particularly threatened by climate change, due to effects on the quality and phenology of host-plants as well as on the butterflies themselves (Roy and Sparks 2000).

Specifically, we explore within-patch mobility to test the hypotheses that (1) movement patterns differ among the sexes, and that (2) movement is strongly affected by prevailing weather conditions. As *L. tityrus* males are territorial (Ebert and Rennwald 1991; Fischer and Fiedler 2000a), we predict that males are more active than females (due to mate location, inspection flights, territorial disputes), while females are

expected to move further away from the point of first encounter than males due to straighter flight trajectories (see above). We further predict that activity is positively related to beneficial (warm, sunny) weather conditions. Addressing such questions may have important implications for dispersal, though note that we are working here on a spatially very restricted scale. To this end, we individually tracked butterflies of both sexes within an extended habitat patch and recorded weather conditions in parallel. As we used customary smartphones in combination with a tracking application, we additionally investigate whether such devices are suitable for investigating insect behaviour.

## Material and Methods

### Study Organism

The sooty copper *L. tityrus* is a widespread butterfly of the temperate zone, ranging from Western Europe to Central Asia (Ebert and Rennwald 1991). Central European low-altitude populations, including the one investigated here, are typically bivoltine with two distinct generations per year. *L. tityrus* inhabits different types of unimproved grass- and wetlands. The most important larval host plant is *Rumex acetosa*, but several congeneric plant species such as *R. acetosella* and *R. scutatus* are utilised as well (Ebert and Rennwald 1991; Tolman and Lewington 1998). Adults feed on a variety of composite (Compositae) and other nectar plants (Ebert and Rennwald 1991; Karl and Fischer 2009). For this study, individuals from a low-altitude population near the city of Greifswald (north-eastern Germany) were examined.

### Study Area

The study was conducted on an extended set-aside field near Greifswald (54°03'02"N, 13°26'26"E), harboring a large population of *L. tityrus*. Hence, the location is close to the Baltic Sea and thus the northern distribution limit of *L. tityrus* (GfS 2017). This part of Germany is characterized by a rather continental climate with an annual mean temperature of 8.0 °C and a precipitation of 566 mm per year (Müller 2006). The study area (ca. 20 ha) is situated 30–40 m above sea level, rather flat, and shows a relatively homogenous vegetation structure without shrubs or trees. The vegetation consists mainly of species indicative of rather nutrient-poor grassland. Host plants (*R. acetosa*, *R. acetosella*) occur in high density throughout the area. The lack of barriers and other disturbing structure renders this study area suitable for tracking individual butterflies.

### Field Methods and Data Analyses

We investigated the movement patterns of *L. tityrus* by tracking the flight paths of individual butterflies with a hand-held GPS device. As GPS device we used Android smartphones and the application “Geo Tracker – GPS tracker”. The following settings were used: accuracy 10 to 20 m, frequency of locating 1 per sec, minimal distance between two points 1 m, maximum distance between two points 500 m. To test for the

accuracy of the GPS application and the smartphones used, tracks of a known distance were compared to measured GPS tracks (Online Resource Fig. 1), revealing a correlation coefficient of 0.995. Furthermore, repeatedly scoring the same position using Geo Tracker without moving revealed identical coordinates. Field work was conducted between mid-May and early June 2017, i.e. within the spring flight period of *L. tityrus*. Data were collected by 12 teams consisting of two persons each, which were consistent over the study period. One person concentrated on observing and tracking the butterfly while the other noted butterfly behavior, time periods, etc. (see further below).

To start a track, an individual of *L. tityrus* was located and its sex determined. Thereafter, the observation was started (to ease sex determination and avoid chasing butterflies, all tracks were started when butterflies were sitting). The observers kept a minimum distance of 2–3 m to the butterfly to minimize possible influences on the animal's behavior. Nevertheless, the actual flight path was tracked as closely as possible by avoiding any shortcuts. Tracking of a butterfly ended when the observer lost sight of the individual or after a maximum of 15 min (i.e. tracking was terminated in case an individual was still available after 15 min). Butterflies showing no flight activity within 15 min were omitted from further analyses.

The following parameters were recorded during tracking: date, sex, cloud cover on a scale between 1 and 8 (in n/8; thus 1/8 equals a cloud cover of one eighth of the sky and 7/8 cloudless), start and end time of tracking, the number of stops, the substrate at the place of alighting (either bare ground, host plant, nectar flower, or other vegetation). Thus, stops separated different flight steps, while crawling within the vegetation was not considered as movement. In total 196 tracks were taken, 106 from males and 90 from females, with each of the 12 observer teams contributing  $\geq 10$  tracks. The tracking application was used to extract the following data: time in motion, total track length, average movement speed, average movement speed in motion, and maximum movement speed. To assess whether a high activity may facilitate dispersal, we additionally measured the linear distance between the starting and end point of each track in Google Earth by exporting the tracks to a computer device. Step length was calculated as total track length divided by the number of stops. Data on temperature, wind speed, relative humidity, and precipitation during each individual track were obtained from the Deutscher Wetterdienst (DWD 2017) and are based on hourly means of the weather station closest to the study area (station Greifswald, distance 5.5 km, coordinates: 54°05'39"N, 13°24'20"E).

## Statistical Analyses

Considering the high density of *L. tityrus* and the large size of the study area, we deem it unlikely that a single individual was observed more than once, although we did not mark the individuals. Therefore, we consider our observations as independent samples representative of the butterfly behavior in this population. To analyze movement-related traits we used general linear mixed models with sex as fixed effect, observer team as random effect, and the covariates tracking time, cloud cover and temperature. Due to correlations with temperature, wind speed and relative humidity were excluded from further analyses. Additionally, precipitation was not considered as it showed essentially no variation, because field work was not carried out under rainy conditions (Online Resource Table 1 and Fig. 2). All statistical tests were performed with Statistica 12.0 (Tulsa, StatSoft, OK). All means are given  $\pm 1$  SE.

## Results

Significant sex differences were found in track length, linear distance between start and end point, time in motion, average speed, average movement speed, and the number of stops on flowers and on host-plants, but not in step length, maximum speed, and the number of stops in total, on other vegetation or on bare ground (Table 1). On average, males compared with females exhibited longer track lengths (Fig. 1a) and distances between start and end point ( $22.3 \pm 2.5$  m vs.  $14.5 \pm 1.6$  m), higher average speeds over the total observation period ( $0.78 \pm 0.08$  km/h vs.  $0.22 \pm 0.03$  km/h) and also in motion (Fig. 1b), spent more time flying (Fig. 1c), and stopped more often on flowers (Fig. 1d) but less often on host-plants (Fig. 1e).

Regarding effects of environmental factors, only linear distance between start and end point, step length, and the number of stops on host-plants were significantly associated with cloud cover, indicating that increasing cloud cover increased distances (beta  $0.17 \pm 0.09$ , Online Resource Fig. 3) and step lengths (beta  $0.18 \pm 0.09$ ) but decreased the number of stops on host-plants (beta  $-0.17 \pm 0.08$ ). Temperature exclusively affected the number of stops on bare ground, which increased with decreasing temperature (beta  $-0.36 \pm 0.11$ ). Track length (beta  $0.17 \pm 0.08$ ), the total number of stops (beta  $0.44 \pm 0.07$ ) and the number of stops on flowers (beta  $0.23 \pm 0.08$ ), host-plants (beta  $0.22 \pm 0.08$ ) and other vegetation (beta  $0.32 \pm 0.08$ ) were all significantly positively related, whereas step length (beta  $-0.19 \pm 0.09$ ) and average speed (beta  $-0.36 \pm 0.07$ ) were negatively related to tracking time, while no significant association was found with any of the other dependent variables. Significant effects of observer team were found in all dependent variables except from step length and the number of stops on bare ground.

## Discussion

Our study revealed, as predicted, sex-specific differences in behavioral patterns of *L. tityrus*. Males spent more time flying and showed longer track lengths than females, in accordance with our expectation. We assume that the longer track lengths result from the males' higher overall activity (see also Kingsolver 1983). The males' higher activity is most likely due to spending much of their time locating mates including frequent inspection and patrolling flights as well as territorial disputes with rivaling males (Wickman 1985; Ebert and Rennwald 1991; Fischer et al. 1999). Females, in contrast, seem to generally spend less time flying and allocate more of their time to oviposition (Kingsolver 1983; Fischer et al. 1999). However, the fact that males also covered longer linear distances between the start and end points of observations is contrary to our expectation. For various reasons we expected females to cover longer distances owing to straighter flight trajectories, while males were supposed to stay within or at least close to their territories. Furthermore, the current data challenge earlier ones obtained from mark-recapture studies in related species, where males were found to be more stationary than females (Fischer et al. 1999; Fischer and Fiedler 2000b). We believe that our current results are biased due to the rather short observation periods (max. 15 min) compared to the mark-recapture studies in which butterflies were observed for several days. Thus, within this rather short period of time, the males'

**Table 1** General linear mixed models for the effects of sex (fixed), observer team (random), and the covariates tracking time, cloud cover and temperature on track length, linear distance between start and end point of track, step length, time in motion, maximum speed, average speed, average movement speed, total number of stops, the number of stops on flowers, the number of stops on host-plants (*Rumex spec.*), the number of stops on other vegetation, and the number of stops on bare ground in the butterfly *Lycaena tityrus*. Significant *p*-values are given in bold

Track length	DF	MS	F	p
Sex	1	55324	29.9	< <b>0.0001</b>
Observer team	11	7336	4.0	< <b>0.0001</b>
Track time	1	9653	5.2	<b>0.0236</b>
Cloud Cover	1	1943	1.5	0.3071
Temperature	1	205	0.1	0.7397
Error	179	1852		
Linear distance	DF	MS	F	p
Sex	1	2443	5.8	<b>0.0173</b>
Observer team	11	929	2.2	<b>0.0165</b>
Track time	1	6	< 0.1	0.9077
Cloud cover	1	1677	4.0	<b>0.0480</b>
Temperature	1	549	1.3	0.2560
Error	177	423		
Step length	DF	MS	F	p
Sex	1	466	1.0	0.3170
Observer team	11	762	1.6	0.0890
Track time	1	2235	4.8	<b>0.0292</b>
Cloud cover	1	1852	4.0	<b>0.0469</b>
Temperature	1	53	0.1	0.7364
Error	179	462		
Time in motion	DF	MS	F	p
Sex	1	145652	26.3	< <b>0.0001</b>
Observer team	11	18491	3.3	<b>0.0003</b>
Track time	1	8850	1.6	0.2077
Cloud cover	1	9056	1.6	0.2025
Temperature	1	11899	2.1	0.1444
Error	179	5536		
Maximum speed	DF	MS	F	p
Sex	1	20.8	2.6	0.1112
Observer team	11	27.8	3.4	<b>0.0002</b>
Track time	1	23.1	2.8	0.0936
Cloud cover	1	10.4	1.3	0.2595
Temperature	1	0.1	< 0.1	0.9255
Error	179	8.1		
Average speed	DF	MS	F	p
Sex	1	4.5	14.1	<b>0.0002</b>

**Table 1** (continued)


Observer team	11	1.5	4.7	<b>&lt; 0.0001</b>
Track time	1	8.0	4.7	<b>&lt; 0.0001</b>
Cloud cover	1	0.3	0.9	0.3342
Temperature	1	< 0.1	< 0.1	0.8840
Error	179	0.3		
Average movement speed	DF	MS	F	p
Sex	1	39.6	6.5	<b>0.0115</b>
Observer team	11	25.2	4.2	<b>0.0002</b>
Track time	1	9.6	1.6	0.2104
Cloud cover	1	15.4	2.5	0.1129
Temperature	1	7.5	1.2	0.2687
Error	179	6.1		
Stops number	DF	MS	F	p
Sex	1	13.8	0.7	0.3985
Observer team	11	64.1	3.3	<b>0.0004</b>
Track time	1	0.6	34.6	<b>&lt; 0.0001</b>
Cloud cover	1	41.9	2.2	0.1423
Temperature	1	32.6	1.7	0.1954
Error	179	19.3		
Flower stops	DF	MS	F	p
Sex	1	70.5	8.7	<b>0.0035</b>
Observer team	11	15.3	1.9	<b>0.0420</b>
Track time	1	59.7	7.4	<b>0.0071</b>
Cloud cover	1	0.4	< 0.1	0.8276
Temperature	1	12.6	1.6	0.2124
Error	179	423.0		
<i>Rumex</i> stops	DF	MS	F	p
Sex	1	62.4	17.6	<b>&lt; 0.0001</b>
Observer team	11	8.2	2.3	<b>0.0115</b>
Track time	1	18.8	8.1	<b>0.0049</b>
Cloud cover	1	17.0	4.8	<b>0.0300</b>
Temperature	1	9.5	2.7	0.1037
Error	179	3.5		
Vegetation stops	DF	MS	F	p
Sex	1	9.0	0.9	0.3382
Observer team	11	36.4	3.7	<b>&lt; 0.0001</b>
Track time	1	157.1	16.0	<b>&lt; 0.0001</b>

**Table 1** (continued)

Cloud cover	1	3.1	0.3	0.5749
Temperature	1	2.6	0.3	0.6040
Error	179	9.8		
Ground stops	DF	MS	F	p
Sex	1	< 0.1	< 0.1	0.8934
Observer team	11	0.8	1.7	0.0744
Track time	1	< 0.1	< 0.1	0.8314
Cloud cover	1	0.6	1.3	0.2514
Temperature	1	5.1	11.0	<b>0.0011</b>
Error	179	0.5		

higher activity may have caused longer linear distances. Note in this context the short absolute distances covered within the observation period. Additionally, males may have abandoned territoriality. Note that territorial behavior in *Lycaena* butterflies depends on context, and that males may switch to patrolling behavior during spells of beneficial weather conditions and at high densities of conspecifics (Fischer and Fiedler 2001), which was the case during the study period.

The finding that males achieved higher speeds than females has probably morphological and physiological reasons. Male (*Lycaena*) butterflies have a lower body mass and concomitantly wing loading, but a higher thorax-abdomen ratio, wing aspect ratio, and relative fat content compared with females (Karl et al. 2008; Saastamoinen et al. 2012). These traits are typically associated with increased flight ability and maneuverability (van Dyck et al. 1998; van Dyck and Wiklund 2002; Berwaerts et al. 2008),



**Fig. 1** Sexual differences in *Lycaena tityrus* in track length (a), average movement speed (b), time in motion (c), and the number of stops on flowers (d) and *Rumex* host-plants (e)

likely of crucial importance for males to succeed in territorial contests and competition for mates. The fact that males alighted more often on flowers than females is presumably caused by the need for nectar to fuel their high flight activity, while the higher affinity of females to host-plants is certainly related to the search for oviposition plants (Fischer et al. 1999).

In contrast to the prevalent sexual differences in movement, effects of environmental factors were less pronounced. While cloud cover significantly affected three variables, temperature affected only one. The negative impact of cloud cover on the number of stops on host plants potentially reflects the high dependence of *L. tityrus* activity on solar radiation, being an extremely heliophilous butterfly (Wickman 2009). Thus, a lack of solar radiation may result in a reduced (egg-laying) activity. Typically, females try to spread their eggs over a wide range as a risk-spreading strategy (Hopper 1999). Contrary to our assumptions, the linear distance covered and step length were positively related to cloud cover, for which we have no straightforward explanation. Perhaps air temperature was still high enough to sustain flight activity. The more frequent stops on bare ground at cooler temperatures are probably related to thermoregulatory needs, thus alighting on relatively warm spots compared with the surrounding vegetation (Thomas and Lewington 2010; Pradel and Fischer 2011). Taken together, these findings lend at least some support to the notion that detrimental weather conditions may negatively affect flight activity and therefore dispersal in insects, as has been also found in other studies (Meyer and Sisk 2001; Dennis and Sparks 2006; Cormont et al. 2011; Kuussaari et al. 2016). Ultimately, this dependence rests on the high thorax temperature needed for flight activity (Pollard and Yates 1994), which cannot be maintained without solar radiation or at cool temperatures.

We additionally investigated the effects of tracking time and observer team. The positive relations between tracking time and several variables were expected, and simply reflect that increasing observation time increases track length, the number of stops, etc. The negative relation between tracking time and step length as well as average speed might indicate that it was easier for the observers to follow slow flying butterflies with short step lengths. Effects of observer team were also widespread, though our statistical approach effectively controlled for both sources of confounding effects.

## Conclusions

Our results show that customary smartphones can be suitable to reveal biologically significant and well interpretable behavioral patterns in insects. In particular, behavioral differences between the sexes could be readily determined. Similarly, earlier studies investigating butterfly flight using harmonic radar (Cant et al. 2005) but also GPS receivers (e.g. Breed and Severns 2015; Fernández et al. 2016) revealed solid data on butterfly flight behavior. More problematic than the technical device used seems to be the rather short observation period of 15 min at max. The principal problem here is that small flying insects are notoriously difficult to follow in the field (cf. Riley and Smith 2002; Cant et al. 2005). Accordingly, many tracks were even substantially shorter than 15 min. In that respect it is actually quite surprising that our study revealed robust data. We thus

conclude that our approach was well appropriate to reveal sexual differences in and impacts of ambient weather on butterfly behaviour. However, tracking butterflies with handheld devices does not seem to be appropriate to estimate rates of dispersal and dispersal distances, such that any conclusions regarding insect dispersal need to be drawn with the greatest care.

## References

- Baguette M, van Dyck H (2007) Landscape connectivity and animal behavior: functional grain as a key determinant for dispersal. *Landsc Ecol* 22:1117–1129
- Baguette M, Blanchet S, Legrand D, Stevens VM, Turlure C (2013) Individual dispersal, landscape connectivity and ecological networks. *Biol Rev* 88:310–326
- Berwaerts K, Matthysen E, van Dyck H (2008) Take-off flight performance in the butterfly *Pararge aegeria* relative to sex and morphology: a quantitative genetic assessment. *Evolution* 62:2525–2533
- Bestion E, Clobert J, Baguette M (2015) Dispersal response to climate change: scaling down to intraspecific variation. *Ecol Lett* 18:1226–1233
- Bonte D, van Dyck H, Bullock JM, Coulon A, Delgado M, Gibbs M, Lehouck V, Matthysen E, Mustin K, Saastamoinen M, Schtickzelle N, Stevens VM, Vandewoestijne S, Baguette M, Barton K, Benton TG, Chaput-Bardy A, Clobert J, Dytham C, Hovestadt T, Meier CM, Palmer SCF, Turlure C, Travis JM (2012) Costs of dispersal. *Biol Rev* 87:290–312
- Bowler DE, Benton TG (2005) Causes and consequences of animal dispersal strategies: relating individual behaviour to spatial dynamics. *Biol Rev* 80:205–225
- Breed GA, Sevens PM (2015) Low relative error in consumer-grade GPS units make them ideal for measuring small-scale animal movement patterns. *Peer J* 3:e1205
- Cant ET, Smith AD, Reynolds DR, Osborne JL (2005) Tracking butterfly flight paths across the landscape with harmonic radar. *Proc Biol Sci* 272:785–790
- Cormont A, Malinowska AH, Kostenko O, Radchuk V, Hemerik L, Wallis DeVries MF, Verboom J (2011) Effect of local weather on butterfly flight behaviour, movement, and colonization: significance for dispersal under climate change. *Biodivers Conserv* 20:483–503
- Cote J, Clobert J (2007) Social information and emigration: lessons from immigrants. *Ecol Lett* 10:411–417
- Dennis RL, Sparks TH (2006) When is a habitat not a habitat? Dramatic resource use changes under differing weather conditions for the butterfly *Plebejus argus*. *Biol Conserv* 129:291–301
- DWD [Deutscher Wetterdienst] (2017) WESTE-XL. [http://www.dwd.de/DE/leistungen/weste/westexl/weste\\_xl.html](http://www.dwd.de/DE/leistungen/weste/westexl/weste_xl.html). Accessed 08 August 2017
- Ducatez S, Humeau A, Congretel M, Fréville H, Baguette M (2014) Butterfly species differing in mobility show different structures of dispersal-related syndromes in the same fragmented landscape. *Ecography* 37:378–389
- Ebert G, Rennwald E (1991) Die Schmetterlinge Baden-Württembergs, vol 2. Ulmer, Stuttgart
- Erhardt A, Thomas JA (1991) Lepidoptera as indicators of change in the semi-natural grasslands of lowland and upland Europe. In: Collins MN (ed) The conservation of insects and their habitats. Academic Press, pp 213–234
- Fernández P, Rodríguez A, Obregón R, de Haro S, Jordano D, Fernández-Haeger J (2016) Fine scale movements of the butterfly *Plebejus argus* in a heterogeneous natural landscape as revealed by GPS tracking. *J Insect Behav* 29:80–98
- Fernández-Chacón A, Stefanescu C, Genovart M, Nichols JD, Hines JE, Paramo F, Turco M, Oro D (2014) Determinants of extinction-colonization dynamics in Mediterranean butterflies: the role of landscape, climate and local habitat features. *J Anim Ecol* 83:276–285
- Fischer K, Fiedler K (2000a) Sex-related differences in reaction norms in the butterfly *Lycaena tityrus* (Lepidoptera: Lycaenidae). *Oikos* 90:372–380
- Fischer K, Fiedler K (2000b) Methodische Aspekte von Fang-Wiederfangstudien am Beispiel der Feuerfalter *Lycaena helle* und *L. hippothoe*. *Beitr Ökol* 4:157–172
- Fischer K, Fiedler K (2001) Resource-based territoriality in the butterfly *Lycaena hippothoe* and environmentally induced behavioural shifts. *Anim Behav* 61:723–732

- Fischer K, Beinlich B, Plachter H (1999) Population structure, mobility and habitat preferences of the violet copper *Lycaena helle* (Lepidoptera: Lycaenidae) in western Germany: implications for conservation. *J Insect Conserv* 3:43–52
- Forester DJ, Machlist GE (1996) Modeling human factors that affect the loss of biodiversity. *Conserv Biol* 10:1253–1263
- GfS [Gesellschaft für Schmetterlingsschutz] (2017) Verbreitungskarte *Lycaena tityrus*. <http://www.ufz.de/european-butterflies/index.php?de=43003>. Accessed 25 July 2017
- Hanski I (1998) Metapopulation dynamics. *Nature* 396:41–49
- Hanski I, Gilpin ME (1997) Metapopulation biology: ecology, genetics, and evolution. Academic Press
- Hanski I, Breuker CJ, Schöps K, Setchfield R, Nieminen M (2002) Population history and life history influence the migration rate of female Glanville fritillary butterflies. *Oikos* 98:87–97
- Hanski I, Erälahti C, Kankare M, Ovaskainen O, Sirén H (2004) Variation in migration propensity among individuals maintained by landscape structure. *Ecol Lett* 7:958–966
- Hickling R, Roy DB, Hill JK, Fox R, Thomas CD (2006) The distributions of a wide range of taxonomic groups are expanding polewards. *Glob Chang Biol* 12:450–455
- Hill JK, Thomas CD, Blakely DS (1999) Evolution of flight morphology in a butterfly that has recently expanded its geographic range. *Oecologia* 121:165–170
- Hopper KR (1999) Risk-spreading and bet-hedging in insect population biology. *Annu Rev Entomol* 44:535–560
- Hovestadt T, Nieminen M (2009) Costs and benefits of dispersal in butterflies. In: Settele J, Shreeve T, Konvicka M, van Dyck H (eds) *Ecology of butterflies in Europe*. Cambridge Univ Press, Cambridge, pp 97–106
- Karl I, Fischer K (2009) Altitudinal and environmental variation in lifespan in the copper butterfly *Lycaena tityrus*. *Funct Ecol* 23:1132–1138
- Karl I, Janowitz SA, Fischer K (2008) Altitudinal life-history variation and thermal adaptation in the copper butterfly *Lycaena tityrus*. *Oikos* 117:778–788
- Kingsolver JG (1983) Ecological significance of flight activity in *Colias* butterflies: implications for reproductive strategy and population structure. *Ecology* 64:546–551
- Kuussaari M, Rytteri S, Heikkinen RK, Heliölä J, von Bagh P (2016) Weather explains high annual variation in butterfly dispersal. *Proc R Soc B* 283:20160413
- Legrand D, Trochet A, Moulherat A, Calvez O, Stevens VM, Ducatez S, Clobert J, Baguette M (2015) Ranking the ecological causes of dispersal in a butterfly. *Ecography* 38:822–831
- Meyer CL, Sisk TD (2001) Butterfly response to microclimatic conditions following ponderosa pine restoration. *Restor Ecol* 9:453–461
- Müller T (2006) *Klima-Journal für Vorpommern, Anklam*
- Murphy DD, Freas KE, Weiss SB (1990) An environment-metapopulation approach to population viability analysis for a threatened invertebrate. *Conserv Biol* 4:41–51
- Niitepõld K, Smith AD, Osborne JL, Reynolds DR, Carreck NL, Martin AP, Marden JH, Ovaskainen O, Hanski I (2009) Flight metabolic rate and Pgi genotype influence butterfly dispersal rate in the field. *Ecology* 90:2223–2232
- Pollard E, Yates TJ (1994) *Monitoring butterflies for ecology and conservation: the British butterfly monitoring scheme*. Springer Science and Business Media, Luxemburg
- Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE (2010) Global pollinator declines: trends, impacts and drivers. *Trends Ecol Evol* 25:345–353
- Pradel K, Fischer K (2011) Living on the edge: habitat and host-plant selection in the butterfly *Lycaena tityrus* (Lepidoptera: Lycaenidae) close to its northern range limit. *J Res Lepid* 44:35–41
- Riley JR, Smith AD (2002) Design considerations for an harmonic radar to investigate the flight of insects at low altitude. *Comput Electron Agric* 35:151–169
- Roy DB, Sparks TH (2000) Phenology of British butterflies and climate change. *Glob Chang Biol* 6:407–416
- Saastamoinen M, Brakefield PM, Ovaskainen O (2012) Environmentally induced dispersal-related life-history syndrome in the tropical butterfly, *Bicyclus anynana*. *J Evol Biol* 25:2264–2275
- Sanderson EW, Jaiteh M, Levy MA, Redford KH, Wannebo AV, Woolmer G (2002) The human footprint and the last of the wild. *Bioscience* 52:891–904
- Stevens V, Turlure C, Baguette M (2010) A meta-analysis of dispersal in butterflies. *Biol Rev* 85:625–642
- Therry L, Zawal A, Bonte D, Stoks R (2014) What factors shape female phenotypes of a poleward-moving damselfly at the edge of its range? *Biol J Linn Soc* 112:556–568
- Thomas JA, Lewington R (2010) *The butterflies of Britain and Ireland*. British Wildlife Publishing, Milton on Stour
- Tolman T, Lewington R (1998) *Die Tagfalter Europas und Nordwestafrikas*. Franckh-Kosmos, Stuttgart

- Travis MJJ, Delgado M, Bocedi G, Baguette M, Barton K, Bonte D, Boulangeat I, Hodgson JA, Kubisch A, Penteriani V, Saastamoinen M, Stevens VM, Bullock JM (2013) Dispersal and species' responses to climate change. *Oikos* 122:1532–1540
- Trochet A, Legrand D, Larranga N, Ducatez S, Calvez O, Cote J, Clobert J, Baguette M (2013) Population sex ratio and dispersal in experimental, two-patch metapopulations of butterflies. *J Anim Ecol* 82:946–955
- van Dyck H, Baguette M (2005) Dispersal behaviour in fragmented landscapes: routine or special movements? *Basic Appl Ecol* 6:535–545
- van Dyck H, Wiklund C (2002) Seasonal butterfly design: morphological plasticity among three developmental pathways relative to sex, flight and thermoregulation. *J Evol Biol* 15:216–225
- van Dyck H, Matthysen E, Wiklund C (1998) Phenotypic variation in adult morphology and pupal colour within and among families of the speckled wood butterfly *Pararge aegeria*. *Ecol Entomol* 23:465–472
- van Swaay CAM (1990) An assessment of the changes in butterfly abundance in the Netherlands during the 20th century. *Biol Conserv* 52:287–302
- Warren MS, Hill JK, Thomas JA, Asher J, Fox R, Huntley B, Roy DB, Telfer MG, Jeffcoate S, Harding P, Jeffcoate G, Willis SG, Greatorex-Davies JN, Moss D, Thomas CD (2001) Rapid responses of British butterflies to opposing forces of climate and habitat change. *Nature* 414:65–69
- Wickman PO (1985) Territorial defence and mating success in males of the small heath butterfly, *Coenonympha pamphilus* L. (Lepidoptera: Satyridae). *Anim Behav* 3:1162–1168
- Wickman PO (2009) Thermoregulation and habitat use in butterflies. In: Settele J, Shreeve T, Konvicka M, van Dyck H (eds) *Ecology of butterflies in Europe*. Cambridge Univ Press, Cambridge, pp 55–61

## **Electronic supplementary material**

**Journal of Insect Behavior**

**Movement patterns differ between sexes and depend on weather conditions in the butterfly *Lycaena tityrus***

Elisabeth Reim\*, Ingo Arnstedt, Isabel Barwisch, Max Baumgarten, Sascha Bock, Julia Eberspach, Julia Ellerbrok, Mulugeta Gebremeskel, Simon Küpper, Lukas Guth, Aninha Lassen, Letro Letro, Rebecca Meth, Maria Möller, Felix Närmann, Inga Neunaber, Alexander Seliger, Wilderich V. Stein, Carolin Vallinga, Philipp Vögele, Martin Wagner & Klaus Fischer

\* Corresponding author:


Zoological Institute and Museum, Greifswald University, D-17489 Greifswald, Germany

E-Mail: [elisabeth.reim@uni-greifswald.de](mailto:elisabeth.reim@uni-greifswald.de)


**Table 1** Range of tested weather variables (hourly means) during the observation period

	Observed minimum value	Observed maximum value
Temperature [°C]	14.4	28.5
Wind speed [m/s]	1.8	9.5
Relative humidity [%]	38	91
Precipitation [mm]	0	1.5

**Fig. 1** Relationship between the exact distance moved and the distance tracked with the application “Geo Tracker – GPS tracker” for 30 tracks with a length of 2.5 to 30 meters (a). The distance measured and the distance tracked were positively correlated ( $r= 0.9945$ ). Tracks were measured at intervals of 2.5 m using a marked square of 100 m<sup>2</sup> (b), see (b) for an example, a track of 15 m (marked in red)



**Fig. 1a**



**Fig. 1b**

**Fig. 2** Relationships between the weather variables (a) temperature and wind, (b) temperature and relative humidity, (c) temperature and precipitation, (d) wind and relative humidity, (e) wind and precipitation, (f) relative humidity and precipitation, (g) cloud cover and temperature, (h) cloud cover and wind, (i) cloud cover and relative humidity, and (j) cloud cover and precipitation




Fig. 2a




Fig. 2b




Fig. 2c




Fig. 2d




Fig. 2e




Fig. 2f




Fig. 2g




Fig. 2h




Fig. 2i






Fig. 2j

**Fig. 3** Relationship between the variables (a) linear distance moved and cloud cover and (b) linear distance moved and temperature



**Fig. 3a**



**Fig. 3b**

## 4.2 Emigration propensity and flight performance are decoupled in a butterfly

Submitted as:

Reim, E., Baguette, M., Günter, F. & Fischer, K. (2018) Emigration propensity and flight performance are decoupled in a butterfly. In revision.



1 Running head: Flight behavior in a butterfly

2

3 **Emigration propensity and flight performance are**  
4 **decoupled in a butterfly**

5

6 Elisabeth Reim<sup>1,3</sup>, Michel Baguette<sup>2</sup>, Franziska Günter<sup>1</sup>, and Klaus Fischer<sup>1</sup>

7

8 <sup>1</sup> Zoological Institute and Museum, Greifswald University, D-17489 Greifswald, Germany.

9 <sup>2</sup> Mus. Natl d'Histoire Naturelle, UMR 7205 Inst. of Systematics, Evolution, Biodiversity, 57,

10 rue Cuvier, FR-75231 Paris Cedex 05, France and also at Centre National de la Recherche

11 Scientifique, SEEM (Station d'Ecologie Expérimentale du CNRS à Moulis), USR2936, Moulis,

12 France.

13

14 <sup>3</sup> Corresponding author:

15 Elisabeth Reim

16 ORCID-ID: 0000-0003-4139-2330

17 Zoological Institute and Museum

18 Greifswald University

19 Soldmannstraße 14

20 D-17489 Greifswald, Germany

21 Tel.: +49-3834-4204297

22 E-Mail: elisabeth.reim@uni-greifswald.de

23

24 *Abstract.* Dispersal, defined as any movements potentially leading to gene flow, is a major  
25 process driving a species' capacity to cope with human-induced environmental modifications.  
26 However, the dispersal process is multi-causal, which currently hinders predictions regarding a  
27 species' resilience to global change. We used a multifaceted approach to disentangle the relative  
28 importance of a suite of dispersal-related factors in the butterfly *Lycaena tityrus*, including  
29 condition- (morphology, behavior) and context- (environmental) dependent factors.  
30 Experiments were conducted at an experimental platform dedicated to study dispersal, the  
31 Metatron, to record emigration propensity in two-patch experimental metapopulations under  
32 different environmental conditions. The butterflies' individual condition was subsequently  
33 assessed in the laboratory. Individual condition did not generally influence emigration  
34 propensity. We detected a significant sex-bias in emigration propensity, being in general higher  
35 in females than in males, but in a context-dependent manner. The environmental context  
36 affected emigration propensity, which was higher when habitat quality was poor. Our results  
37 show that emigration is not random in *L. tityrus*, but is rather an active process triggered by sex  
38 and habitat quality. Our main finding was that individual condition, and particularly flight  
39 ability measured by a performance test, was not related to emigration propensity in  
40 experimental metapopulations. Our results may have important implications for forecasting  
41 future species distributions, as deteriorating environmental contexts are likely to increase  
42 emigration whatever the individual condition is.

43

44 *Keywords:* Climate change; condition; context-dependence; dispersal; flight ability; *Lycaena*  
45 *tityrus*; metapopulation; Metatron; movement

46

## INTRODUCTION

47  
48 Dispersal, often defined as any movements potentially leading to gene flow (Ronce 2007), is a  
49 complex process serving different functions and involving a variety of underlying mechanisms  
50 (Bowler and Benton 2005, Bonte et al. 2012, Matthysen 2012). The multi-causality of dispersal  
51 though has been fully appreciated in recent years only (reviewed by Matthysen 2012),  
52 suggesting that complex interactions between several factors rather than a single one influence  
53 the dispersal process (Legrand et al. 2015). Currently there is a lack of studies aiming at  
54 disentangling the relative importance of multiple factors relevant to dispersal including  
55 individual condition (e.g. morphology, behavior) and the environmental context (e.g. habitat  
56 quality, weather; but see e.g. Legrand et al. 2015, Turlure et al. 2016). Also, growing evidence  
57 indicates that dispersal is not a fixed but a highly plastic trait (e.g. Ronce 2007). In invertebrates,  
58 for instance, flight behavior is influenced by food availability (e.g. Saastamoinen et al. 2010)  
59 and weather conditions (Niitepõld et al. 2009). Consequently, organisms may collect  
60 information on their current environment influencing their dispersal decision (Clobert et al.  
61 2009). However, only a few studies have focused on such informative cues that are likely to  
62 trigger dispersal decisions (but see Cote and Clobert 2007, Baguette et al. 2011).

63  
64 The above deficiencies in our current understanding of dispersal are worrying given its crucial  
65 ecological and evolutionary importance, ranging from metapopulation dynamics through  
66 metacommunity regulation to gene flow and speciation (Clobert et al. 2012). Dispersal is also  
67 of utmost importance in driving a species' capacity to respond to human-induced global change  
68 like unpredictable climatic variation or habitat loss and fragmentation (Cote and Clobert 2007,  
69 Travis et al. 2013). Persistence under such dramatic changes requires a species to either stay  
70 and adapt to the new conditions or to leave by moving along the waves of change (Berg et al.  
71 2010). In the latter case, dispersal is a key process allowing individuals to track their shifting

72 climate niche (resulting in species range shifts; Warren et al. 2001, Hickling et al. 2006) or to  
73 maintain meta-population connectivity (Baguette et al. 2013).

74

75 Against this background, we use a multifaceted approach to uncover the relative role of factors  
76 that potentially affect dispersal, including both individual condition (morphology and flight  
77 performance) and environmental context. This approach is in line with the conceptual  
78 framework for movement ecology developed by Nathan et al. (2008), which integrates the basic  
79 components and processes involved in the movement of individual organisms. Consequently,  
80 the movement path of an organism results from dynamic interactions between the internal state,  
81 motion and navigation capacity, and external factors. As our study organism we used the Sooty  
82 Copper butterfly *Lycaena tityrus*, a currently northward expanding species (Brunzel et al. 2008,  
83 Settele et al. 2008), rendering the investigation of its movement drivers particularly interesting  
84 in the current era of climate change.

85

86 We first examined the influence of different environmental factors on the first step of the  
87 dispersal process, i.e. the propensity to leave a habitat patch (emigration propensity). Further  
88 steps are the transfer phase (movement) and the settlement in a novel habitat (see Bonte et al.  
89 2012 for a thorough description of dispersal as a three step process). We used the Metatron, a  
90 unique experimental platform designed to study dispersal in terrestrial organisms in  
91 experimental metapopulations (Legrand et al. 2012). Second, we measured flight performance  
92 (here: time in flight under stress) to examine connections between emigration propensity under  
93 semi-natural conditions and physiological flight performance under laboratory conditions  
94 (Ducatez et al. 2012, Legrand et al. 2015). Third, we assessed individual variation in several  
95 morphological parameters (e.g. thorax-abdomen ratio, wing loading, fat content) that may be  
96 associated with movement ability (Legrand et al. 2016). Note that morphological differences

97 may have important consequences for flight performance (Arnold 1983, Gilchrist 1990,  
98 Berwaerts et al. 2002), and even for dispersal and colonization success in butterflies (Hill et al.  
99 1999, Sekar 2012). How frequently and to what extent such differences directly affect dispersal  
100 behavior though is currently not clear. A study of Hanski et al. (2002), for instance, found no  
101 effect of morphology on dispersal rates.

102

103 We aim to disentangle the relative importance of individual condition versus environmental  
104 context to answer the following questions: (1) To what extent is emigration propensity affected  
105 by the environmental context potentially indicative of habitat quality? (2) Is emigration  
106 propensity in experimental metapopulations related to flight performance, i.e. do individuals  
107 with better flight performance emigrate more readily? (3) Which condition-dependent  
108 individual factors (e.g. morphology, sex) are related to variation in emigration propensity and  
109 flight performance? We predict (1) increased emigration rates if habitat quality is poor, (2) a  
110 high correlation between flight performance and emigration propensity, and consequently (3)  
111 that flight performance and emigration propensity are tightly linked to morphological traits.  
112 Answering these questions might be important for improving predictions on species responses  
113 to environmental change.

114

115

116

## METHODS

117

### *Study organism*

118 *Lycaena tityrus* (Poda, 1761) is a widespread temperate-zone butterfly ranging from Western  
119 Europe to central Asia (Ebert and Rennwald 1991). This species is currently expanding its range  
120 towards higher latitudes and altitudes, which is assumed to be largely driven by anthropogenic  
121 climate change (Brunzel et al. 2008, Settele et al. 2008). The species is bivoltine with two

122 discrete generations per year in most parts of its range, although populations with one or three  
123 generations per year occur. Overwintering takes place as half-grown larva. The principal larval  
124 host-plant is *Rumex acetosa* L. (Polygonaceae), but several congeneric *Rumex* species are  
125 utilized as well (Ebert and Rennwald 1991, Tolman and Lewington 1998, Settele et al. 2008).  
126 Adults are nectar feeders (Ebert and Rennwald 1991). For the following experiments, a total of  
127 35 freshly eclosed, mated females from a bivoltine German population (vicinity of Greifswald,  
128 north-eastern Germany; 54°02N, 13°26E) were caught in June 2015. All females were  
129 transferred to a climate chamber at Greifswald University for egg laying.

130

### 131 *Animal rearing*

132 Caught females were kept at 26°C, 60% relative humidity, and a L18:D6 photoperiod. For  
133 oviposition, females were placed separately in translucent 1 L plastic pots covered with gauze,  
134 and were provided with *Rumex acetosa* (oviposition substrate), fresh flowers, water, and a  
135 highly concentrated sucrose solution (for adult feeding). Deposited eggs were collected daily  
136 and transferred, separated by female, to small plastic boxes and kept under the same conditions  
137 used for oviposition. After hatching, larvae were placed in groups of 10-15 individuals in larger,  
138 transparent plastic boxes (500 ml), containing moistened filter paper and fresh cuttings of *R.*  
139 *acetosa* in ample supply. Boxes were checked daily and larvae were supplied with fresh food  
140 as necessary. Resulting pupae were transferred to a climate cabinet set to 15°C (60% r.h.,  
141 L18:D6) in order to retard further development. Once most larvae had pupated, pupae were  
142 transferred to the ‘Station for Theoretical and Experimental Ecology’ (SETE) of the CNRS in  
143 Ariège, south-western France. There, pupae were randomly divided into three groups and  
144 initially also kept at 15°C. Three days before the start of the respective experiment (see further  
145 below), pupae were transferred to warmer conditions (26-28°C) to facilitate quick and  
146 synchronized adult eclosion. After adult eclosion, butterflies were individually marked by

147 writing a number on their left hindwing using a water-proof felt-tip pen. Sexes were kept  
148 separately until the start of experiments. Butterflies were collected over up to three eclosion  
149 days to reach sufficiently high numbers, being fed with flowers and a highly concentrated  
150 sucrose solution. Consequently, butterflies were 1-3 days old upon their release into the  
151 Metatron (see below).

152

### 153 *Metatron setup and experimental design*

154 The Metatron is an experimental platform consisting of interconnected mesh cages ('patches')  
155 operated by the 'Station for Theoretical and Experimental Ecology' (SETE) of the CNRS. It  
156 allows for the manipulation of spatial and environmental factors in a semi-natural environment  
157 (Legrand et al. 2012). Each cage is 200 m<sup>3</sup> (10 x 10 x 2 m) large and is covered by insect-proof  
158 gaze. The ground within the cages is covered by natural vegetation. We carried out three  
159 experiments in the Metatron as outlined below. Each experiment involved two-patch  
160 experimental meta-populations, consisting of a departure and an arrival cage connected by a  
161 single, S-shaped corridor (19 m long). To start the experiments, butterflies were released into  
162 the departure cage, in which they could either stay or which they could leave through the  
163 corridor leading to the arrival cage. Thereafter, butterfly positions were monitored through  
164 several recapture sessions (see below). To allow for an appropriate discrimination between  
165 dispersers and resident individuals, corridors were designed to be challenging (corridor entries  
166 represent less than 2% of the total vertical surface area of patches; Legrand et al. 2012).  
167 Moreover, the S-shape of the corridor prevented dispersing butterflies to see the arrival cage.  
168 Emigration propensity was scored as the number of butterflies leaving the departure cage, i.e.  
169 being re-captured either in the arrival cage or corridor at least once. The number of repeated  
170 dispersers, i.e. butterflies moving back and forth between the cages more than once, was low  
171 (~ 2%). The vegetation within both cages and the corridors was cut (turf height 10 cm) prior to

172 the start of experiments to allow for the manipulation of host and nectar plant availability (see  
173 below).

174

175 In the first two experiments we tested the effect of one environmental factor each, by using 4  
176 cages to create 2 two-patch metapopulations per experiment. In experiment 1, we exclusively  
177 manipulated resource availability, i.e. one of the departure cages entailed resources (25 pots  
178 with flowering nectar plants and additionally 20 pots with host plants) while the other did not.  
179 All cages were exposed to full sunlight ('sunny'), conditions suitable for high flight activity.  
180 Throughout all experiments, arrival cages had identical conditions with resources and sunny  
181 conditions being available, while no resources were available within the corridors. In  
182 experiment 2, we exclusively manipulated solar radiation. Here, individuals in both departure  
183 cages had access to an equal though limited number of flowering nectar plants (8 pots, no access  
184 to host plants), while one departure cage provided sunny and the other one shady and therefore  
185 cooler conditions. Shady conditions were achieved through closing motor-driven shutters  
186 positioned above the cages (Legrand et al. 2012).

187

188 In the third experiment, we investigated both environmental factors to test for additive /  
189 interactive effects. Here, we used in total 8 cages, i.e. 4 two-patch metapopulations. We  
190 manipulated resource availability and solar radiation in the departure cages. Two of the  
191 departure cages contained resources (25 pots with flowering nectar plants and 20 pots with host  
192 plants), while the other two did not. Per resource type, one cage provided sunny and the other  
193 one shady conditions. Consequently, we had four types of departure cages: (1) resources and  
194 sun, (2) resources and shade, (3) no resources and sun, (4) no resources and shade. Experiments  
195 were conducted between 15<sup>th</sup> and 29<sup>th</sup> July 2015.

196

197 We released 138 and 150 into the two departure cages in experiment 1, 92 and 102 in  
198 experiment 2, and 107, 108, 113 and 119 butterflies into the 4 cages in experiment 3 (n = 929  
199 in total). Numbers of males and females were equal per cage. In general, *L. tityrus* is a rather  
200 sedentary species, which may occur in high local densities in high-quality habitat patches. In  
201 experiment 1 and 2 all butterflies were 2 days old upon release into the Metatron, while in  
202 experiment 3 age ranged between 1-3 days. The release of butterflies always took place in the  
203 morning at 9 a.m. Recapture sessions started in the afternoon (3 p.m.) of the same day and were  
204 thereafter performed twice a day (between 9-12 a.m. and 3-6 p.m.; except for rainy conditions).  
205 The position (i.e. departure cage, corridor, or arrival cage) of each individual was recorded per  
206 session for the following 2.5 (experiments 1 and 2) or 3.5 days (experiment 3). The short  
207 observation periods resulted from high mortality within the Metatron due to biotic (predators,  
208 e.g. spiders) and abiotic (e.g. heat, thunderstorms) factors (mortality rates: experiment 1 = 68%,  
209 experiment 2 = 53%, experiment 3 = 83%). Mortality rates were assessed at the end of each  
210 experiment. Butterflies found dead were replaced as long as there were enough freshly eclosed  
211 butterflies available. During the last recapture session, all remaining butterflies were caught and  
212 subjected to a flight performance test. As *L. tityrus* is a native species in Ariège where the  
213 Metatron is located, special care was taken to prevent the escape of butterflies to control the  
214 risk of admixture between experimental and wild individuals.

215

216

#### *Flight performance test*

217

218

219

220

221

Butterflies were tested for flight performance using a well-established assay (cf. Ducatez et al.  
2012). Each individual was placed into a plastic chamber (30 x 16 x 14 cm), which was  
perforated at its base and fixed to a rapid agitator (IKA Vortex 4 digital). After a habituation  
period of 30 seconds, the vortex was switched on to strongly shake the chamber, preventing the  
butterfly from holding onto the walls. Thus, during the test, butterflies had to fly or were lying

222 uncomfortably at the bottom of the shaking chamber. The time an individual butterfly spent  
223 flying was recorded for 60 seconds, with higher values reflecting a better flight performance.  
224 Experiments were performed at  $22 \pm 0.5^\circ\text{C}$ , mimicking summer (July) conditions in  
225 Greifswald. After the performance test, butterflies were frozen in liquid nitrogen for later  
226 analyses of potentially flight-related parameters.

227

228

### *Morphological analyses*

229 To test for associations between emigration propensity in the Metatron or flight performance  
230 and other traits, we measured several morphological traits. Therefore, frozen butterflies were  
231 back-transferred to Greifswald University. First, adult body mass was determined to the nearest  
232 0.01 mg (Sartorius CPA225D). Then, wings, heads and legs were removed. Thorax and  
233 abdomen were separated and afterwards weighed. Abdomen fat content was measured  
234 following Fischer et al. (2003), but using the less poisonous acetone instead of dichloromethane.  
235 Abdomen were first dried to constant weight for two days at  $70^\circ\text{C}$ . Abdomen dry mass was  
236 measured. Afterwards, fat was extracted twice, each time for 48 h using 1 ml of acetone  
237 ( $\text{C}_3\text{H}_6\text{O}$ ) per butterfly. Solutions were exchanged between both fat extractions. Then, abdomen  
238 were again dried for two days at  $70^\circ\text{C}$ , after which the fat-free dry mass was measured. Absolute  
239 (mg) and relative (%) fat content were determined as the mass difference between abdomen dry  
240 mass and the remaining dry mass after the two fat extractions. Forewing area and length (from  
241 basis to apex) were measured using digital images of left forewings (captured from ventrally  
242 with a digital camera mounted on a stereo microscope) and NIS elements software. Wing  
243 loading was calculated as total body mass divided by forewing area, and wing aspect ratio as  $4$   
244  $\times$  forewing length<sup>2</sup> divided by the forewing area (Berwaerts et al. 2002). Thorax-abdomen ratio  
245 was calculated by dividing thorax through abdomen mass.

246

247 *Statistical analyses*

248 Data on morphological traits were analyzed with general linear mixed models (GLMMs) using  
249 resources, sun and sex as fixed effects and family as a random effect. In experiment 3, age  
250 (fixed factor) was used additionally as covariate (note that experiments 1 and 2 all individuals  
251 were 2-days old; see above). Data on individual flight performance and emigration propensity  
252 (departure cage left: yes / no) were analyzed using generalized linear models (GzLMs) with  
253 either a normal error distribution and a log-link function (flight performance) or a binomial  
254 error distribution and logit-link function (emigration propensity), using the same factors as  
255 above. Only butterflies where both emigration propensity and flight performance data were  
256 available were used for the analyses. All above models were constructed by a stepwise  
257 backwards elimination of non-significant factors with p-values > 0.1. Note that an AIC  
258 approach yielded identical results.

259  
260 To additionally test for the impact of continuous variables (morphology) on flight performance  
261 and emigration propensity, we first performed, separately for males and females, principal  
262 component analyses (PCAs) owing to strong inter-correlations among traits (Appendix S1:  
263 Table S1). The sexes were analyzed separately because GLMM results indicated that males and  
264 females differ strongly in their morphological traits (see below). We used the first four principal  
265 components (PCs) for further analyses, having Eigenvalues between 4.1 and 0.7 for males and  
266 between 5.1 and 1.0 for females (Appendix S1: Table S2). Thus, all PCs explaining  $\geq 7\%$  of  
267 the variance were included. We then constructed another set of models by including the  
268 respective PCs (and flight performance for emigration propensity) as covariates to the models  
269 mentioned above, again followed by a stepwise backwards elimination of non-significant  
270 factors. All statistical tests were performed with Statistica 12.0 (Tulsa, StatSoft, OK). All means  
271 are given  $\pm 1$  SE.

272

273

274

## RESULTS

275

### *Experiment 1*

276 Significant sex differences were found for all morphological traits except from thorax mass,  
277 absolute fat content, forewing length, and wing loading (Appendix S1: Table S3), indicating  
278 that females had on average higher abdomen and adult body masses, a lower relative fat content,  
279 thorax-abdomen ratio and wing aspect ratio, and a larger wing area than males (Table 1). A  
280 significant effect of resource availability was found for thorax mass only, which was higher in  
281 individuals released into departure cages with resources. However, resource availability was  
282 involved in significant interactions with sex for total body mass, abdomen mass, thorax-  
283 abdomen ratio, absolute fat content, and wing loading. These interactions indicate that sexual  
284 differences were more pronounced in or restricted to individuals being released into cages with  
285 resources (Table 1). Differences among families were significant for thorax mass, relative and  
286 absolute fat content, wing length, and wing area, and tendencies were found for total body mass  
287 and wing aspect ratio.

288

289 Emigration propensity in the Metatron was significantly affected by resource availability only  
290 (Table 2a). Individuals with no access to resources showed a higher emigration propensity (38%  
291 of 39 vs. 15% of 47; Fig. 1a). Including principal components and flight performance as  
292 covariates revealed no significant effects on emigration propensity in either males or females,  
293 but the factor resources was still included in both models (Table 2a). Additionally, the factor  
294 flight performance was included into the model for males, indicating a non-significant tendency  
295 towards a negative correlation between flight performance and emigration propensity (beta -  
296  $0.11 \pm 0.11$ ).

297

298 Flight performance was significantly affected by the factors sex and resource availability (Table  
299 2b). Males showed a longer flight endurance than females ( $36.2 \pm 7.6 > 25.6 \pm 3.6$  s), and  
300 individuals that were released into a departure cage with resources flew longer than individuals  
301 that were released into a departure cage without resources ( $38.4 \pm 4.7 > 23.4 \pm 6.5$  s; Fig. 1b).  
302 Including principal components as covariates additionally revealed a significant impact of PC2  
303 and family for males and PC4 for females (Table 2b). Flight performance was negatively  
304 correlated with PC2 in males (beta  $-0.38 \pm 0.22$ ). PC2 in turn was most strongly (positively)  
305 correlated with wing area ( $r = 0.912$ ), wing length ( $r = 0.865$ ), and thorax mass ( $r = 0.883$ ;  
306 Appendix S1: Table S2a). Thus, smaller males had a better flight performance. Flight  
307 performance was negatively correlated with PC4 in females (beta  $-0.10 \pm 0.13$ ). PC4 in turn  
308 was most strongly (negatively) correlated with aspect ratio ( $r = -0.972$ ). Thus, a higher aspect  
309 ratio increased flight performance in females (Appendix S1: Table S2b).

310

311

### *Experiment 2*

312 Significant sex differences were found for all morphological traits except from thorax mass,  
313 relative fat content, and forewing length (Appendix S1: Table S4). Females had on average a  
314 higher abdomen and adult body masses, a lower thorax-abdomen ratio, a higher absolute fat  
315 content, a larger wing area, a higher wing loading, and a lower wing aspect ratio than males  
316 (Table 3). A significant effect of sunny versus shady conditions was found for abdomen mass,  
317 thorax-abdomen ratio, and absolute and relative fat content. Individuals that were released into  
318 sunny cages had on average a lower abdomen mass, concomitantly a higher thorax-abdomen  
319 ratio, and a lower absolute and relative fat content. Furthermore, significant sun by sex  
320 interactions were found for abdomen mass, thorax-abdomen ratio, relative fat content, and wing  
321 loading (Appendix S1: Table S4). They show that sexual differences were more pronounced

322 under sunny as compared with shady conditions, except for relative fat content (Table 3).  
323 Differences among families were significant for thorax mass, wing length, and wing area.

324

325 Emigration propensity was significantly affected by sunny versus shady conditions only (Table  
326 4a). Individuals showed a higher emigration propensity when released into a shady rather than  
327 sunny departure cage (36.7% of 49 > 4.8% of 42; Fig. 2a). This pattern also prevailed in females  
328 when being analyzed separately. Including principal components and flight performance as  
329 covariates did not change the above results, as no covariate was included within the final  
330 models.

331

332 Flight performance was not affected by the factors sex, sun or family in the first model.  
333 Including covariates revealed no significant influence of any factors on females, but a  
334 significant impact of PC1 to 4 on males. Additionally, sun and family tended to influence flight  
335 performance in males (Table 4b). Males that were released into shady departure cages tended  
336 to fly marginally longer than those released into sunny departure cages ( $33.1 \pm 8.6 > 30.3 \pm 7.4$   
337 s; Fig. 2b). Male flight performance was negatively correlated with PC1 (beta  $-0.27 \pm 0.27$ ),  
338 PC2 (beta  $-0.05 \pm 0.26$ ) and PC4 (beta  $-0.05 \pm 0.30$ ) and positively with PC3 (beta  $0.10 \pm 0.30$ ).  
339 PC1 in turn was most strongly (positively) correlated with abdomen mass ( $r = 0.953$ ), PC2 most  
340 strongly (positively) with wing area ( $r = 0.912$ ), PC3 and PC4 most strongly (positively) with  
341 aspect ratio (PC3:  $r = 0.743$ , PC4:  $r = 0.563$ ; Appendix S1: Table S2a). Thus, males with a  
342 lower abdomen mass and thus wing loading and a smaller wing area showed a better flight  
343 performance, while aspect ratio showed no clear association with flight performance.

344

### Experiment 3

345

346 Significant sex differences were found for all morphological traits measured except from thorax  
347 mass, relative fat content, and forewing length (Appendix S1: Table S5), indicating that females  
348 had on average higher abdomen and total body masses, a lower thorax-abdomen ratio, a higher  
349 absolute fat content (mg), a larger wing area, a higher wing loading, and a lower wing aspect  
350 ratio than males (Table 5). A significant effect of resource availability was found for relative  
351 and absolute fat content only, showing that individuals that had access to resources had a lower  
352 relative ( $7.9 \pm 0.6\%$  vs.  $15.1 \pm 1.4\%$ ) and absolute ( $0.31 \pm 0.03$  mg vs.  $0.70 \pm 0.08$  mg) abdomen  
353 fat content. Effects of sunny versus shady conditions, in contrast, were non-significant  
354 throughout, as was the case for all interaction terms. Differences among families were  
355 significant for wing aspect ratio only. A significant effect of age was found for wing area only,  
356 indicating that older butterflies had larger wings than younger ones (thus, largest butterflies  
357 eclosed first).

358

359 Emigration propensity was significantly affected by the factors resource availability, the  
360 interaction between resources and sex, and the three-way interaction with sun (Table 6a). Thus,  
361 individuals that were released into departure cages without resources showed a higher  
362 emigration propensity (46% of 28 > 24% of 45). The resources by sex interaction shows that  
363 females had a higher emigration propensity than males if resources were present in the departure  
364 cage, while it was the other way round if no resources were present (Fig. 3a). Consequently,  
365 emigration propensity increased much more strongly in males than in females when no  
366 resources were present. The significant three-way interaction additionally shows that males  
367 were more dispersive than females only if a lack of resources coincided with sunny conditions  
368 (60% of 5 vs. 22% of 9). Under all other conditions, females were the more dispersive sex  
369 (resources + sun: 60% of 10 vs. 0% of 7; resources + shade: 22% of 18 vs. 10% of 10; no

370 resources + shade: 58% of 12 vs. 50% of 2). Including principal components and flight  
371 performance as covariates revealed an effect of the interaction between resources and sun for  
372 females and PC3 for males (Table 6a). The resources by sun interaction for females shows that  
373 females had a higher emigration propensity if resources and sun were present (60% of 10) or  
374 both absent (58% of 12) in the departure cage, while emigration propensity was lower when  
375 one of the factors was absent (resources + shade: 22% of 18; no resources + sun: 22% of 9). In  
376 males, PC3 showed a positive correlation with emigration propensity (beta  $0.50 \pm 0.19$ ). PC3  
377 in turn was most strongly (positively) correlated with aspect ratio ( $r = 0.743$ ), indicating that  
378 emigration propensity was positively associated with aspect ratio in males (Appendix S1: Table  
379 S2a).

380

381 Flight performance in the vortex was significantly affected by the factors sex, sun, and family  
382 (Table 6b). Males showed a better flight performance than females ( $38.3 \pm 7.0 > 25.6 \pm 4.8$  s),  
383 and individuals that were released into shady departure cages flew longer than those released  
384 into sunny departure cages ( $39.0 \pm 5.3 > 24.9 \pm 6.5$  s; Fig. 3b). Additionally including principal  
385 components as covariates revealed a significant impact of PC1 for females and PC3 for males.  
386 Flight performance was negatively correlated with PC3 in males (beta  $-0.24 \pm 0.22$ ). PC3 in  
387 males was most strongly (positively) correlated with aspect ratio ( $r = 0.743$ , Appendix S1: Table  
388 S2a), meaning that flight performance was negatively correlated with aspect ratio in males.  
389 Flight performance was negatively correlated with PC1 in females (beta  $-0.27 \pm 0.14$ ). PC1 in  
390 females was in turn most strongly (negatively) correlated with body mass ( $r = -0.974$ ; Appendix  
391 S1: Table S2b). Thus, flight performance was positively related to body mass in females.

392

393

## DISCUSSION

394

395 We start our discussion with variation in morphological traits, as these have been implied to  
396 affect flight ability in insects and may thus underlie variation in emigration propensity.

397

398

### *Sexual differences in morphological traits*

399 The morphological traits measured were, as expected, strongly affected by sex. Our results  
400 showed consistently that females were heavier than males, which is usually explained by  
401 fecundity selection for large body size in females, as both traits are often positively related  
402 (Honek 1993, Blanckenhorn 2000). Male insects, in contrast, often favor early adult emergence  
403 to maximize mating opportunities, resulting in rapid development which is in turn traded off  
404 against body size (selection for protandry; Wiklund and Fagerström 1977, Karl and Fischer  
405 2008). In accordance with above, mass differences between the sexes were caused by  
406 pronounced variation in abdomen mass, enabling high fecundity and large amounts of storage  
407 reserves to fuel egg production in females (Eaton 1988, Tammaru et al. 1996, Berwaerts et al.  
408 2002). Note here the higher absolute fat content found in females, at least under favorable  
409 conditions, while relative fat content is typically higher in males in order to fuel flight activity  
410 (Zera et al. 1998, Karl and Fischer 2008).

411

412 The females' higher mass results in a higher wing loading compared with males, which is likely  
413 detrimental to flight performance (Wickman 2009, Saastamoinen et al. 2012). Males and  
414 females also differed clearly in wing morphology. Despite similar wing lengths, females  
415 showed larger wing areas resulting from having wider wings, while those of males are more  
416 elongated as evidenced by a higher wing aspect ratio (Dudley 2000, Hassall 2015). Thorax  
417 mass, in contrast, did not differ among sexes, demonstrating that males have relatively larger  
418 thoraxes than females, resulting in a consistently increased thorax-abdomen ratio (Berwaerts et

419 al. 2002, Karl et al. 2008). The above sex-specific differences in morphology may result from  
420 differential selective pressures favoring excellent flight performance to succeed in competition  
421 for mates in males (Van Dyck and Wiklund 2002, Berwaerts et al. 2006).

422

### 423 *Environmental differences in morphological traits*

424 As expected for a study carried out under semi-natural conditions, effects of environmental  
425 conditions were generally weaker and less consistent than sexual differences. Storage reserves,  
426 namely fat content, were reduced if resources were present in the departure cages in experiment  
427 3. We assume that this counterintuitive result stems from butterflies remaining largely inactive  
428 when being deprived of food in order to save energy and to endure the critical phase  
429 (Saastamoinen and Hanski 2008). Furthermore, females need nectar for egg production in  
430 copper butterflies, without which less energy will be allocated to reproduction (Arrese and  
431 Soulages 2010, Karl et al. 2011). Both mechanisms result in a reduced consumption of storage  
432 reserves, at least initially (note here the young age of the butterflies tested). The above  
433 interpretation of an important impact of activity on storage reserves is further supported by the  
434 fact that the highest mean fat content was found in individuals being deprived of food under  
435 shady conditions, as a lack of solar radiation will certainly further reduce butterfly activity  
436 (Wickman 2009, Kuussaari et al. 2016). Furthermore, experiment 2 yielded identical results, as  
437 individuals released into sunny cages had on average a lower abdomen mass, a higher thorax-  
438 abdomen ratio, and a lower fat content. However, in experiment 1 the above patterns did not  
439 prevail, for which we have no explanation.

440

441 In experiments 1 and 2, sexual differences were more pronounced when individuals had access  
442 to food or when they were exposed to sunny conditions. The former indicates that females  
443 gained more mass when having access to food, probably as a result of egg maturation (Karl et

444 al. 2011). The latter indicates that males lost more mass and fat under sunny conditions than  
445 females, which may reflect their high flight activity (see above) under such beneficial  
446 conditions.

447

448 *Effects on emigration propensity*

449 A lack of resources needed for adult feeding and oviposition caused, as expected, a higher  
450 emigration propensity. Thus, individuals apparently showed a higher motivation to leave the  
451 departure patch in search for vital resources. Also, emigration propensity was higher under  
452 shady as compared with sunny conditions. This is intriguing since shade / lack of opportunity  
453 for sun basking will certainly reduce overall flight activity (e.g. Fischer and Fiedler 2001).  
454 Butterflies require high muscle temperatures between 30-38°C for flight activity, which is  
455 achieved through basking in the sun (e.g. Watt 1995, Wickman 2009). Taken together, these  
456 findings clearly suggest that emigration in *L. tityrus* is not merely driven by chance effects, but  
457 that butterflies can rather accurately assess patch quality and actively decide to leave a patch in  
458 case of degraded habitat quality. Such conditional, context-dependent behavior allows to escape  
459 from patches of unfavorable quality (Matthysen 2012). Accordingly, directed flight has been  
460 observed to occur more often in unsuitable habitats (Dennis 2004).

461

462 Overall, females were the more dispersive sex in *L. tityrus*. In experiment 3, females showed a  
463 clearly higher emigration propensity than males if resources were present. Several reasons may  
464 explain why females should indeed be more dispersive than males: (1) single mated females  
465 are, in contrast to males, able to found a new population, (2) females may emigrate more readily  
466 to escape from male harassment or as a risk spreading strategy by distributing their eggs over a  
467 wider area (Hill et al. 1999, Hopper 1999, Trochet et al. 2013). When being confronted with a  
468 lack of resources, both sexes increased emigration rates. However, this increase was much more

469 pronounced in males, causing an even higher emigration propensity under such conditions than  
470 in females. This is likely caused by males searching for appropriate sites to establish a territory,  
471 which typically cover dense patches of flowering nectar plants (Ebert and Rennwald 1991,  
472 Fischer and Fiedler 2001). The fact that male emigration was most strongly promoted when a  
473 lack of resources coincided with sunny conditions probably indicates that males took advantage  
474 of beneficial conditions for flight to search for territories. However, sex differences were not  
475 significant in experiments 1 and 2, but note that the principal pattern described above prevailed  
476 also in experiment 2 (not attaining significance) and to some extent even in experiment 1.

477

478 The impact of condition on emigration propensity though was remarkably low. Only in one out  
479 of 6 of our analyses (males in experiment 3), a PC (PC3) had a significant impact, indicating a  
480 positive association between wing aspect ratio and emigration propensity. Note that flight  
481 performance was only once included into the final models (though not attaining significance),  
482 reinforcing the above notion that emigration in *L. tityrus* is in the first place governed by  
483 motivation (i.e. context) rather than flight ability.

484

#### 485 *Variation in flight performance*

486 In two out of three experiments, males showed better flight performance than females. We  
487 cannot explain the results obtained in experiment 2, but assume that males indeed typically  
488 show a better flight performance than females. This would be in line with the above  
489 morphological differences between sexes. Furthermore, individuals released into shady  
490 compared with sunny departure cages had a better flight performance. Thus, conditions  
491 promoting flight activity diminish subsequent flight performance, presumably as a result of  
492 exhaustion and resource depletion. This is further supported by results from experiment 1, in  
493 which a lack of resources had a negative effect on flight performance.

494

495 Morphological traits had apparently a larger impact on flight performance than on emigration  
496 propensity (significant impact of PCs in 5 out of 6 analyses). This suggests that differences in  
497 flight performance between males and females can indeed be readily explained by  
498 morphological differences as has been repeatedly suggested (Van Dyck and Wiklund 2002,  
499 Berwaerts et al. 2006). In all three experiments, at least one PC turned out to be significant for  
500 males. Accordingly, smaller (lighter) males and those having a lower wing loading had a better  
501 flight performance. Additionally, lower abdomen masses increase the thorax-abdomen-ratio,  
502 which has often been found to enhance flight performance (see above). Wing aspect ratio, in  
503 contrast, showed no consistent association with flight performance. Regarding females, one PC  
504 turned out to be significant in two of the experiments. Flight performance was positively related  
505 to aspect ratio (experiment 1) and body mass (experiment 3). Here, a high body mass (with an  
506 accordingly high fat content to fuel flight) may reflect an overall good condition.

507

508

### *Conclusions*

509 Despite problems with high mortality restricting our observations to the first days of adult life,  
510 our results show some clear patterns which thus likely bear biological significance. Note that  
511 adult life span in copper butterflies is relatively short, typically ranging between 8-10 days on  
512 average in nature (Fischer et al. 1999). Based on experiments performed under semi-natural  
513 conditions, we obtained some interesting results. First, males and females differed substantially  
514 in morphology, with males showing traits typically associated with better flight performance,  
515 which we could verify in a flight performance test. Such differences most likely result from  
516 selection on males for increased acceleration speed, maneuverability and endurance in order to  
517 succeed in aerial combats with rivalling males and ultimately competition for females  
518 (Saastamoinen et al. 2012). Second and more interestingly, the traits underlying increased flight

519 performance (i.e. condition) did not affect emigration propensity. For instance, although  
520 females showed traits associated with diminished flight performance, they showed a higher  
521 emigration propensity than males (when resources were present), which can be readily  
522 explained in ecological and evolutionary contexts. This indicates that flight performance and  
523 the first step of dispersal, i.e. the actual decision whether to disperse or not, do not necessarily  
524 equate, casting doubt on the intuitive assumption of a general, positive relationship between the  
525 two. Note that the study design did not allow to examine the other stages of dispersal. Thus, we  
526 cannot assess the effects of flight performance on dispersal success. Third, environmental  
527 conditions (i.e. the context) affected condition, flight performance, but also emigration  
528 propensity. Notably, conditions indicative of poor habitat quality such as shade and a lack of  
529 resources promoted emigration. This suggests that emigration in *L. tityrus* is not merely a  
530 random process, but is governed by an active decision in a context-dependent manner,  
531 following an accurate assessment of patch quality. In summary, our results suggest that  
532 emigration propensity in this butterfly is a highly plastic, context-dependent trait triggered  
533 largely by habitat quality rather than by individual condition (Saastamoinen et al. 2012, Legrand  
534 et al. 2015). Besides abiotic factors, resource availability, and harassment, this may also involve  
535 density dependence as has been previously shown in other studies (Enfjäll and Leimar 2005,  
536 Trochet et al. 2013). Importantly, deteriorating habitat quality promotes emigration propensity  
537 (Legrand et al. 2015), which should be considered when trying to forecast future species  
538 distribution. For instance, climate change is likely to reduce habitat quality, but this will at the  
539 same time likely increase emigration propensity and thereby potentially facilitate range shifts  
540 in flying insects (Travis et al. 2013).

541

542

543

## ACKNOWLEDGEMENTS

544 We thank Delphine Legrand, Olivier Guillaume, and Olivier Calvez for technical support and  
545 helpful suggestions. This research was funded by the DFG research training group RESPONSE  
546 (DFG GRK 2010). MB was funded by the Agence Nationale de la Recherche (ANR) through  
547 the INDHET (open call, 2012) grant. MB is part of the Laboratoire d'Excellence TULIP (ANR-  
548 10-LABX-41).

549

550

551

## LITERATURE CITED

- 552 Arnold, S. J. 1983. Morphology, performance and fitness. *American Zoologist* 23:347–361.
- 553 Arrese, E. L., and J. L. Soulages. 2010. Insect fat body: energy, metabolism, and regulation.  
554 *Annual Review of Entomology* 55:207–225.
- 555 Baguette, M., J. Clobert, and N. Schtickzelle. 2011. Metapopulation dynamics of the bog  
556 fritillary butterfly: experimental changes in habitat quality induced negative density-  
557 dependent dispersal. *Ecography* 34:170–176.
- 558 Baguette, M., S. Blanchet, D. Legrand, V. M. Stevens, and C. Turlure. 2013. Individual  
559 dispersal, landscape connectivity and ecological networks. *Biological Reviews* 88:310–  
560 326.
- 561 Berg, M. P., E. Toby Kiers, G. Driessen, M. van der Heijden, B. W. Kooi, F. Kuenen, M.  
562 Liefting, H. A. Verhoef, and J. Ellers. 2010. Adapt or disperse: understanding species  
563 persistence in a changing world. *Global Change Biology* 16:587–598.
- 564 Berwaerts, K., P. Aerts, and H. Van Dyck. 2006. On the sex-specific mechanisms of butterfly  
565 flight: flight performance relative to flight morphology, wing kinematics, and sex in  
566 *Pararge aegeria*. *Biological Journal of the Linnean Society* 89:675–687.
- 567 Berwaerts, K., H. Van Dyck, and P. Aerts. 2002. Does flight morphology relate to flight

568 performance? An experimental test with the butterfly *Pararge aegeria*. *Functional*  
569 *Ecology* 16:484–491.

570 Blanckenhorn, W. U. 2000. The evolution of body size. What keeps organisms small? *The*  
571 *Quarterly Review of Biology* 75:385–407.

572 Bonte, D., H. Van Dyck, J. M. Bullock, A. Coulon, M. Delgado, M. Gibbs, V. Lehouck, E.  
573 Matthysen, K. Mustin, M. Saastamoinen, N. Schtickzelle, V. M. Stevens, S.  
574 Vandewoestijne, M. Baguette, K. Barton, T. G. Benton, A. Chaput-Bardy, J. Clobert, C.  
575 Dytham, T. Hovestadt, C. M. Meier, S. C. F. Palmer, C. Turlure, and J. M. J. Travis. 2012.  
576 Costs of dispersal. *Biological Reviews* 87:290–312.

577 Bowler, D. E., and T. G. Benton. 2005. Causes and consequences of animal dispersal strategies:  
578 relating individual behaviour to spatial dynamics. *Biological Reviews* 80:205–225.

579 Brunzel, S., M. Bussmann, and H. Obergruber. 2008. Deutliche Veränderungen von  
580 Tagfalterzönosen als Folge von Ausbreitungsprozessen. Erste Ergebnisse eines  
581 Monitorings über 17 Jahre. *Natur und Landschaft* 6:280–287.

582 Clobert, J., J. F. Le Galliard, J. Cote, S. Meylan, and M. Massot. 2009. Informed dispersal,  
583 heterogeneity in animal dispersal syndromes and the dynamics of spatially structured  
584 populations. *Ecology Letters* 12:197–209.

585 Clobert, J., M. Baguette, T. G. Benton, and J. M. Bullock. 2012. *Dispersal ecology and*  
586 *evolution*. Oxford University Press, Oxford, UK.

587 Cote, J., and J. Clobert. 2007. Social information and emigration: lessons from immigrants.  
588 *Ecology Letters* 10:411–417.

589 Dennis, R. L. H. 2004. Butterfly habitats, broad-scale biotope affiliations, and structural  
590 exploitation of vegetation at finer scales: the matrix revisited. *Ecological Entomology*  
591 29:744–752.

592 Ducatez, S., D. Legrand, A. Chaput-Bardy, V. M. Stevens, H. Fréville, and M. Baguette. 2012.

593 Inter-individual variation in movement: is there a mobility syndrome in the large white  
594 butterfly *Pieris brassicae*? *Ecological Entomology* 37:377–385.

595 Dudley, R. 2000. *The biomechanics of insect flight*. Princeton University Press, Princeton,  
596 USA.

597 Eaton, J. L. 1988. *Lepidopteran anatomy*. John Wiley and Sons, New York, USA.

598 Ebert, G., and E. Rennwald. 1991. *Die Schmetterlinge Baden-Württembergs*. Vol. 2. Ulmer,  
599 Stuttgart, Germany.

600 Enfjäll, K., and O. Leimar. 2005. Density-dependent dispersal in the Glanville fritillary,  
601 *Melitaea cinxia*. *Oikos* 108:465–472.

602 Fischer, K., B. Beinlich, and H. Plachter. 1999. Population structure, mobility and habitat  
603 preferences of the violet copper *Lycaena helle* (Lepidoptera: Lycaenidae) in Western  
604 Germany: implications for conservation. *Journal of Insect Conservation* 52:43–52.

605 Fischer, K., and K. Fiedler. 2001. Resource-based territoriality in the butterfly *Lycaena*  
606 *hippotoe* and environmentally induced behavioural shifts. *Animal Behaviour* 61:723–  
607 732.

608 Fischer, K., P. M. Brakefield, and B. J. Zwaan. 2003. Plasticity in butterfly egg size: why larger  
609 offspring at lower temperatures? *Ecology* 84:3138–3147.

610 Gilchrist, G. W. 1990. The consequences of sexual dimorphism in body size for butterfly flight  
611 and thermoregulation. *Functional Ecology* 4:475–487.

612 Hanski, I., C. J. Breuker, K. Schops, R. Setchfield, and M. Nieminen. 2002. Population history  
613 and life history influence the migration rate of female Glanville fritillary butterflies. *Oikos*  
614 98:87–97.

615 Hassall, C. 2015. Strong geographical variation in wing aspect ratio of a damselfly, *Calopteryx*  
616 *maculata* (Odonata: Zygoptera). *PeerJ* 3:e1219.

617 Hickling, R., D. B. Roy, J. K. Hill, R. Fox, and C. D. Thomas. 2006. The distributions of a wide

618 range of taxonomic groups are expanding polewards. *Global Change Biology* 12:450–455.

619 Hill, J. K., C. D. Thomas, and D. S. Blakeley. 1999. Evolution of flight morphology in a  
620 butterfly that has recently expanded its geographic range. *Oecologia* 121:165–170.

621 Honek, A. 1993. Intraspecific variation in body size and fecundity in insects: a general  
622 relationship. *Oikos* 66:483–492.

623 Hopper, K. R. 1999. Risk-spreading and bet-hedging in insect population biology. *Annual*  
624 *Review of Entomology* 44:535–560.

625 Karl, I., and K. Fischer. 2008. Why get big in the cold? Towards a solution to a life-history  
626 puzzle. *Oecologia* 155:215–225.

627 Karl, I., S. A. Janowitz, and K. Fischer. 2008. Altitudinal life-history variation and thermal  
628 adaptation in the copper butterfly *Lycaena tityrus*. *Oikos* 117:778–788.

629 Karl, I., R. Stoks, M. De Block, S. A. Janowitz, and K. Fischer. 2011. Temperature extremes  
630 and butterfly fitness: conflicting evidence from life history and immune function. *Global*  
631 *Change Biology* 17:676–687.

632 Kuussaari, M., S. Rytteri, R. K. Heikkinen, J. Helio, and P. von Bagh. 2016. Weather explains  
633 high annual variation in butterfly dispersal. *Proceedings of the Royal Society B*  
634 283:20160413.

635 Legrand, D., O. Guillaume, M. Baguette, J. Cote, A. Trochet, O. Calvez, S. Zajitschek, F.  
636 Zajitschek, J. Lecomte, Q. Bénard, J.-F. Le Galliard, and J. Clobert. 2012. The Metatron:  
637 an experimental system to study dispersal and metaecosystems for terrestrial organisms.  
638 *Nature Methods* 9:828–833.

639 Legrand, D., A. Trochet, S. Moulherat, O. Calvez, V. M. Stevens, S. Ducatez, J. Clobert, and  
640 M. Baguette. 2015. Ranking the ecological causes of dispersal in a butterfly. *Ecography*  
641 38:822–831.

642 Legrand, D., N. Larranaga, R. Bertrand, S. Ducatez, O. Calvez, V. M. Stevens, and M. Baguette.

643 2016. Evolution of a butterfly dispersal syndrome. *Proceedings of the Royal Society B:*  
644 *Biological Sciences* 283:20161533.

645 Matthysen, E. 2012. Multicausality of dispersal: a review. Pages 1–18 in J. Clobert, M.  
646 Baguette, T. G. Benton, J. M. Bullock, editors. *Dispersal Ecology and Evolution*. Oxford  
647 University Press, Oxford, UK.

648 Nathan, R., W. M. Getz, E. Revilla, M. Holyoak, R. Kadmon, D. Saltz, and P. E. Smouse. 2008.  
649 A movement ecology paradigm for unifying organismal movement research. *Proceedings*  
650 *of the National Academy of Sciences* 105:19052–19059.

651 Niitepõld, K., A. D. Smith, J. L. Osborne, D. R. Reynolds, N. L. Carreck, A. P. Martin, J. H.  
652 Marden, O. Ovaskainen, and I. Hanski. 2009. Flight metabolic rate and *pgi* genotype  
653 influence butterfly dispersal rate in the field. *Ecology* 90:2223–2232.

654 Ronce, O. 2007. How does it feel to be like a rolling stone? Ten questions about dispersal  
655 evolution. *Annual Review of Ecology, Evolution, and Systematics* 38:231–253.

656 Saastamoinen, M., and I. Hanski. 2008. Genotypic and environmental effects on flight activity  
657 and oviposition in the Glanville fritillary butterfly. *The American Naturalist* 171:701–712.

658 Saastamoinen, M., D. van der Sterren, N. Vastenhout, B. J. Zwaan, and P. M. Brakefield. 2010.  
659 Predictive adaptive responses: condition-dependent impact of adult nutrition and flight in  
660 the tropical butterfly *Bicyclus anynana*. *The American Naturalist* 176:686–698.

661 Saastamoinen, M., P. M. Brakefield, and O. Ovaskainen. 2012. Environmentally induced  
662 dispersal-related life-history syndrome in the tropical butterfly, *Bicyclus anynana*. *Journal*  
663 *of Evolutionary Biology* 25:2264–2275.

664 Sekar, S. 2012. A meta-analysis of the traits affecting dispersal ability in butterflies: can  
665 wingspan be used as a proxy? *Journal of Animal Ecology* 81:174–184.

666 Settele, J., O. Kudrna, A. Harpke, I. Kühn, and C. Van Swaay. 2008. Climatic risk atlas of  
667 European butterflies. *Biodiversity and ecosystem risk assessment 1*. Pensoft, Sofia,

668 Bulgaria.

669 Tammaru, T., P. Kaitaniemi, and K. Ruohomaki. 1996. Realized fecundity in *Epirrita*  
670 *autumnata* (Lepidoptera: Geometridae): relation to body size and consequences to  
671 population dynamics. *Oikos* 77:407–416.

672 Tolman, T., and R. Lewington. 1998. *Die Tagfalter Europas und Nordwestafrikas*. Franckh-  
673 Kosmos, Stuttgart, Germany.

674 Travis, J. M. J., M. Delgado, G. Bocedi, M. Baguette, K. Bartoń, D. Bonte, I. Boulangeat, J. A.  
675 Hodgson, A. Kubisch, V. Penteriani, M. Saastamoinen, V. M. Stevens, and J. M. Bullock.  
676 2013. Dispersal and species' responses to climate change. *Oikos* 122:1532–1540.

677 Trochet, A., D. Legrand, N. Larranaga, S. Ducatez, O. Calvez, J. Cote, J. Clobert, and M.  
678 Baguette. 2013. Population sex ratio and dispersal in experimental, two-patch  
679 metapopulations of butterflies. *Journal of Animal Ecology* 82:946–955.

680 Turlure, C., N. Schtickzelle, H. Van Dyck, B. Seymoure, and R. Rutowski. 2016. Flight  
681 morphology, compound eye structure and dispersal in the bog and the cranberry fritillary  
682 butterflies: An inter- and intraspecific comparison. *PLOS ONE* 11:e0158073.

683 Van Dyck, H., and C. Wiklund. 2002. Seasonal butterfly design: morphological plasticity  
684 among three developmental pathways relative to sex, flight and thermoregulation. *Journal*  
685 *of Evolutionary Biology* 15:216–225.

686 Warren, M. S., J. K. Hill, J. A. Thomas, J. Asher, R. Fox, B. Huntley, D. B. Roy, M. G. Telfer,  
687 S. Jeffcoate, P. Harding, G. Jeffcoate, S. G. Willis, J. N. Greatorex-Davies, D. Moss, and  
688 C. D. Thomas. 2001. Rapid responses of British butterflies to opposing forces of climate  
689 and habitat change. *Nature* 414:65–69.

690 Watt, W. B. 1995. Adaptive significance of pigment polymorphism in *Colias* butterflies. I.  
691 Variation of melanin pigment in relation to thermoregulation. *Evolution* 22:437–458.

692 Wickman, P. O. 2009. Thermoregulation and habitat use in butterflies. Pages 55–61 in J.

693 Settele, T. G. Shreeve, M. Konvicka, and H. Van Dyck, editors. Ecology of Butterflies in  
694 Europe. Cambridge University Press, Cambridge, UK.

695 Wiklund, C., and T. Fagerström. 1977. Why do males emerge before females? *Oecologia*  
696 31:153–158.

697 Zera, A. J., J. Potts, and K. Kobus. 1998. The physiology of life-history trade-offs: experimental  
698 analysis of a hormonally induced life-history trade-off in *Gryllus assimilis*. *The American*  
699 *Naturalist* 152:7–23.

700

701

## TABLES

702 TABLE 1. Morphological traits of *Lycaena tityrus* in relation to sex and the presence of  
 703 resources in the departure cage. Given are means  $\pm$  1 SE. Group sample sizes are 16 for fed  
 704 males, 7 for unfed males, 31 for fed females and 34 for unfed females. Different superscript  
 705 letters indicate significant differences among groups.

706

Trait	Resources		No resources	
	Males	Females	Males	Females
Adult body mass (mg)	29.2 $\pm$ 0.98 <sup>a</sup>	37.3 $\pm$ 1.20 <sup>b</sup>	28.1 $\pm$ 3.04 <sup>a</sup>	32.4 $\pm$ 1.45 <sup>a</sup>
Thorax mass (mg)	15.2 $\pm$ 0.62 <sup>a</sup>	14.3 $\pm$ 0.40 <sup>a</sup>	12.6 $\pm$ 1.10 <sup>a</sup>	13.2 $\pm$ 0.57 <sup>a</sup>
Abdomen mass (mg)	7.9 $\pm$ 0.76 <sup>a</sup>	16.7 $\pm$ 0.71 <sup>b</sup>	10.3 $\pm$ 2.71 <sup>ac</sup>	13.3 $\pm$ 0.82 <sup>c</sup>
Thorax-abdomen ratio	2.1 $\pm$ 0.20 <sup>a</sup>	0.9 $\pm$ 0.03 <sup>b</sup>	1.7 $\pm$ 0.34 <sup>a</sup>	1.0 $\pm$ 0.04 <sup>b</sup>
Fat (mg)	0.4 $\pm$ 0.10 <sup>ab</sup>	0.5 $\pm$ 0.04 <sup>b</sup>	0.3 $\pm$ 0.16 <sup>ab</sup>	0.3 $\pm$ 0.04 <sup>a</sup>
Fat (%)	13.3 $\pm$ 2.67 <sup>a</sup>	8.8 $\pm$ 0.60 <sup>ab</sup>	12.9 $\pm$ 4.27 <sup>a</sup>	5.5 $\pm$ 0.56 <sup>b</sup>
Wing length (mm)	14.7 $\pm$ 0.17 <sup>a</sup>	14.8 $\pm$ 0.13 <sup>a</sup>	14.2 $\pm$ 0.39 <sup>a</sup>	14.9 $\pm$ 0.14 <sup>a</sup>
Wing area (mm <sup>2</sup> )	80.8 $\pm$ 1.85 <sup>a</sup>	88.1 $\pm$ 1.51 <sup>b</sup>	75.7 $\pm$ 4.20 <sup>a</sup>	89.1 $\pm$ 1.62 <sup>b</sup>
Wing loading	36.2 $\pm$ 1.08 <sup>a</sup>	42.3 $\pm$ 1.01 <sup>b</sup>	37.7 $\pm$ 4.55 <sup>ab</sup>	36.1 $\pm$ 1.21 <sup>a</sup>
Wing aspect ratio	10.7 $\pm$ 0.10 <sup>a</sup>	10.0 $\pm$ 0.05 <sup>b</sup>	10.7 $\pm$ 0.13 <sup>a</sup>	10.0 $\pm$ 0.05 <sup>b</sup>

707

708 TABLE 2. Results of generalized linear models for the effects of resources, sex, and family on  
 709 (a) emigration propensity (logit-link function and binomial error distribution) and (b) flight  
 710 performance (flight duration in a vortex; log-link function and normal error distribution) in the  
 711 butterfly *Lycaena tityrus* (models 1). Second models were constructed separately for males and  
 712 females by additionally including principal components (PCs) as covariates. All models were  
 713 constructed by a stepwise backwards elimination of non-significant factors. Significant p-  
 714 values are given in bold.

715

716 TABLE 2a

<b>Model 1</b>	<b>d.f.</b>	<b>Odds ratio</b>	<b>95% CI</b>	<b>Wald Chi<sup>2</sup></b>	<b>p</b>
Resources	1	0.280	0.010-0.784	5.9	<b>0.0154</b>
<b>Model 2 Males</b>	<b>d.f.</b>	<b>Odds ratio</b>	<b>95% CI</b>	<b>Wald Chi<sup>2</sup></b>	<b>p</b>
Flight performance	1	1.121	0.985-1.276	3.0	0.0837
Resources	1	0.005	< 0.001-1.674	3.2	0.0737
<b>Model 2 Females</b>	<b>d.f.</b>	<b>Odds ratio</b>	<b>95% CI</b>	<b>Wald Chi<sup>2</sup></b>	<b>p</b>
Resources	1	0.360	0.107-1.215	2.7	0.0996

717

718 TABLE 2b

<b>Model 1</b>	<b>d.f.</b>	<b>Parameter estimate</b>	<b>SE</b>	<b>Wald Chi<sup>2</sup></b>	<b>p</b>
Resources	1	-0.25	0.09	8.7	<b>0.0032</b>
Sex	1	0.18	0.07	6.5	<b>0.0109</b>
Family	6			11.3	0.0808
<b>Model 2 Males</b>	<b>d.f.</b>	<b>Parameter estimate</b>	<b>SE</b>	<b>Wald Chi<sup>2</sup></b>	<b>p</b>
Resources	1	-0.24	0.10	5.5	<b>0.0195</b>
PC2	1	-0.26	0.08	10.8	<b>0.0010</b>
Family	5			21.0	<b>0.0008</b>
<b>Model 2 Females</b>	<b>d.f.</b>	<b>Parameter estimate</b>	<b>SE</b>	<b>Wald Chi<sup>2</sup></b>	<b>p</b>
Resources	1	-0.43	0.13	11.3	<b>0.0008</b>
PC4	1	-0.16	0.08	3.8	<b>0.0498</b>

719

720 TABLE 3. Morphological traits of *Lycaena tityrus* in relation to sex and sunny versus shady  
 721 conditions in the departure cage. Given are means  $\pm$  1 SE. Group sample size is 11 for males  
 722 and 31 for females at sunny, and 8 for males and 31 for females at shady conditions. Different  
 723 superscript letters indicate significant differences among groups.

724

Trait	Sun		Shade	
	Males	Females	Males	Females
Adult body mass (mg)	27.8 $\pm$ 0.61 <sup>a</sup>	40.0 $\pm$ 1.53 <sup>b</sup>	32.2 $\pm$ 2.68 <sup>ab</sup>	39.4 $\pm$ 1.94 <sup>b</sup>
Thorax mass (mg)	14.9 $\pm$ 0.49 <sup>a</sup>	14.8 $\pm$ 0.54 <sup>a</sup>	13.7 $\pm$ 0.90 <sup>a</sup>	14.3 $\pm$ 0.59 <sup>a</sup>
Abdomen mass (mg)	7.0 $\pm$ 0.74 <sup>a</sup>	19.2 $\pm$ 1.00 <sup>b</sup>	13.0 $\pm$ 2.28 <sup>ab</sup>	19.2 $\pm$ 1.32 <sup>b</sup>
Thorax-abdomen ratio	2.3 $\pm$ 0.22 <sup>a</sup>	0.8 $\pm$ 0.03 <sup>b</sup>	1.4 $\pm$ 0.30 <sup>c</sup>	0.8 $\pm$ 0.05 <sup>b</sup>
Fat (mg)	0.2 $\pm$ 0.07 <sup>a</sup>	1.0 $\pm$ 0.15 <sup>b</sup>	1.2 $\pm$ 0.26 <sup>b</sup>	1.1 $\pm$ 0.11 <sup>b</sup>
Fat (%)	11.6 $\pm$ 2.86 <sup>a</sup>	15.2 $\pm$ 1.43 <sup>a</sup>	29.1 $\pm$ 3.24 <sup>b</sup>	18.4 $\pm$ 1.30 <sup>a</sup>
Wing length (mm)	14.9 $\pm$ 0.14 <sup>a</sup>	14.8 $\pm$ 0.12 <sup>a</sup>	14.5 $\pm$ 0.15 <sup>a</sup>	14.8 $\pm$ 0.13 <sup>a</sup>
Wing area (mm <sup>2</sup> )	82.9 $\pm$ 1.49 <sup>ab</sup>	88.5 $\pm$ 1.64 <sup>b</sup>	77.4 $\pm$ 1.66 <sup>a</sup>	88.8 $\pm$ 1.57 <sup>b</sup>
Wing loading	33.7 $\pm$ 1.08 <sup>a</sup>	45.0 $\pm$ 1.30 <sup>b</sup>	41.5 $\pm$ 2.97 <sup>ab</sup>	43.8 $\pm$ 1.65 <sup>b</sup>
Wing aspect ratio	10.8 $\pm$ 0.06 <sup>a</sup>	9.9 $\pm$ 0.03 <sup>b</sup>	10.9 $\pm$ 0.12 <sup>a</sup>	9.9 $\pm$ 0.05 <sup>b</sup>

725

726 TABLE 4. Results of generalized linear models for the effects of sun, sex and family on (a)  
727 emigration propensity (logit-link function and binomial error distribution) and (b) flight  
728 performance (flight duration in a vortex; log-link function and normal error distribution) in the  
729 butterfly *Lycaena tityrus* (model 1). Second models were constructed separately for males and  
730 females by additionally including principal components (PCs) as covariates. All models were  
731 constructed by a stepwise backwards elimination of non-significant factors. For (a), no PC was  
732 included for females and no factor at all was selected for males within the second models. For  
733 (b), no factor was selected in model 1 and in model 2 for females. Significant p-values are given  
734 in bold.

735

736 TABLE 4a

<b>Model 1</b>	<b>d.f.</b>	<b>Odds ratio</b>	<b>95% CI</b>	<b>Wald Chi<sup>2</sup></b>	<b>p</b>
Sun	1	0.086	0.019-0.399	9.8	<b>0.0017</b>
<b>Model 2 Females</b>	<b>d.f.</b>	<b>Odds ratio</b>	<b>95% CI</b>	<b>Wald Chi<sup>2</sup></b>	<b>p</b>
Sun	1	0.111	0.023-0.544	7.4	<b>0.0067</b>

737

738 TABLE 4b

<b>Model 2 Males</b>	<b>d.f.</b>	<b>Parameter estimate</b>	<b>SE</b>	<b>Wald Chi<sup>2</sup></b>	<b>p</b>
PC1	1	-0.37	0.16	5.6	<b>0.0181</b>
PC2	1	-0.43	0.22	4.1	<b>0.0437</b>
PC3	1	0.68	0.28	6.3	<b>0.0122</b>
PC4	1	0.88	0.42	4.5	<b>0.0343</b>
Sun	1	0.33	0.20	2.8	0.0956
Family	5			9.3	0.0980

739

740 TABLE 5. Morphological traits of *Lycaena tityrus* in relation to sex. Given are means  $\pm$  1 SE.  
 741 Group sample size is 24 for males and 49 for females. Different superscript letters indicate  
 742 significant differences among groups.

743

<b>Trait</b>	<b>Males</b>	<b>Females</b>
Adult body mass (mg)	27.9 $\pm$ 0.98 <sup>a</sup>	37.2 $\pm$ 1.31 <sup>b</sup>
Thorax mass (mg)	14.9 $\pm$ 0.06 <sup>a</sup>	14.3 $\pm$ 0.04 <sup>a</sup>
Abdomen mass (mg)	7.1 $\pm$ 0.04 <sup>a</sup>	16.9 $\pm$ 0.09 <sup>b</sup>
Thorax-abdomen ratio	2.3 $\pm$ 0.14 <sup>a</sup>	0.9 $\pm$ 0.04 <sup>b</sup>
Fat (mg)	0.2 $\pm$ 0.04 <sup>a</sup>	0.6 $\pm$ 0.06 <sup>b</sup>
Fat (%)	11.2 $\pm$ 1.41 <sup>a</sup>	10.5 $\pm$ 0.91 <sup>b</sup>
Wing length (mm)	14.5 $\pm$ 0.16 <sup>a</sup>	14.8 $\pm$ 0.11 <sup>a</sup>
Wing area (mm <sup>2</sup> )	78.5 $\pm$ 1.70 <sup>a</sup>	86.8 $\pm$ 1.19 <sup>b</sup>
Wing loading	35.6 $\pm$ 0.87 <sup>a</sup>	42.3 $\pm$ 1.35 <sup>b</sup>
Wing aspect ratio	10.8 $\pm$ 0.07 <sup>a</sup>	9.9 $\pm$ 0.04 <sup>b</sup>

744

745 TABLE 6. Results of generalized linear models for the effects of resources, sun, sex, family  
 746 and age on (a) emigration propensity (logit-link function and binomial error distribution) and  
 747 (b) flight performance (flight duration in a vortex; log-link function and normal error  
 748 distribution) in the butterfly *Lycaena tityrus* (models 1). Second models were constructed  
 749 separately for males and females by additionally including principal components (PCs) as  
 750 covariates. All models were constructed by a stepwise backwards elimination of non-significant  
 751 factors. Significant p-values are given in bold.

752

753 TABLE 6a

<b>Model 1</b>	<b>Odds ratio</b>	<b>95% CI</b>	<b>d.f.</b>	<b>Wald Chi<sup>2</sup></b>	<b>p</b>
Resources	0.222	0.061-0.811	1	5.2	<b>0.0227</b>
Resources * Sex			1	5.3	<b>0.0217</b>
Resources * Sun * Sex			1	5.6	<b>0.0183</b>
<b>Model 2 Males</b>	<b>Odds ratio</b>	<b>95% CI</b>	<b>d.f.</b>	<b>Wald Chi<sup>2</sup></b>	<b>p</b>
PC3	0.259	0.067-1.003	1	3.8	0.0505
<b>Model 2 Females</b>	<b>Odds ratio</b>	<b>95% CI</b>	<b>d.f.</b>	<b>Wald Chi<sup>2</sup></b>	<b>p</b>
Resources * Sun			1	5.1	<b>0.0243</b>

754

755 TABLE 6b

<b>Model 1</b>	<b>Parameter estimate</b>	<b>SE</b>	<b>d.f.</b>	<b>Wald Chi<sup>2</sup></b>	<b>p</b>
Sun	0.22	0.09	1	5.9	<b>0.0153</b>
Sex	0.28	0.09	1	8.9	<b>0.0028</b>
Family			5	12.1	<b>0.0341</b>
<b>Model 2 Males</b>	<b>Parameter estimate</b>	<b>SE</b>	<b>d.f.</b>	<b>Wald Chi<sup>2</sup></b>	<b>p</b>
PC3	-0.62	0.28	1	4.9	<b>0.0276</b>
Family			4	8.4	0.0794
<b>Model 2 Females</b>	<b>Parameter estimate</b>	<b>SE</b>	<b>d.f.</b>	<b>Wald Chi<sup>2</sup></b>	<b>p</b>
PC1	-0.10	0.05	1	3.9	<b>0.0470</b>

756

FIGURE LEGENDS

757

758

759 FIGURE 1. Emigration propensity (%), a) and flight performance (means + 1 SE, b) of *Lycaena*  
760 *tityrus* in relation to the presence or absence of resources in the departure cage. Males: filled  
761 bars; females: open bars.

762


763 FIGURE 2. Emigration propensity (%), a) and flight performance (means + 1 SE, b) of *Lycaena*  
764 *tityrus* in relation to having experienced sunny or shady conditions in the departure cage. Males:  
765 filled bars; females: open bars.

766

767 FIGURE 3. Emigration propensity (%), a) in relation to the presence or absence of resources  
768 and flight performance (means + 1 SE, b) in relation to having experienced sunny or shady  
769 conditions in the departure cage in *Lycaena tityrus*. Males: filled bars; females: open bars.


770

771 FIG. 1a




772

773 FIG. 1b




774

775 FIG. 2a




776

777 FIG. 2b




778

779 FIG. 3a



780

781 FIG. 3b



782

APPENDIX S1

TABLE S1. Correlations among morphological parameters in *L. tityrus* for experiment 1 (a), 2 (b) and 3 (c). Significant p-values are given in bold.

TABLE S1a

	Adult body mass	Abdomen mass	Thorax mass	Fat (mg)	Fat (%)	Wing length	Wing area	Thorax-abdomen ratio	Wing loading	Wing aspect ratio
Adult body mass	1.000	<b>0.904</b>	<b>0.742</b>	<b>0.656</b>	<b>0.239</b>	<b>0.549</b>	<b>0.614</b>	<b>-0.439</b>	<b>0.869</b>	<b>-0.296</b>
Abdomen mass	<b>0.904</b>	1.000	<b>0.394</b>	<b>0.655</b>	0.185	<b>0.316</b>	<b>0.438</b>	<b>-0.740</b>	<b>0.859</b>	<b>-0.383</b>
Thorax mass	<b>0.742</b>	<b>0.394</b>	1.000	<b>0.404</b>	<b>0.266</b>	<b>0.647</b>	<b>0.591</b>	0.214	<b>0.561</b>	-0.017
Fat (mg)	<b>0.656</b>	<b>0.655</b>	<b>0.404</b>	1.000	<b>0.791</b>	<b>0.221</b>	<b>0.231</b>	<b>-0.357</b>	<b>0.686</b>	-0.067
Fat (%)	<b>0.239</b>	0.185	<b>0.266</b>	<b>0.791</b>	1.000	0.011	-0.114	0.083	<b>0.396</b>	<b>0.341</b>
Wing length	<b>0.549</b>	<b>0.316</b>	<b>0.647</b>	<b>0.221</b>	0.010	1.000	<b>0.930</b>	0.059	0.120	-0.073
Wing area	<b>0.614</b>	<b>0.438</b>	<b>0.591</b>	<b>0.232</b>	-0.114	<b>0.930</b>	1.000	-0.129	0.153	<b>-0.431</b>
Thorax-abdomen ratio	<b>-0.439</b>	<b>-0.740</b>	0.214	<b>-0.357</b>	0.083	0.059	-0.129	1.000	<b>-0.477</b>	<b>0.470</b>
Wing loading	<b>0.869</b>	<b>0.859</b>	<b>0.561</b>	<b>0.686</b>	<b>0.396</b>	0.121	0.153	<b>-0.477</b>	1.000	-0.097
Wing aspect ratio	<b>-0.296</b>	<b>-0.383</b>	-0.017	-0.067	<b>0.341</b>	-0.073	<b>-0.431</b>	<b>0.470</b>	-0.097	1.000

TABLE S1b

	Adult body mass	Abdomen mass	Thorax mass	Fat (mg)	Fat (%)	Wing length	Wing area	Thorax-abdomen ratio	Wing loading	Wing aspect ratio
Adult body mass	1.000	<b>0.946</b>	<b>0.737</b>	<b>0.456</b>	0.088	<b>0.581</b>	<b>0.727</b>	<b>-0.546</b>	<b>0.933</b>	<b>-0.493</b>
Abdomen mass	<b>0.946</b>	1.000	<b>0.484</b>	<b>0.543</b>	0.187	<b>0.436</b>	<b>0.638</b>	<b>-0.742</b>	<b>0.906</b>	<b>-0.570</b>
Thorax mass	<b>0.737</b>	<b>0.485</b>	1.000	0.118	-0.139	<b>0.658</b>	<b>0.637</b>	0.059	<b>0.640</b>	-0.155
Fat (mg)	<b>0.456</b>	<b>0.543</b>	0.118	1.000	<b>0.753</b>	<b>0.270</b>	<b>0.327</b>	<b>-0.482</b>	<b>0.427</b>	-0.197
Fat (%)	0.088	0.187	-0.139	<b>0.753</b>	1.000	0.075	0.001	-0.190	0.104	0.173
Wing length	<b>0.581</b>	<b>0.436</b>	<b>0.658</b>	<b>0.270</b>	0.075	1.000	<b>0.896</b>	< 0.001	<b>0.301</b>	-0.083
Wing area	<b>0.727</b>	<b>0.638</b>	<b>0.637</b>	<b>0.327</b>	0.001	<b>0.896</b>	1.000	<b>-0.274</b>	<b>0.438</b>	<b>-0.514</b>
Thorax-abdomen ratio	<b>-0.546</b>	<b>-0.742</b>	0.059	<b>-0.482</b>	-0.190	< 0.001	<b>-0.274</b>	1.000	<b>-0.583</b>	<b>0.603</b>
Wing loading	<b>0.933</b>	<b>0.906</b>	<b>0.640</b>	<b>0.427</b>	0.104	<b>0.301</b>	<b>0.438</b>	<b>-0.583</b>	1.000	<b>-0.383</b>
Wing aspect ratio	<b>-0.493</b>	<b>-0.570</b>	-0.155	-0.197	0.173	-0.083	<b>-0.514</b>	<b>0.603</b>	<b>-0.383</b>	1.000

TABLE S1c

	Adult body mass	Abdomen mass	Thorax mass	Fat (mg)	Fat (%)	Wing length	Wing area	Thorax-abdomen ratio	Wing loading	Wing aspect ratio
Adult body mass	1.000	<b>0.917</b>	<b>0.665</b>	<b>0.534</b>	0.038	<b>0.454</b>	<b>0.589</b>	<b>-0.521</b>	<b>0.927</b>	<b>-0.379</b>
Abdomen mass	<b>0.917</b>	1.000	<b>0.319</b>	<b>0.598</b>	0.076	<b>0.261</b>	<b>0.455</b>	<b>-0.762</b>	<b>0.891</b>	<b>-0.479</b>
Thorax mass	<b>0.665</b>	<b>0.319</b>	1.000	0.145	-0.045	<b>0.540</b>	<b>0.495</b>	0.183	<b>0.567</b>	0.009
Fat (mg)	<b>0.534</b>	<b>0.598</b>	0.145	1.000	<b>0.699</b>	0.134	0.225	<b>-0.464</b>	<b>0.542</b>	<b>-0.236</b>
Fat (%)	0.038	0.076	-0.045	<b>0.699</b>	1.000	-0.144	-0.208	-0.013	0.139	0.163
Wing length	<b>0.454</b>	<b>0.261</b>	<b>0.540</b>	0.134	-0.144	1.000	<b>0.893</b>	0.002	0.135	0.056
Wing area	<b>0.589</b>	<b>0.455</b>	<b>0.495</b>	0.225	-0.208	<b>0.893</b>	1.000	<b>-0.254</b>	<b>0.249</b>	<b>-0.396</b>
Thorax-abdomen ratio	<b>-0.521</b>	<b>-0.762</b>	0.183	<b>-0.464</b>	-0.013	0.002	<b>-0.254</b>	1.000	<b>-0.517</b>	<b>0.571</b>
Wing loading	<b>0.927</b>	<b>0.891</b>	<b>0.567</b>	<b>0.543</b>	0.139	0.135	<b>0.249</b>	<b>-0.517</b>	1.000	<b>-0.277</b>
Wing aspect ratio	<b>-0.379</b>	<b>-0.479</b>	0.009	<b>-0.236</b>	0.163	0.056	<b>-0.396</b>	<b>0.571</b>	<b>-0.277</b>	1.000

TABLE S2. Principal component analysis (PCA) results for males (a) and females (b) including Eigenvalues, percentage of the variance explained, cumulative percentage of the variance explained, and r-values of correlations with continuous variables (rows 4-13) for principal components 1-4. r-values > 0.7 in bold.

TABLE S2a

No.	Parameter	PC1	PC2	PC3	PC4
1	Eigenvalue	4.1	3.0	1.5	0.7
2	Variance explained (%)	41.2	30.0	15.2	7.0
3	Cumulative variance (%)	41.2	71.2	86.4	93.4
4	Adult body mass	0.697	0.624	-0.320	0.126
5	Thorax mass	-0.151	<b>0.883</b>	-0.256	0.052
6	Abdomen mass	<b>0.953</b>	0.052	-0.174	0.096
7	Thorax-abdomen ratio	<b>-0.818</b>	0.354	0.071	-0.211
8	Fat (mg)	<b>0.811</b>	0.157	0.416	-0.333
9	Fat (%)	0.687	0.210	0.536	-0.388
10	Wing length	-0.198	<b>0.865</b>	0.359	0.184
11	Wing area	-0.264	<b>0.912</b>	0.124	0.007
12	Wing loading	<b>0.869</b>	0.134	-0.397	0.135
13	Wing aspect ratio	0.203	-0.188	<b>0.743</b>	0.563

TABLE S2b

No.	Parameter	PC1	PC2	PC3	PC4
1	Eigenvalue	5.1	2.0	1.1	1.0
2	Variance explained (%)	50.7	20.2	11.1	10.2
3	Cumulative variance (%)	50.7	70.9	82.0	92.2
4	Adult body mass	<b>-0.974</b>	0.094	0.180	-0.041
5	Thorax mass	<b>-0.796</b>	0.399	0.112	-0.030
6	Abdomen mass	<b>-0.952</b>	-0.129	0.212	-0.038
7	Thorax-abdomen ratio	0.628	0.491	-0.271	0.055
8	Fat (mg)	-0.636	-0.534	-0.495	0.058
9	Fat (%)	-0.427	-0.699	-0.509	0.072
10	Wing length	-0.617	0.634	-0.393	-0.130
11	Wing area	-0.656	0.623	-0.342	0.149
12	Wing loading	<b>-0.874</b>	-0.166	0.385	-0.133
13	Wing aspect ratio	0.171	-0.033	-0.155	<b>-0.972</b>

TABLE S3. Results of general linear mixed models for the effects of resources, sex (fixed factors), and family (random) on morphological traits in *Lycaena tityrus* for individuals released in experiment 1. Models were constructed using a stepwise backwards elimination of non-significant factors. Significant p-values are given in bold.

<b>Adult body mass</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sex	3.9 * 10 <sup>-4</sup>	1	8.9	<b>0.0037</b>
Resources * Sex	1.4 * 10 <sup>-4</sup>	2	3.1	<b>0.0493</b>
Family	0.9 * 10 <sup>-4</sup>	6	2.1	0.0595
Error	0.4 * 10 <sup>-4</sup>	78		
<b>Thorax mass</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Resources	0.31 * 10 <sup>-4</sup>	1	5.0	<b>0.0275</b>
Family	0.22 * 10 <sup>-4</sup>	6	3.5	<b>0.0040</b>
Error	0.06 * 10 <sup>-4</sup>	80		
<b>Abdomen mass</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sex	5.6 * 10 <sup>-4</sup>	1	30.6	<b>&lt;0.0001</b>
Resources * Sex	2.2 * 10 <sup>-4</sup>	1	11.9	<b>0.0009</b>
Error	0.2 * 10 <sup>-4</sup>	85		
<b>Thorax-abdomen ratio</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sex	14.9	1	86.0	<b>&lt;0.0001</b>
Resources * Sex	0.6	2	3.5	<b>0.0357</b>
Family	0.3	6	1.8	0.1076
Error	0.2	78		

Table S3 continued

<b>Fat (mg)</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Resources * Sex	0.31	3	4.4	<b>0.0063</b>
Family	0.20	6	2.8	<b>0.0151</b>
Error	0.07	78		
<b>Fat (%)</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sex	373	1	12.8	<b>0.0006</b>
Resources * Sex	77	2	2.7	0.0767
Family	114	6	3.9	<b>0.0018</b>
Error	29	78		
<b>Wing length</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Family	3.6	6	10.3	<b>&lt;0.0001</b>
Error	0.3	77		
<b>Wing area</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sex	855	1	20.4	<b>&lt;0.0001</b>
Family	496	6	11.8	<b>&lt;0.0001</b>
Error	42	76		
<b>Wing loading</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Resources * Sex	5.8	1	13.5	<b>0.0004</b>
Error	0.4	82		

Table S3 continued

<b>Wing aspect ratio</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sex	6.54	1	76.9	<b>&lt;0.0001</b>
Family	0.19	6	2.2	0.0545
Error	0.09	76		

TABLE S4. Results of general linear mixed models for the effects of sun, sex (fixed factors), and family (random) on morphological traits in *Lycaena tityrus* for individuals released in experiment 2. Models were constructed using stepwise backwards elimination of non-significant factors. Significant p-values are given in bold.

<b>Adult body mass</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sex	12.2 * 10 <sup>-4</sup>	1	15.4	<b>0.0002</b>
Family	1.7 * 10 <sup>-4</sup>	6	2.2	0.0525
Error	0.8 * 10 <sup>-4</sup>	80		
<b>Thorax mass</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Family	0.2 * 10 <sup>-4</sup>	6	2.4	<b>0.0373</b>
Error	0.1 * 10 <sup>-4</sup>	81		
<b>Abdomen mass</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sun	1.6 * 10 <sup>-4</sup>	1	4.1	<b>0.0452</b>
Sex	9.7 * 10 <sup>-4</sup>	1	25.3	<b>&lt;0.0001</b>
Sun * Sex	1.7 * 10 <sup>-4</sup>	1	4.5	<b>0.0365</b>
Family	0.7 * 10 <sup>-4</sup>	6	1.9	0.1000
Error	0.4 * 10 <sup>-4</sup>	78		

Table S4 continued

<b>Thorax-abdomen ratio</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sun	3.26	1	20.3	<b>&lt;0.0001</b>
Sex	12.93	1	80.6	<b>&lt;0.0001</b>
Sun * Sex	3.44	1	21.4	<b>&lt;0.0001</b>
Family	0.23	6	1.5	0.2021
Error	0.16	78		
<b>Fat (mg)</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sun	2.17	1	4.2	<b>0.0445</b>
Sex	2.91	1	5.6	<b>0.0203</b>
Error	0.52	85		
<b>Fat (%)</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sun	1509	1	22.7	<b>&lt;0.0001</b>
Sun * Sex	647	1	9.7	<b>0.0025</b>
Error	67	85		
<b>Wing length</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Family	1.28	6	3.5	<b>0.0042</b>
Error	0.37	76		
<b>Wing area</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sex	735.1	1	13.3	<b>0.0005</b>
Family	249.6	6	4.5	<b>0.0006</b>
Error	55.4	75		

Table S4 continued

<b>Wing loading</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sun	1.54	1	2.3	0.1324
Sex	6.69	1	10.1	<b>0.0021</b>
Sun * Sex	2.88	1	4.3	<b>0.0406</b>
Error	0.66	79		
<b>Wing aspect ratio</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sex	11.15	1	185.9	<b>&lt;0.0001</b>
Family	0.13	6	2.1	0.0635
Error	0.06	75		

TABLE S5. Results of general linear mixed models for the effects of resources, sun, sex (fixed factors), family (random) and age (covariate) on morphological traits in *Lycaena tityrus* for individuals released in experiment 3. Models were constructed using a stepwise backwards elimination of non-significant factors. Significant p-values are given in bold.

<b>Adult body mass</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sex	14.1 * 10 <sup>-4</sup>	1	21.9	<b>&lt;0.0001</b>
Error	0.6 * 10 <sup>-4</sup>	72		
<b>Thorax mass</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sun * Sex	0.25 * 10 <sup>-4</sup>	1	3.0	0.0869
Error	0.08 * 10 <sup>-4</sup>	72		
<b>Abdomen mass</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sex	15.8 * 10 <sup>-4</sup>	1	53.3	<b>&lt;0.0001</b>
Error	0.3 * 10 <sup>-4</sup>	72		
<b>Thorax-abdomen ratio</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sex	28.32	1	147.4	<b>&lt;0.0001</b>
Age	0.56	1	2.9	0.0933
Error	0.19	71		
<b>Fat (mg)</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Resources	2.10	1	23.0	<b>&lt;0.0001</b>
Sex	1.24	1	13.6	<b>0.0004</b>
Error	0.09	71		

Table S5 continued

<b>Fat (%)</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Resources	874.6	1	29.3	<b>&lt;0.0001</b>
Sex	60.8	1	2.0	0.1582
Resources * Sun	69.2	1	2.3	0.1324
Error	29.9	70		
<b>Wing length</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Age	1.55	1	2.9	0.0938
Error	0.54	68		
<b>Wing area</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sex	255.9	1	4.0	<b>0.0492</b>
Age	1160.6	1	18.2	<b>&lt;0.0001</b>
Error	63.8	67		
<b>Wing loading</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sex	6.73	1	10.3	<b>0.0020</b>
Error	0.65	68		
<b>Wing aspect ratio</b>	<b>MS</b>	<b>d.f.</b>	<b>F</b>	<b>p</b>
Sex	8.89	1	122.4	<b>&lt;0.0001</b>
Family	0.31	5	4.2	<b>0.0023</b>
Error	0.07	63		

#### 4.3 Nutritional stress reduces flight performance and exploratory behavior in a butterfly

Published as:

Reim, E., Eichhorn, D., Roy, J.D., Steinhoff, P.O.M., Fischer, K. (2018) Nutritional stress reduces flight performance and exploratory behavior in a butterfly. *Insect Science*, DOI: 10.1111/1744-7917.12596.



## ORIGINAL ARTICLE

## Nutritional stress reduces flight performance and exploratory behavior in a butterfly

Elisabeth Reim , Danny Eichhorn, Jan D. Roy, Philip O. M. Steinhoff  and Klaus Fischer

Zoological Institute and Museum, Greifswald University, Greifswald, Germany

**Abstract** Anthropogenic global change, including agricultural intensification and climate change, poses a substantial challenge to many herbivores due to a reduced availability of feeding resources. The concomitant food stress is expected to detrimentally affect performance, amongst others in dispersal-related traits. Thus, while dispersal is of utmost importance to escape from deteriorating habitat conditions, such conditions may negatively feedback on the ability to do so. Therefore, we here investigate the impact of larval and adult food stress on traits related to dispersal ability, including morphology, physiology, flight performance, and exploratory behavior, in a butterfly. We show that inadequate nutrition during development and in the adult stage diminishes flight performance, despite some re-allocation of somatic resources. Detrimental effects of food stress on flight performance were mainly caused by reductions in body mass and storage reserves. Similar results were found for exploratory behavior. Furthermore, exploratory behavior was found to be (moderately) repeatable at the individual level, which might indicate the existence of a personality trait. This notion is further supported by the fact that flight performance and exploratory behavior were positively correlated, potentially suggesting the existence of a dispersal syndrome. In summary, our findings may have important implications for dispersal in natural environments, as the conditions requiring dispersal the most impair flight ability and thereby likely dispersal rates.

**Key words** condition; dispersal syndrome; flight ability; food stress; habitat degradation; resource allocation

## Introduction

Dispersal, often defined as any movements potentially leading to gene flow (Ronce, 2007), defines the potential of individuals to spread, for instance away from local unfavorable conditions, to avoid competition or to more suitable habitats (Matthysen, 2012). Its incidence is predicted to increase with deteriorating environmental conditions (Bowler & Benton, 2005; Legrand *et al.*, 2015). Consequently, organisms are expected to be able to base their dispersal decisions on environmental circumstances (Dall *et al.*, 2005; Clobert *et al.*, 2009). Dispersal is therefore a

crucial process to deal with heterogeneous environments (Cote & Clobert, 2007; Travis *et al.*, 2013). In the long term, it may allow individuals to track their shifting climate niche (Warren *et al.*, 2001; Hickling *et al.*, 2006) and to maintain metapopulation connectivity (Hanski & Gilpin, 1997; Baguette *et al.*, 2013).

Though dispersal is consequently an ecologically important process, the motivation to disperse and dispersal ability may differ strongly among and within populations (van Dyck & Baguette, 2005; Baguette & van Dyck, 2007; Ducatez *et al.*, 2014; Bestion *et al.*, 2015). Increasing evidence suggests that dispersing individuals are not a random subset of a given population, but that they may differ in various phenotypic traits from conspecifics, indicating the existence of so called dispersal syndromes (Bonte & Saastamoinen, 2012; Stevens *et al.*, 2013). Factors potentially affecting dispersal ability include

Correspondence: Elisabeth Reim, Zoological Institute and Museum, Greifswald University, Soldmannstraße 14, D-17489 Greifswald, Germany. Tel: +49 3834 4204297; fax: +49 3834 4204252; email: elisabeth.reim@uni-greifswald.de

morphology, physiology, and overall condition (Bowler & Benton, 2005; Bonte *et al.*, 2012; Therry *et al.*, 2014). In most insects and many other animals, dispersal is related to flight ability. Traits that have been associated with flight, and thereby at least potentially dispersal ability in flying insects, include wing size, wing aspect ratio, body mass, wing loading, thorax mass, thorax–abdomen ratio, storage reserves and flight metabolic rate (Betts & Wootton, 1988; Berwaerts *et al.*, 2002; Niitepõld *et al.*, 2009). Many of these traits show environmentally induced plasticity, which may in turn feedback on dispersal ability. Thus, if environmental conditions change, feedback loops are expected between flight-related morphology/physiology and dispersal ability.

Due to widespread effects on body size and storage reserves, food stress is certainly an important environmental factor which may detrimentally affect flight performance, as flight is a highly energy demanding process (Candy *et al.*, 1997; Suarez, 2000). In holometabolous insects, the respective energy may stem from larval storage reserves or adult income (Boggs & Freeman, 2005). Food stress in the larval phase of insect herbivores may arise from either limited availability or inadequate quality of host plants. For example, a recent study dealing with the effects of drought on plant quality revealed negative impacts of larval food stress on butterfly flight muscles (Gibbs & Weir, 2017). Larval food stress may not only diminish overall performance, but additionally trigger a reallocation of resources among competing body parts and functions such as reproduction, storage, and flight-related machinery, thus shaping the adult's physiological state (McNamara & Houston, 1996; Boggs, 2009).

Similar to larval diet, adult income also affects overall performance, as it is often of crucial importance to fuel flight, egg production, and longevity (Hill & Pierce, 1989; Fischer & Fiedler, 2001a; Bauerfeind & Fischer, 2005, 2009). Insect flight is initially fueled by carbohydrates, while longer flights rely on the lipid metabolism (Canavoso *et al.*, 2003). Consequently, inadequate nutrition during larval development and in the adult stage may exhibit negative effects on flight-related morphology and physiology. Investigating such effects seems important within the current era of anthropogenic climate change. Additionally, modern agricultural land-use practices, including fertilization, pesticide use and high mowing frequencies, will strongly reduce the availability of nectar resources for adult feeding, but also of host plants for larval feeding (Pleasants & Oberhauser, 2013; Lebeau *et al.*, 2016).

To some extent though, organisms are able to handle short-term limitations in resource availability for instance through compensatory growth (Metcalf & Monaghan,

2001). Such compensation typically involves costs that are often apparent later in life only (Metcalf & Monaghan, 2001; Stoks *et al.*, 2006; Block & Stoks, 2016). For instance, Saastamoinen *et al.* (2013) could demonstrate delayed costs of compensatory growth, as food-deprived individuals showed either reduced fecundity or lifespan. Likewise, shortage of larval-derived nutrients (i.e., nitrogenous compounds) limited reproduction in the tropical butterfly *Bicyclus anynana* (Bauerfeind & Fischer, 2005). Thus, the conditions experienced during development may affect the adult phenotype as a predictive adaptive response in order to be better prepared for dealing with detrimental conditions (e.g., Monaghan, 2008).

Against this background, we here investigate the effects of food stress on flight-related parameters in the nonpest butterfly *Lycaena tityrus*, a currently northward expanding species. As dispersal is expected to be a multicausal process (Matthysen, 2012; Legrand *et al.*, 2015), we target an array of different traits. As quantifying dispersal is challenging (Saastamoinen *et al.*, 2010), we here focus on proxies of dispersal ability such as morphology, physiology, and physiological flight performance (e.g., Zera *et al.*, 1998; Berwaerts *et al.*, 2002, 2006; Ducatez *et al.*, 2013; Kehl *et al.*, 2015). Additionally, we investigate exploratory behavior, a trait that may reflect different personalities in butterflies (Ducatez *et al.*, 2012, 2014). We set out to answer the following questions: (i) Does larval food stress negatively affect adult condition, flight performance, and exploratory behavior or are the butterflies able to compensate for a period of food deprivation during development? (ii) To what extent does adult starvation influence condition and flight performance? We predict that larval as well as adult food stress will negatively affect dispersal-related traits.

## Materials and methods

### Study organism

*Lycaena tityrus* (Poda, 1761) is a widespread temperate-zone butterfly ranging from Western Europe to central Asia (Ebert & Rennwald, 1991). This species is currently expanding its range toward higher latitudes and altitudes, which is assumed to be largely driven by anthropogenic climate change (Brunzel *et al.*, 2008; Settele *et al.*, 2008). The species is bivoltine with two discrete generations per year in most parts of its range. Overwintering takes place as half-grown larva. The principal larval host plant is *Rumex acetosa* L. (Polygonaceae), but several congeneric *Rumex* species are utilized as well (Ebert & Rennwald, 1991; Tolman & Lewington, 1998; Settele *et al.*, 2008).

Adults are nectar feeders, using a wide array of flowers (Ebert & Rennwald, 1991). For the following experiments, a total of 34 freshly eclosed, mated females from a bivoltine German population (vicinity of Greifswald, northeastern Germany; 54°02N, 13°26E) were caught in August 2015. The size of the respective habitat is ca. 20 ha, harboring a large population of *L. tityrus*. All females were transferred to a climate chamber at Greifswald University for egg laying.

### Experimental design

Caught females were kept at 26 °C, 60% relative humidity, and a L 18 : D 6 photoperiod. Unless otherwise stated, resulting offspring was reared under the same conditions. For oviposition, females were placed separately in translucent 1 L plastic pots covered with gauze and were provided with *Rumex acetosa* (oviposition substrate), fresh flowers (e.g., *Achillea millefolium* L.), water and a highly concentrated sucrose solution (20%, v/v) for adult feeding. Deposited eggs were collected daily and transferred, separated by female, to small plastic boxes. Resulting larvae were reared under the same conditions used for oviposition. Two separate experiments were performed to investigate the effects of larval and adult food stress on the flight performance of the resulting butterflies.

#### Experiment 1: Larval food stress

After hatching, the larvae of each female were randomly divided among two feeding treatments (split brood design). One group had access to food in ample supply until pupation, whereas the other group experienced a period of food deprivation (30 h) in the middle of the last larval instar. This time period was used to mimic a situation in which a larva has to leave a deteriorated host plant and search for a new one. All larvae were individually reared in small plastic boxes (125 mL). Boxes contained moistened filter paper and fresh cuttings of *R. acetosa* and *acetosella*. Boxes were checked daily and larvae were supplied with fresh food as necessary. For the starvation period, food was removed from the respective boxes. Afterward, larvae were allowed to feed again until pupation. Resulting pupae were transferred to 15 °C in order to retard further development for logistic reasons, and were kept individually in plastic cups (125 mL). Once all larvae had pupated, they were back-transferred to the warmer conditions outlined above until adult eclosion. Butterflies were kept individually and were provided with moistened cotton wool for drinking. One-day-old butterflies were first subjected to a tunnel test and

afterward to a flight performance test (see below). Thus, each individual was used for both flight tests. Flight performance tests were performed at 18, 22, and 26 ± 0.5 °C mimicking cold, medium, and warm summer conditions in Greifswald, while tunnel tests were performed at warm (26 ± 0.5 °C) conditions only because lower conditions likely prevent butterflies from flying. We included relatively low temperatures to test for flight performance under suboptimal conditions, which seems to be closely related to general flight performance in butterflies, and because too beneficial conditions may mask differences among treatments (Merckx *et al.*, 2006; Karl *et al.*, 2008).

#### Experiment 2: Adult food stress

Larvae were raised in translucent plastic boxes (125 mL) in groups of 8–10 individuals until pupation, being provided with food (*R. acetosa*) in ample supply throughout. Pupae were individually transferred to 15 °C in order to retard further development for logistic reasons. Once all larvae had pupated, pupae were back-transferred to 26 °C to induce adult eclosion. After eclosion, butterflies were randomly divided into two feeding treatments and placed individually in translucent 1 L plastic pots covered with gauze. The individuals of the control group had access to food throughout, being provided with fresh flowers, water, and a highly concentrated sucrose solution. The butterflies of the adult food stress treatment experienced a period of food deprivation for 2 d (i.e., the entire period from eclosion to testing), during which they were provided with water only. On day 2 after adult eclosion, butterflies were tested individually for flight performance at either 18 °C or 26 °C, as no significant difference was found in flight endurance between 22 and 26 °C in experiment 1 (see below). As above, butterflies were first subjected to the tunnel test and afterward to the flight performance test. Tunnel tests were also performed at 18 °C and 26 °C to test the above assumption.

#### Tunnel test

We used the tunnel test as previously described by Ducatez *et al.* (2012, 2014). The aim of this test is to investigate flight behavior/willingness to fly under novel and challenging conditions, thereby testing for exploratory behavior (individual boldness or shyness). To this end, we used a 3 m long opaque PVC-U pipe with a diameter of 80 cm in experiment 1. The pipe was placed on a table within a climate chamber set at a constant temperature of 26 ± 0.5 °C. In experiment 2, we used a slightly modified approach with a thinner pipe (diameter 40 cm) to increase

difficulty for the butterflies. In the climate chamber, all lights were turned off, except for a small daylight emitting light source (Beurer TL 40) at the end of the tunnel. Butterflies were individually released from handheld plastic cups in the center of the entrance of the pipe, that is, at a height of 40 cm (experiment 1) or 20 cm (experiment 2), to avoid any take-off effort. Butterflies were not touched directly to minimize handling effects. Thus, the values measured are mainly based on the behavioral decision to fly or not. Each individual was tested three times to assess behavioral repeatability (i.e., whether behavior is different between individuals but consistent across time; Bell *et al.*, 2009). For each trial, we recorded how far the butterfly flew into the tunnel toward the light source. Trials were terminated once an individual did not move for >5 s. Butterflies that flew through the entire tunnel were assigned the tunnel length of 3 m. The mean of the three trials was used in subsequent analyses.

#### *Flight performance test*

Butterflies were tested for flight performance using a well-established assay (cf. Ducatez *et al.*, 2012, 2013). Each individual was placed into a plastic chamber (30 cm × 16 cm × 14 cm), which was perforated at its base and fixed to a rapid agitator (IKA Vortex 4 digital). After a habituation period of 30 s, the vortex was switched on to strongly shake the chamber, preventing the butterfly from holding on to the walls. This method forces the butterflies to fly during the test, as they were otherwise lying uncomfortably on the bottom of the shaking chamber. The time an individual butterfly spent flying was recorded for 60 s, with higher values reflecting a better flight performance. After the performance test, butterflies were frozen for later analyses of potentially flight-related parameters.

#### *Morphological and physiological analyses*

To test for associations between flight behavior/performance and other traits, we measured several morphological and physiological traits. First, adult body mass was determined to the nearest 0.01 mg (Sartorius CPA225D). Then, wings, heads, and legs were removed. Thorax and abdomen were separated and afterward weighed. Abdomen fat content was measured following Fischer *et al.* (2003), but using the less poisonous acetone instead of dichloromethane. Abdomens were first dried to constant weight for 2 d at 70 °C. Abdomen dry mass was measured. Afterward, fat was extracted for two times 48 h using 1 mL of acetone for each butterfly. Solutions were exchanged between both fat extractions. Then,

abdomens were again dried for 2 d at 70 °C, after which fat-free dry mass was measured. Absolute (mg) and relative (%) abdomen fat content were determined as the mass difference between abdomen dry mass and the remaining dry mass after the two fat extractions. Forewing area and length (from basis to apex) were measured using digital images of left forewings (captured from ventrally with a digital camera mounted on a stereo microscope) and NIS elements software. Wing loading was calculated as total body mass divided by forewing area, and wing aspect ratio as  $4 \times \text{forewing length}^2$  divided by the forewing area (Berwaerts *et al.*, 2002). Thorax–abdomen ratio was calculated by dividing thorax through abdomen mass.

#### *Statistical analyses*

Data on developmental, morphological and physiological traits were analyzed with general linear mixed models (GLMMs; experiment 1) and general linear models (GLMs; experiment 2), using feeding treatment and sex as well as the respective interaction as fixed factors. In experiment 1, family (i.e., the offspring of each individual female) was used additionally as a random effect. Only families with  $n > 7$  offspring were considered to cover within-family variation. In experiment 2 consideration of family was not possible as according data were not available. In experiment 1, data on abdomen mass and in experiment 2 data on abdomen mass, thorax–abdomen ratio, and absolute fat content were LN transformed prior to analysis to meet model requirements. Data on exploratory behavior and flight performance were also analyzed with GLMs, using the same factors mentioned above and additionally test temperature for flight endurance in the vortex and the tunnel test in experiment 2. For experiment 1, data on the flight distance in the tunnel were LN transformed and on flight endurance  $x^3$  transformed prior to analyses. Interaction terms between feeding treatment, sex, and test temperature (if used) were also included.

To additionally test for the impact of continuous variables (morphology, physiology) on exploratory behavior and flight performance, we first performed principal component analyses (PCAs) owing to strong intercorrelations among traits. PCAs were calculated separately for males and females, as sexes differed strongly in a variety of traits (see below). In experiment 1, we used the first five principal components (PCs) for further analyses, having Eigenvalues between 3.1 and 0.9 for males and between 3.7 and 1.0 for females (cf. Table 1). In experiment 2, we used the first four PCs for further analyses, having Eigenvalues between 3.9 and 1.0 for males and between 5.7 and 0.8 for females (cf. Table 2). As PCAs had to be

**Table 1** Results of a principal component analyses for larval starved males (A) and females (B) including Eigen values, percentage of the variance explained, cumulative percentage of the variance explained, and *r* values of correlations with continuous variables (rows 4–13) for principal components (PC) 1–5 (experiment 1). *r* values > 0.7 in bold.

No.	Parameter	PC1	PC2	PC3	PC4	PC5
<b>(A) Males</b>						
1	Eigen value	3.1	2.4	1.9	1.7	0.9
2	Variance explained (%)	30.8	23.8	19.2	16.6	8.5
3	Cumulative variance (%)	30.8	54.6	73.8	90.4	98.9
4	Adult body mass (mg)	<b>0.956</b>	0.261	−0.061	0.038	0.017
5	Thorax mass (mg)	<b>0.862</b>	0.248	−0.111	−0.379	0.093
6	Abdomen mass (mg)	<b>0.723</b>	0.121	0.102	0.650	−0.139
7	Thorax–abdomen ratio	0.095	0.157	−0.187	<b>−0.930</b>	0.234
8	Fat (mg)	0.297	−0.318	<b>0.880</b>	−0.098	0.112
9	Fat (%)	0.070	−0.352	<b>0.903</b>	−0.147	0.141
10	Wing length (mm)	0.009	<b>0.784</b>	0.346	−0.222	−0.464
11	Wing area (mm <sup>2</sup> )	−0.048	<b>0.951</b>	0.230	0.029	0.003
12	Wing loading (mg/cm <sup>2</sup> )	<b>0.884</b>	−0.386	−0.241	−0.004	0.032
13	Wing aspect ratio	0.118	−0.568	−0.015	−0.376	<b>−0.722</b>
<b>(B) Females</b>						
1	Eigen value	3.7	2.3	1.7	1.3	1.0
2	Variance explained (%)	37.0	22.6	17.0	13.0	9.6
3	Cumulative variance (%)	37.0	59.7	76.6	89.5	99.1
4	Adult body mass (mg)	<b>0.963</b>	0.174	−0.188	−0.016	0.042
5	Thorax mass (mg)	<b>0.850</b>	0.234	−0.361	0.194	0.216
6	Abdomen mass (mg)	<b>0.925</b>	0.078	0.078	−0.293	−0.197
7	Thorax–abdomen ratio	−0.265	0.173	−0.503	0.630	0.497
8	Fat (mg)	0.618	−0.339	0.620	0.275	0.113
9	Fat (%)	0.206	−0.449	<b>0.702</b>	0.458	0.174
10	Wing length (mm)	0.139	<b>0.855</b>	0.206	0.343	−0.299
11	Wing area (mm <sup>2</sup> )	0.133	<b>0.928</b>	0.336	0.024	0.074
12	Wing loading (mg/cm <sup>2</sup> )	<b>0.817</b>	−0.410	−0.396	−0.023	0.008
13	Wing aspect ratio	−0.013	−0.256	−0.285	0.606	−0.697

constructed separately for males and females, PCs could not be included in the above models testing for sexual differences in flight performance. We therefore constructed another set of models separately for each sex, by including the respective PCs and the results of tunnel or vortex test as covariates, followed by a stepwise backward elimination of nonsignificant factors. All statistical tests were performed with Statistica 12.0 (Tulsa, StatSoft, OK, USA). All means are given  $\pm 1$  SE and are based on untransformed data for easier reference.

## Results

### *Experiment 1: Larval food stress*

Developmental and morphological traits were strongly affected by feeding treatment and sex (Table 3). Feed-

ing treatment significantly affected larval development time, adult body mass, abdomen mass, thorax mass, wing length, wing area, and wing loading, but not pupal development time, thorax–abdomen ratio, abdomen fat content, and wing aspect ratio. Food stress resulted in longer larval development times, lower body masses, smaller wings, and a lower wing loading (for mean values see Table S1). Feeding treatment by sex interactions were significant for larval and pupal development time only, showing that females responded more strongly to food stress than males. Significant sex differences were found for all traits except from thorax mass, wing length, and wing area, indicating that females had on average longer development times, higher abdomen and concomitantly total body masses, a lower thorax–abdomen ratio, higher absolute but lower relative abdomen fat contents, a higher wing loading, and a lower wing aspect ratio than males. Additionally,

**Table 2** Results of a principal component analyses for adult starved males (A) and females (B) including Eigen values, percentage of the variance explained, cumulative percentage of the variance explained, and *r* values of correlations with continuous variables (rows 4–13) for principal components (PC) 1–4 (experiment 2). *r* values > 0.7 in bold.

No.	Parameter	PC1	PC2	PC3	PC4
<b>(A) Males</b>					
1	Eigen value	3.9	2.9	1.3	1.0
2	Variance explained (%)	39.4	29.0	13.2	10.1
3	Cumulative variance (%)	39.4	68.4	81.6	91.7
4	Adult body mass (mg)	<b>0.953</b>	−0.084	0.256	−0.060
5	Thorax mass (mg)	0.588	−0.670	0.002	−0.287
6	Abdomen mass (mg)	<b>0.820</b>	0.420	0.297	−0.001
7	Thorax–abdomen ratio	−0.299	<b>−0.771</b>	−0.308	−0.284
8	Fat (mg)	0.655	0.323	−0.636	0.105
9	Fat (%)	0.580	0.187	<b>−0.737</b>	0.199
10	Wing length (mm)	0.425	<b>−0.831</b>	0.053	0.234
11	Wing area (mm <sup>2</sup> )	0.509	<b>−0.759</b>	−0.006	−0.076
12	Wing loading (mg/cm <sup>2</sup> )	<b>0.840</b>	0.336	0.309	−0.027
13	Wing aspect ratio	−0.094	0.397	0.159	<b>0.857</b>
<b>(B) Females</b>					
1	Eigen value	5.7	2.0	1.0	0.8
2	Variance explained (%)	57.1	20.1	10.0	7.7
3	Cumulative variance (%)	57.1	77.3	87.3	95.0
4	Adult body mass (mg)	<b>0.981</b>	−0.050	−0.040	−0.105
5	Thorax mass (mg)	<b>0.804</b>	−0.404	−0.119	0.242
6	Abdomen mass (mg)	<b>0.964</b>	0.088	−0.090	−0.202
7	Thorax–abdomen ratio	<b>−0.785</b>	−0.387	0.040	0.427
8	Fat (mg)	<b>0.770</b>	0.413	0.279	0.228
9	Fat (%)	0.432	0.586	0.468	0.453
10	Wing length (mm)	0.584	<b>−0.763</b>	0.204	0.069
11	Wing area (mm <sup>2</sup> )	<b>0.725</b>	−0.606	−0.123	0.214
12	Wing loading (mg/cm <sup>2</sup> )	<b>0.920</b>	0.205	0.014	−0.237
13	Wing aspect ratio	−0.262	−0.427	<b>0.790</b>	−0.343

females tended to have larger wings than males. Based on the above data we performed two PCAs, one for each sex. In both sexes the resulting first five PCs were most strongly related to body mass (PC1), wing size (PC2), abdomen fat content (PC3), thorax–abdomen ratio (PC4), and wing aspect ratio (PC5; Table 1).

Flight performance traits were first analyzed with GLMMs including the factors feeding treatment, sex, and family. In the tunnel test, exploratory behavior was significantly affected by the factors feeding treatment, sex, and family (Table 4, part A). Control individuals and males covered longer distances than food-stressed individuals and females, respectively (Fig. 1A). In a next step, we calculated GLMMs separately for males and females to include PCs as covariates. For males, including covariates revealed significant effects

of feeding treatment, PC3 (reflecting fat content), and the results of the vortex test, while family was not included anymore. Flight distance was positively related to PC3 (beta  $0.17 \pm 0.09$ ), suggesting that flight distance tended to be positively related to abdomen fat content in males. Furthermore, flight distance was positively correlated with flight performance (beta  $0.17 \pm 0.08$ ). Hence, males that covered longer distances in the tunnel also showed a better flight performance in the vortex. For females, in contrast, no parameters were selected. Overall, the repeatability of the tunnel test was moderate but significant, with correlations among the three replicated measures being positive with *r* values ranging between 0.464 and 0.568 (all  $P < 0.0001$ ,  $n = 301$ ).

In the vortex test, flight performance was significantly affected by test temperature and feeding treatment. Flight

**Table 3** General linear mixed model results for the effects of larval feeding treatment, sex (both fixed), and family (random factor) on developmental and morphological traits in *Lycaena tityrus* (experiment 1). When not attaining significance, interactions and the random effect “family” were removed in a stepwise manner. Significant *P* values are given in bold.


Traits	MS	DF	DF error	<i>F</i>	<i>P</i>
<b>Larval time (d)</b>					
Feeding treatment	43.0	1	277	17.5	<b>&lt;0.0001</b>
Sex	438.5	1	277	178.3	<b>&lt;0.0001</b>
Feeding × sex	16.7	1	277	6.8	<b>0.0097</b>
<b>Pupal time (d)</b>					
Feeding treatment	15.7	1	277	2.5	0.1166
Sex	184.0	1	277	29.1	<b>&lt;0.0001</b>
Feeding × sex	122.0	1	277	19.3	<b>&lt;0.0001</b>
<b>Adult body mass (mg)</b>					
Feeding treatment	1711.4	1	279	78.9	<b>&lt;0.0001</b>
Sex	1659.2	1	279	76.5	<b>&lt;0.0001</b>
<b>Abdomen mass (mg)</b>					
Feeding treatment	1.8	1	279	34.0	<b>&lt;0.0001</b>
Sex	14.4	1	279	279.8	<b>&lt;0.0001</b>
<b>Thorax mass (mg)</b>					
Feeding treatment	514.5	1	279	77.1	<b>&lt;0.0001</b>
Sex	15.7	1	279	2.4	0.1257
<b>Thorax–abdomen ratio</b>					
Feeding treatment	0.5	1	279	3.5	0.0613
Sex	37.9	1	279	283.9	<b>&lt;0.0001</b>
<b>Fat (absolute)</b>					
Feeding treatment	0.01	1	279	1.0	0.3218
Sex	0.53	1	279	59.3	<b>&lt;0.0001</b>
<b>Fat (relative)</b>					
Feeding treatment	39.2	1	279	3.1	0.0800
Sex	342.8	1	279	27.0	<b>&lt;0.0001</b>
<b>Wing length (mm)</b>					
Feeding treatment	35.8	1	279	77.4	<b>&lt;0.0001</b>
Sex	0.4	1	279	0.9	0.3432
<b>Wing area (mm<sup>2</sup>)</b>					
Feeding treatment	5363	1	279	69.8	<b>&lt;0.0001</b>
Sex	280	1	279	3.7	0.0571
<b>Wing loading (mg/cm<sup>2</sup>)</b>					
Feeding treatment	279	1	279	5.2	<b>0.0228</b>
Sex	1795	1	279	33.8	<b>&lt;0.0001</b>
<b>Wing aspect ratio</b>					
Feeding treatment	0.4	1	279	1.1	0.3057
Sex	1.8	1	279	4.6	<b>0.0334</b>

endurance increased with increasing test temperature and was higher in control than in starved individuals (Table 4, part B, Fig. 1B). Additionally, females tended to show a shorter flight endurance than males. Including covariates in the subsequent analyses separated by sex revealed significant effects of temperature and PC1 (reflecting body mass) in both sexes, and the performance in the vortex test in males, while feeding treatment was not included anymore. The latter indicates that the effects of

**Table 4** General linear model results for (A) the effects of larval feeding treatment, sex, and family on the flight distance covered in a tunnel and (B) for the effects of larval feeding treatment, test temperature, sex, and family on flight endurance in a vortex in *Lycaena tityrus* (experiment 1). Second models were constructed by additionally including PCs 1–5 and the flight performance in the tunnel or in the vortex as covariates, followed by a stepwise backward elimination of nonsignificant factors. The latter models were constructed separately for males and females, owing to strong variation in morphological traits among the sexes (cf. Table S1). Throughout, family was included as random effect, all others as fixed effects. For (A) no factor was selected in model 2 for females. Significant *P* values are given in bold.

Models	MS	DF	DF error	<i>F</i>	<i>P</i>
<b>(A)</b>					
Tunnel test model 1					
Feeding treatment	5.0	1	277.5	8.5	<b>0.0039</b>
Sex	3.7	1	277.4	6.4	<b>0.0121</b>
Family	1.1	22	256.0	2.0	<b>0.0061</b>
Males model 2					
Feeding treatment	2.9	1	141	4.5	<b>0.0351</b>
PC3 (fat content)	2.6	1	141	4.1	<b>0.0439</b>
Vortex test	3.0	1	141	4.7	<b>0.0320</b>
Females model 2					
No factors selected					
<b>(B)</b>					
Vortex test model 1					
Feeding treatment	$1.5 \times 10^{10}$	1	275	4.1	<b>0.0429</b>
Temperature	$8.2 \times 10^{10}$	1	275	22.5	<b>&lt;0.0001</b>
Sex	$1.2 \times 10^{10}$	1	275	3.3	0.0708
Males model 2					
Temperature	$5.7 \times 10^{10}$	2	140	16.7	<b>&lt;0.0001</b>
PC1 (body mass)	$1.7 \times 10^{10}$	1	140	5.1	<b>0.0256</b>
Tunnel test	$1.7 \times 10^{10}$	1	140	5.0	<b>0.0272</b>
Females model 2					
Temperature	$4.0 \times 10^{10}$	2	130	11.7	<b>&lt;0.0001</b>
PC1 (body mass)	$1.6 \times 10^{10}$	1	130	4.6	<b>0.0339</b>
Tunnel test	$1.2 \times 10^{10}$	1	130	3.5	0.0646

feeding treatment are mediated by variation in PC1, that is, body mass. Flight endurance was positively related to PC1 (males: beta  $0.17 \pm 0.08$ ; females: beta  $0.20 \pm 0.08$ ). Thus, high body mass enhanced flight endurance. Flight performance was positively correlated with the tunnel test results in males (beta  $0.17 \pm 0.08$ ). Hence, males that covered longer distances in the tunnel also showed a better flight performance. In females, flight distance in the tunnel test tended to be positively correlated to flight performance (beta  $0.15 \pm 0.09$ ). In general, results of the tunnel test and vortex test were positively related (Spearman’s correlation:  $r = 0.19$ ,  $P < 0.05$ ; Fig. S1A).



**Fig. 1** Flight distance covered in a tunnel (A) and flight endurance in a vortex (B, means + 1 SE) in *Lycaena tityrus* males and females in relation to larval feeding treatment and test temperature (in B, experiment 1). Group sample sizes range between 74 and 77 (in A) and 24 and 26 (in B). Males: filled bars; females: open bars.

### Experiment 2: Adult food stress

Regarding morphological and physiological traits, adult feeding treatment significantly affected adult body mass, abdomen mass, thorax mass, thorax–abdomen ratio, absolute and relative fat content, wing loading, and wing aspect ratio, but not wing length and wing area (Table 5). Absence of adult food sources resulted in lower abdomen, thorax, and body masses, a higher thorax–abdomen ratio and reduced wing loading, lower absolute and relative fat contents, and a higher wing aspect ratio (for means see Table S2). Significant sex differences were found for all traits measured, except for wing length. Body and abdomen mass, absolute fat content, wing area, and wing loading were higher while thorax mass, thorax–abdomen ratio and wing aspect ratio were lower in females than in males. Females tended to have smaller wings than males. Sexual differences in adult body mass and concomitantly wing loading though were restricted to control individuals (significant feeding treatment by sex interactions). Again, PCAs were performed for both sexes separately. In males, the extracted PCs represent in the first place body mass (PC1), wing size and thorax–abdomen ratio (PC2), fat content (PC3), and wing aspect ratio (PC4). In females, the PCs reflect mainly body mass (PC1), wing size (PC2), wing aspect ratio (PC3), and fat content and thorax–abdomen ratio (PC4; cf. Table 2).

In the GLM including the factor sex, the distance covered in the tunnel was significantly affected by the factor temperature only (Table 6, part A), with butterflies covering longer distances at the higher temperature (26 °C:  $30.7 \pm 3.3$  cm, 18 °C:  $2.3 \pm 6.3$  cm; Fig. 2A). Including subsequently principal components and the flight performance in the vortex as covariates did not reveal additional significant effects. Overall, the repeatability of the tunnel

test was weak but significant in two out of three cases, with  $r$  values ranging between 0.142 and 0.295 (trial 1 and 2:  $P = 0.0045$ ; trial 1 and 3:  $P = 0.0001$ ; trial 2 and 3:  $P = 0.0685$ ;  $n = 165$ ).

In the vortex test, flight performance was significantly affected by feeding treatment and temperature, with flight endurance being higher at the higher temperature and in control than in starved individuals (Table 6, part B; Fig. 2B). Including principal components and the results of the tunnel test as covariates revealed significant effects of feeding treatment, PC1 (reflecting body mass), and the tunnel test results for males, and of temperature for females. Additionally, PC4 (reflecting relative fat content and thorax–abdomen ratio) tended to affect flight performance in females. In males, flight performance was positively correlated with PC1 (beta  $0.28 \pm 0.12$ ) and the tunnel test results (beta  $0.32 \pm 0.11$ ). Thus, high body mass increased flight performance and males that covered longer distances in the tunnel also had a better flight performance. In females, flight performance tended to be positively correlated with PC4 (beta  $0.17 \pm 0.09$ ). Thus, a high relative fat content and thorax–abdomen ratio tended to increase flight performance. In general, results of the tunnel test and vortex test were positively related (Spearman's correlation:  $r = 0.22$ ,  $P < 0.05$ ; Fig. S1B).

## Discussion

### *Effects of feeding treatment on morphological and physiological traits*

Food stress during larval development resulted, as expected, in longer development time, lower body mass and

**Table 5** General linear model results for the effects of adult feeding treatment and sex (both fixed) on morphological and physiological traits in *Lycaena tityrus* (experiment 2). Non-significant interactions were removed from models in a stepwise manner. Significant *P* values are given in bold.


Traits	MS	DF	<i>F</i>	<i>P</i>
<b>Adult body mass (mg)</b>				
Feeding treatment	$12.3 \times 10^{-4}$	1	26.0	<b>&lt;0.0001</b>
Sex	$5.6 \times 10^{-4}$	1	11.8	<b>0.0008</b>
Feeding $\times$ sex	$3.6 \times 10^{-4}$	1	7.7	<b>0.0061</b>
Error	$0.5 \times 10^{-4}$	162		
<b>Abdomen mass (mg)</b>				
Feeding treatment	5.3	1	53.9	<b>&lt;0.0001</b>
Sex	8.7	1	88.1	<b>&lt;0.0001</b>
Error	0.1	163		
<b>Thorax mass (mg)</b>				
Feeding treatment	$0.33 \times 10^{-4}$	1	4.0	<b>0.0485</b>
Sex	$0.68 \times 10^{-4}$	1	8.2	<b>0.0048</b>
Error	$0.08 \times 10^{-4}$	163		
<b>Thorax–abdomen ratio</b>				
Feeding treatment	3.3	1	31.8	<b>&lt;0.0001</b>
Sex	10.8	1	105.7	<b>&lt;0.0001</b>
Error	0.1	163		
<b>Fat (absolute)</b>				
Feeding treatment	30.4	1	44.4	<b>&lt;0.0001</b>
Sex	13.2	1	19.3	<b>&lt;0.0001</b>
Error	0.7	163		
<b>Fat (relative)</b>				
Feeding treatment	475.8	1	18.8	<b>&lt;0.0001</b>
Sex	116.1	1	4.6	<b>0.0339</b>
Error	25.4	163		
<b>Wing length (mm)</b>				
Feeding treatment	0.3	1	0.6	0.4371
Sex	1.8	1	3.6	<b>0.0582</b>
Error	0.5	161		
<b>Wing area (mm<sup>2</sup>)</b>				
Feeding treatment	25	1	0.4	0.5084
Sex	464	1	8.1	<b>0.0050</b>
Error	57	161		
<b>Wing loading (mg/cm<sup>2</sup>)</b>				
Feeding treatment	1429	1	41.8	<b>&lt;0.0001</b>
Sex	295	1	8.6	<b>0.0038</b>
Feeding $\times$ sex	228	1	6.7	<b>0.0107</b>
Error	34	160		
<b>Aspect ratio</b>				
Feeding treatment	1.6	1	12.6	<b>0.0005</b>
Sex	18.8	1	148.2	<b>&lt;0.0001</b>
Error	0.1	161		

**Table 6** General linear model results for the effects of adult feeding treatment, temperature, and sex (all fixed) on the flight distance covered in a tunnel (A) and on flight endurance in a vortex (B) in *Lycaena tityrus* (experiment 2). Second models were constructed by additionally including PCs 1–4 and the flight performance in tunnel or vortex as covariates, followed by a stepwise backward elimination of nonsignificant factors. The latter models were constructed separately for males and females, owing to strong variation in morphological traits among the sexes (cf. Table S2). For (A), no PCs were included in either males or females. Significant *P* values are given in bold.

Models	MS	DF	<i>F</i>	<i>P</i>
<b>(A)</b>				
Tunnel test model 1				
Temperature	28699.8	1	23.96	<b>&lt;0.0001</b>
Error	1198.0	164		
Males model 2				
No factors selected				
Females model 2				
No factors selected				
<b>(B)</b>				
Vortex test model 1				
Feeding treatment	2192.1	1	7.33	<b>0.0075</b>
Temperature	9711.6	1	32.46	<b>&lt;0.0001</b>
Error	299.2	163		
Males model 2				
Feeding treatment	1228.4	1	4.2	<b>0.0445</b>
PC1 (body mass)	1709.9	1	5.9	<b>0.0185</b>
Tunnel test	2394.0	1	8.2	<b>0.0058</b>
Error	291.3	58		
Females model 2				
Temperature	7766.3	1	27.1	<b>&lt;0.0001</b>
PC4 (aspect ratio)	1079.1	1	3.8	0.0551
Error	286.5	99		

wing loading, and an overall smaller adult size. Thus, despite a prolonged larval period, individuals remained smaller (accompanied by lower wing loadings) compared with individuals that developed under control conditions (Blanckenhorn, 1999; Fischer & Fiedler, 2001b; Boggs & Freeman, 2005; Bauerfeind & Fischer, 2009). This could suggest that development time and body size are both critically important to fitness. Otherwise larvae should further increase larval time to avoid size reductions (in case of a selective premium on body size) or not extend the larval time at all (in case of a selective premium on larval time; Bauerfeind & Fischer, 2005).

Interestingly, only the females showed longer larval and pupal development times after larval food stress,



**Fig. 2** Flight distance covered in a tunnel (A) and flight endurance in a vortex (B, means + 1 SE) in *Lycaena tityrus* in relation to adult feeding treatment and test temperature (experiment 2). Group sample sizes range between 26 and 59 individuals. Food: filled bars; no food: open bars.

suggesting that the above general patterns differ among the sexes. Thus, males seem to have a selective premium on fast development to ensure early emergence that is assumed to increase mating success (protandry selection; Wiklund & Fagerström, 1977; Fischer & Fiedler, 2000; Karl & Fischer, 2008), which likely explains the lack of response in development time found here (Karl *et al.*, 2010). Females, in contrast, did prolong development time, suggesting a premium on body size, which is often associated with increased fecundity (Bauerfeind & Fischer, 2005). In general, the observed sex differences with females being larger than males reflect well-known and expected patterns found in many insects, which most likely result from differential selective pressures (Gilchrist, 1990; van Dyck & Wiklund, 2002; Merckx & van Dyck, 2005; Berwaerts *et al.*, 2006). It is striking that abdomen fat content remained unaffected by larval food stress, which may aid dispersal and/or investment into reproduction as

fat is the principal energy store in insects (e.g., Karl & Fischer, 2008).

Effects of adult food stress on body mass were stronger in females than in males, which is likely attributable to a lack of egg production under food deprivation. Note that Copper butterflies are strongly relying on adult feeding for egg production (Fischer & Fiedler, 2001a). Although attaining significance, the effect of food stress on thorax mass was comparatively weak, suggesting that butterflies may have tried to preserve thoracic muscles and thus flight performance (high thorax–abdomen ratio; Marden, 1989; Thomas *et al.*, 1998; Berwaerts *et al.*, 2002) to enhance the ability to disperse to a more suitable habitat (Clobert *et al.*, 2009). However, note that in insects storage reserves are mainly conserved within the abdomen, such that effects of food shortage are less likely to affect thorax as compared with abdomen mass. We have no explanation for the effect of adult food stress on wing aspect ratio, which is determined prior to adult eclosion. Taken together, the above results indicate some changes in somatic allocation between control and food-stressed individuals in order to maintain dispersal ability. Furthermore, they clearly show that our feeding treatments were successful in manipulating morphology and physiology, a necessary prerequisite to examine such variation on flight performance.

#### *Effects on exploratory behavior and flight performance*

Flight behavior was measured here using two established laboratory assays, a tunnel test and flight endurance. Interestingly, the results of both tests were statistically (though weakly) related, such that individuals with a higher flight endurance also performed better in the tunnel test. These findings suggest that either the tunnel test measures flight ability along with exploratory behavior, or that individuals in better condition (as evidenced by a better performance in the vortex) are more likely to exhibit exploratory behavior. Anyway, it is striking that in our study both approaches yielded relatively similar results.

We were able to demonstrate that starvation during development as well as in the adult stage may have detrimental consequences for flight performance and thus potentially dispersal ability. Flight endurance in the vortex was in both experiments reduced in food-stressed individuals. Larval starvation additionally negatively affected the distance covered in the tunnel. A lack of adequate nutrition has often negative impacts on adult physiology and overall performance of organisms (Boggs, 2009; Hamel *et al.*, 2009), as also indicated in our study. Interestingly, a study of Saastamoinen *et al.* (2010) revealed an opposite

pattern, namely that food-stressed individuals performed better during a flight test than control individuals, perhaps due to an altered resource allocation. In our study though flight performance was clearly reduced through food stress, which might be problematic in the current era of anthropogenic global change. While increasing habitat change and fragmentation warrant dispersal, rising temperatures and habitat degradation diminish habitat quality, which may negatively feedback on dispersal ability through increased food stress. Note that such effects may be exaggerated by (interactions with) additional stressors associated with climate change, including desiccation and heat stress. For instance, exposure of fruit flies to food and heat stress decreased flight distance and duration in comparison to flies experiencing no-stress treatments (Wang *et al.*, 2009). Likewise, an exposure to high temperatures during development decreased flight metabolic rates and hence likely flight performance in the Glanville fritillary butterfly (Mattila & Hanski, 2014).

Measuring morphological and physiological traits and including them as covariates in statistical models may shed further light on the mechanistic underpinning of such reduced performance under food stress. After including covariates, feeding treatment only remained marginally significant for larval food-stressed males in the tunnel test and adult food-stressed males in the vortex test. These findings suggest that the poorer performance of food-stressed individuals is mechanistically in the first place caused by stress-induced changes in morphology and physiology. Specifically, body mass and relative fat content seemed to be most relevant here. Body mass was positively related to flight endurance (vortex test) in males in both experiments, and in females having experienced larval food stress. Moreover, flight endurance after adult food stress tended to be positively related to relative fat content in females. These findings suggest that individuals that managed to maintain high body mass and thereby amounts of storage reserves in periods of food stress showed a better flight performance. Storage reserves such as lipids are commonly used to fuel flight activity, reflect a generally good condition, and are related to acceleration capacity (Zera *et al.*, 1998; Berwaerts *et al.*, 2002; Karl & Fischer, 2008). Similar results were obtained in the tunnel test with males that had experienced larval starvation, in which the distance covered in the tunnel was positively related to relative fat content.

Sex differences in flight performance were obvious in the larval food stress experiment only. Here, females showed a tendency to perform worse in comparison with males in both flight tests. This might be due to their higher mass and thus wing loading, which leads to higher energetic costs during flight (Wickman, 2009).

This does not necessarily contradict our above conclusions. While those individuals dealing with food stress best may have a generally better condition, female flight ability may nevertheless suffer from their generally higher wing loading.

Ambient temperature, the most important environmental factor for ectotherms (e.g., Deutsch *et al.*, 2008), is mainly determining body temperature (Wickman, 2009). Body temperature in turn influences the flight performance of ectotherms because physiology is temperature dependent (Huey & Kingsolver, 1993; Gilchrist, 1996; Forsman, 1999). Our results thus confirm other findings, which revealed positive relationships between high temperatures and flight performance (Forsman, 1999).

## Conclusions

In our study, we were able to demonstrate that inadequate nutrition negatively affected flight performance. This is true despite some re-allocation of somatic resources in favor of dispersal-related traits. Furthermore, higher temperatures increased flight performance. This suggests that flight performance, and presumably also dispersal, are strongly dependent on environmental conditions such as ambient temperatures. Detrimental effects of starvation on flight performance seem to be conveyed in the first place by reductions in body mass and storage reserves, and thus the total amount of resources being available to the given individual. Our findings may have important implications for dispersal in natural environments. While deteriorating habitat conditions are expected to promote dispersal as an adaptive response (Legrand *et al.*, 2015), they may at the same time interfere with an individual's ability to do so.

The tunnel test used here was previously developed to detect possible behavioral differences between individuals under unfavorable conditions (Ducatez *et al.*, 2012, 2014; Ducatez & Baguette, 2016), while the vortex test addresses flight ability. Here, the latter produced more clear-cut results. Nevertheless, the tunnel test was shown to produce (moderately) repeatable results at the individual level, which might indicate the existence of personality traits in butterflies. Interestingly, both tests yielded similar results and were statistically correlated, which might point toward the existence of a dispersal syndrome (e.g., for *Pieris brassicae* in Ducatez *et al.*, 2012). Thus, the individuals performing repeatedly better than others in different types of flight tests might be the ones more apt to disperse under natural conditions (Delattre *et al.*, 2013).

## Acknowledgments

We thank three anonymous reviewers for helpful comments. This research was funded by the DFG research training group RESPONSE (DFG GRK 2010).

## Disclosure

The authors declare no conflict of interest.

## References

- Baguette, M., Blanchet, S., Legrand, D., Stevens, V.M. and Turlure, C. (2013) Individual dispersal, landscape connectivity and ecological networks. *Biological Reviews*, 88, 310–326.
- Baguette, M. and van Dyck, H. (2007) Landscape connectivity and animal behavior: functional grain as a key determinant for dispersal. *Landscape Ecology*, 22, 1117–1129.
- Bauerfeind, S.S. and Fischer, K. (2005) Effects of food stress and density in different life stages on reproduction in a butterfly. *Oikos*, 111, 514–524.
- Bauerfeind, S.S. and Fischer, K. (2009) Effects of larval starvation and adult diet-derived amino acids on reproduction in a fruit-feeding butterfly. *Entomologia Experimentalis et Applicata*, 130, 229–237.
- Bell, A.M., Hankison, S.J. and Laskowski, K.L. (2009) The repeatability of behaviour: a meta-analysis. *Animal Behaviour*, 77, 771–783.
- Berwaerts, K., Aerts, P. and van Dyck, H. (2006) On the sex-specific mechanisms of butterfly flight: flight performance relative to flight morphology, wing kinematics, and sex in *Pararge aegeria*. *Biological Journal of the Linnean Society*, 89, 675–687.
- Berwaerts, K., van Dyck, H. and Aerts, P. (2002) Does flight morphology relate to flight performance? An experimental test with the butterfly *Pararge aegeria*. *Functional Ecology*, 16, 484–491.
- Bestion, E., Clobert, J. and Cote, J. (2015) Dispersal response to climate change: scaling down to intraspecific variation. *Ecology Letters*, 18, 1226–1233.
- Betts, C.R. and Wootton, R.J. (1988) Wing shape and flight behaviour in butterflies (Lepidoptera: Papilionoidea and Hesperioidea): a preliminary analysis. *Journal of Experimental Biology*, 138, 271–288.
- Blanckenhorn, W.U. (1999) Different growth responses to temperature and resource limitation in three fly species with similar life histories. *Evolutionary Ecology*, 13, 395–409.
- Block, M.D.E. and Stoks, R. (2016) Fitness effects from egg to reproduction: bridging the life history transition. *Ecology*, 86, 185–197.
- Boggs, C.L. (2009) Understanding insect life histories and senescence through a resource allocation lens. *Functional Ecology*, 23, 27–37.
- Boggs, C.L. and Freeman, K.D. (2005) Larval food limitation in butterflies: effects on adult resource allocation and fitness. *Oecologia*, 144, 353–361.
- Bonte, D., van Dyck, H., Bullock, J.M., Coulon, A., Delgado, M., Gibbs, M. et al. (2012) Costs of dispersal. *Biological Reviews*, 87, 290–312.
- Bonte, D. and Saastamoinen, M. (2012) Dispersal syndromes in butterflies and spiders. *Dispersal Ecology and Evolution* (eds J. Clobert, M. Baguette, T.G. Benton & J.M. Bullock), pp. 161–170. Oxford University Press, Oxford.
- Bowler, D.E. and Benton, T.G. (2005) Causes and consequences of animal dispersal strategies: relating individual behaviour to spatial dynamics. *Biological Reviews*, 80, 205–225.
- Brunzel, S., Bussmann, M. and Obergruber, H. (2008) Deutliche Veränderungen von Tagfalterzönosen als Folge von Ausbreitungsprozessen. Erste Ergebnisse eines Monitorings über 17 Jahre. *Natur und Landschaft*, 6, 280–287.
- Canavoso, L.E., Stariolo, R. and Rubiolo, E.R. (2003) Flight metabolism in *Panstrongylus megistus* (Hemiptera: Reduviidae): the role of carbohydrates and lipids. *Memorias do Instituto Oswaldo Cruz*, 98, 909–914.
- Candy, D.J., Becker, A. and Wegener, G. (1997) Coordination and integration of metabolism in insect flight. *Comparative Biochemistry and Physiology—B Biochemistry and Molecular Biology*, 117, 497–512.
- Clobert, J., Le Galliard, J.F., Cote, J., Meylan, S. and Massot, M. (2009) Informed dispersal, heterogeneity in animal dispersal syndromes and the dynamics of spatially structured populations. *Ecology Letters*, 12, 197–209.
- Cote, J. and Clobert, J. (2007) Social information and emigration: lessons from immigrants. *Ecology Letters*, 10, 411–417.
- Dall, S.R.X., Giraldeau, L.A., Olsson, O., McNamara, J.M. and Stephens, D.W. (2005) Information and its use by animals in evolutionary ecology. *Trends in Ecology and Evolution*, 20, 187–193.
- Delattre, T., Baguette, M., Burel, F., Stevens, V.M., Quénot, H. and Vernon, P. (2013) Interactive effects of landscape and weather on dispersal. *Oikos*, 122, 1576–1585.
- Deutsch, C.A., Tewksbury, J.J., Huey, R.B., Sheldon, K.S., Ghalambor, C.K., Haak, D.C. et al. (2008) Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of the National Academy of Sciences USA*, 105, 6668–6672.
- Ducatez, S. and Baguette, M. (2016) Inter-individual variation in shivering behaviour in the migratory painted lady *Vanessa cardui*. *Ecological Entomology*, 41, 131–137.
- Ducatez, S., Baguette, M., Trochet, A., Chaput-Bardy, A., Legrand, D., Stevens, V. et al. (2013) Flight endurance and

- heating rate vary with both latitude and habitat connectivity in a butterfly species. *Oikos*, 122, 601–611.
- Ducatez, S., Humeau, A., Congretel, M., Fréville, H. and Baguette, M. (2014) Butterfly species differing in mobility show different structures of dispersal-related syndromes in the same fragmented landscape. *Ecography*, 37, 378–389.
- Ducatez, S., Legrand, D., Chaput-Bardy, A., Stevens, V.M., Fréville, H. and Baguette, M. (2012) Inter-individual variation in movement: is there a mobility syndrome in the large white butterfly *Pieris brassicae*? *Ecological Entomology*, 37, 377–385.
- Ebert, G. and Rennwald, E. (1991) *Die Schmetterlinge Baden-Württembergs*. Vol. 2. Ulmer, Stuttgart.
- Fischer, K. and Fiedler, K. (2000) Sex-related differences in reaction norms in the butterfly *Lycaena tityrus* (Lepidoptera: Lycaenidae). *Oikos*, 90, 372–380.
- Fischer, K. and Fiedler, K. (2001a) Effects of adult feeding and temperature regime on fecundity and longevity in the butterfly *Lycaena hippothoe*. *Journal of the Lepidopterists Society*, 54, 91–95.
- Fischer, K. and Fiedler, K. (2001b) Effects of larval starvation on adult life-history traits in the butterfly species *Lycaena tityrus* (Lepidoptera: Lycaenidae). *Entomologia Generalis*, 25, 249–254.
- Fischer, K., Brakefield, P.M. and Zwaan, B.J. (2003) Plasticity in butterfly egg size: why larger offspring at lower temperatures? *Ecology*, 84, 3138–3147.
- Forsman, A. (1999) Temperature influence on escape behaviour in two species of pygmy grasshoppers. *Ecoscience*, 6, 214–222.
- Gibbs, M. and Weir, L. (2017) Sub-lethal viral exposure and growth on drought stressed host plants changes resource allocation patterns and life history costs in the speckled wood butterfly, *Pararge aegeria*. *Journal of Invertebrate Pathology*, 150, 106–113.
- Gilchrist, G.W. (1990) The consequences of sexual dimorphism in body size for butterfly flight and thermoregulation. *Functional Ecology*, 4, 475–487.
- Gilchrist, G.W. (1996) A quantitative genetic analysis of thermal sensitivity in the locomotor performance curve of *Aphidius ervi*. *Evolution*, 50, 1560–1572.
- Hamel, S., Côté, S.D., Gaillard, J.M. and Festa-Bianchet, M. (2009) Individual variation in reproductive costs of reproduction: high-quality females always do better. *Journal of Animal Ecology*, 78, 182–190.
- Hanski, I. and Gilpin, M.E. (1997) *Metapopulation Biology: Ecology, Genetics, and Evolution*. Academic Press, New York.
- Hickling, R., Roy, D.B., Hill, J.K., Fox, R. and Thomas, C.D. (2006) The distributions of a wide range of taxonomic groups are expanding polewards. *Global Change Biology*, 12, 450–455.
- Hill, C.J. and Pierce, N.E. (1989) The effect of adult diet on the biology of butterflies—1. The common imperial blue, *Jalmenus evagoras*. *Oecologia*, 81, 249–257.
- Huey, R.B. and Kingsolver, J.G. (1993) Evolution of resistance to high temperature in ectotherms. *The American Naturalist*, 142, S21–S46.
- Karl, I. and Fischer, K. (2008) Why get big in the cold? Towards a solution to a life-history puzzle. *Oecologia*, 155, 215–225.
- Karl, I., Hoffmann, K.H. and Fischer, K. (2010) Food stress sensitivity and flight performance across phosphoglucose isomerase enzyme genotypes in the Sooty Copper butterfly. *Population Ecology*, 52, 307–315.
- Karl, I., Janowitz, S.A. and Fischer, K. (2008) Altitudinal life-history variation and thermal adaptation in the Copper butterfly *Lycaena tityrus*. *Oikos*, 117, 778–788.
- Kehl, T., Bensch, J., Böhm, F., Kniepkamp, B.O., Leonhardt, V., Schwieger, S. et al. (2015) Fat and sassy: factors underlying male mating success in a butterfly. *Entomologia Experimentalis et Applicata*, 155, 257–265.
- Lebeau, J., Wesselingh, R.A. and Van Dyck, H. (2016) Floral resource limitation severely reduces butterfly survival, condition and flight activity in simplified agricultural landscapes. *Oecologia*, 180, 421–427.
- Legrand, D., Trochet, A., Moulherat, S., Calvez, O., Stevens, V.M., Ducatez, S. et al. (2015) Ranking the ecological causes of dispersal in a butterfly. *Ecography*, 38, 822–831.
- Marden, J.H. (1989) Bodybuilding dragonflies: costs and benefits of maximizing flight muscle. *Physiological Zoology*, 62, 505–521.
- Matthysen, E. (2012) Multicausality of dispersal: a review. *Dispersal Ecology and Evolution* (eds. J. Clobert, M. Baguette, T.G. Benton & J.M. Bullock), pp. 1–18. Oxford University Press, Oxford.
- Mattila, A.L.K. and Hanski, I. (2014) Heritability of flight and resting metabolic rates in the Glanville fritillary butterfly. *Journal of Evolutionary Biology*, 27, 1733–1743.
- McNamara, J.M. and Houston, A.I. (1996) State-dependent life histories. *Nature*, 380, 215–221.
- Merckx, T. and Van Dyck, H. (2005) Mate location behaviour of the butterfly *Pararge aegeria* in woodland and fragmented landscapes. *Animal Behaviour*, 70, 411–416.
- Merckx, T., Karlsson, B. and Van Dyck, H. (2006) Sex- and landscape-related differences in flight ability under suboptimal temperatures in a woodland butterfly. *Functional Ecology*, 20, 436–441.
- Metcalfe, N.B. and Monaghan, P. (2001) Compensation for a bad start: grow now, pay later? *Trends in Ecology and Evolution*, 16, 254–260.
- Monaghan, P. (2008) Early growth conditions, phenotypic development and environmental change. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 363, 1635–1645.

- Niitepõld, K., Smith, A.D., Osborne, J.L., Reynolds, D.R., Carreck, N.L., Martin, A.P. *et al.* (2009) Flight metabolic rate and Pgi genotype influence butterfly dispersal rate in the field. *Ecology*, 90, 2223–2232
- Pleasants, J.M. and Oberhauser, K.S. (2013) Milkweed loss in agricultural fields because of herbicide use: effect on the monarch butterfly population. *Insect Conservation and Diversity*, 6, 135–144.
- Ronce, O. (2007) How does it feel to be like a rolling stone? Ten questions about dispersal evolution. *Annual Review of Ecology, Evolution, and Systematics*, 38, 231–253.
- Saastamoinen, M., Hrai, N. and van Nouhuys, S. (2013) Direct and trans-generational responses to food deprivation during development in the Glanville fritillary butterfly. *Oecologia*, 171, 93–104.
- Saastamoinen, M., van der Sterren, D., Vastenhout, N., Zwaan, B.J. and Brakefield, P.M. (2010) Predictive adaptive responses: condition-dependent impact of adult nutrition and flight in the tropical butterfly *Bicyclus anynana*. *The American Naturalist*, 176, 686–698.
- Settele, J., Kudrna, O., Harpke, A., Kühn, I. and van Swaay, C. (2008) *Climatic Risk Atlas of European Butterflies. Biodiversity and Ecosystem Risk Assessment 1*. Pensoft, Sofia.
- Stevens, V.M., Trochet, A., Blanchet, S., Moulherat, S., Clobert, J. and Baguette, M. (2013) Dispersal syndromes and the use of life-histories to predict dispersal. *Evolutionary Applications*, 6, 630–642.
- Stoks, R., De Block, M. and McPeck, M.A. (2006) Physiological costs of compensatory growth in a damselfly. *Ecology*, 87, 1566–1574.
- Suarez, R.K. (2000) Energy metabolism during insect flight: biochemical design and physiological performance. *Physiological and Biochemical Zoology*, 73, 765–771.
- Therry, L., Zawal, A., Bonte, D. and Stoks, R. (2014) What factors shape female phenotypes of a poleward-moving damselfly at the edge of its range? *Biological Journal of the Linnean Society*, 112, 556–568.
- Thomas, C.D., Hill, J.K. and Lewis, O.T. (1998) Evolutionary consequences of habitat fragmentation in a localized butterfly. *Journal of Animal Ecology*, 67, 485–497.
- Tolman, T. and Lewington, R. (1998) *Die Tagfalter Europas und Nordwestafrikas*. Franckh-Kosmos, Stuttgart.
- Travis, J.M.J., Delgado, M., Bocedi, G., Baguette, M., Bartoń, K., Bonte, D. *et al.* (2013) Dispersal and species' responses to climate change. *Oikos*, 122, 1532–1540.
- van Dyck, H. and Baguette, M. (2005) Dispersal behaviour in fragmented landscapes: routine or special movements? *Basic and Applied Ecology*, 6, 535–545.
- van Dyck, H. and Wiklund, C. (2002) Seasonal butterfly design: morphological plasticity among three developmental pathways relative to sex, flight and thermoregulation. *Journal of Evolutionary Biology*, 15, 216–225.
- Wang, X.G., Johnson, M.W., Daane, K.M. and Opp, S. (2009) Combined effects of heat stress and food supply on flight performance of olive fruit fly (Diptera: Tephritidae). *Annals of the Entomological Society of America*, 102, 727–734.
- Warren, M.S., Hill, J.K., Thomas, J.A., Asher, J., Fox, R., Huntley, B. *et al.* (2001) Rapid responses of British butterflies to opposing forces of climate and habitat change. *Nature*, 414, 65–69.
- Wickman, P.O. (2009) Thermoregulation and habitat use in butterflies. *Ecology of Butterflies in Europe* (eds. J. Settele, T.G. Shreeve, M. Konvicka & H. van Dyck), pp. 55–61. Cambridge University Press, Cambridge.
- Wiklund, C. and Fagerström, T. (1977) Why do males emerge before females? *Oecologia*, 31, 153–158.
- Zera, A.J., Potts, J. and Kobus, K. (1998) The physiology of life-history trade-offs: experimental analysis of a hormonally induced life-history trade-off in *Gryllus assimilis*. *The American Naturalist*, 152, 7–23.

Manuscript received January 23, 2018

Final version received March 23, 2018

Accepted March 29, 2018

## Supporting Information

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

**Table S1** Developmental and morphological traits in relation to larval feeding regime (starved individuals were deprived of food for 30 h in the last larval instar) and sex in *Lycaena tityrus* (experiment 1).

**Table S2** Morphological and physiological traits in relation to adult feeding regime and sex in *Lycaena tityrus* (experiment 2).

**Fig. S1** Relationship between the tunnel and vortex test for experiment 1(A) and 2(B).

## Supporting Information

**Table S1** Developmental and morphological traits in relation to larval feeding regime (starved individuals were deprived of food for 30 h in the last larval instar) and sex in *Lycaena tityrus* (experiment 1). Given are means  $\pm$  1 SE. Group sample sizes range between 74 and 78 individuals. Different superscript letters (a, b, c, d) within a row indicate significant differences among groups (Tukey HSD after GLM). Thus, means followed by the same letter (e.g. 'a') do not differ significantly, while means followed by different letters (e.g. 'a' and 'b') do differ significantly.

Trait	Males fed	Females fed	Males starved	Females starved
Larval time (d)	19.7 $\pm$ 0.16 <sup>a</sup>	21.7 $\pm$ 0.20 <sup>b</sup>	20.0 $\pm$ 0.22 <sup>a</sup>	23.0 $\pm$ 0.16 <sup>c</sup>
Pupal time (d)	26.3 $\pm$ 0.25 <sup>ab</sup>	26.6 $\pm$ 0.32 <sup>a</sup>	25.5 $\pm$ 0.26 <sup>b</sup>	28.4 $\pm$ 0.36 <sup>c</sup>
Adult body mass (mg)	29.6 $\pm$ 0.51 <sup>a</sup>	35.0 $\pm$ 0.55 <sup>b</sup>	25.2 $\pm$ 0.50 <sup>c</sup>	29.5 $\pm$ 0.66 <sup>a</sup>
Abdomen mass (mg)	7.6 $\pm$ 0.31 <sup>a</sup>	12.1 $\pm$ 0.30 <sup>b</sup>	6.6 $\pm$ 0.18 <sup>c</sup>	10.1 $\pm$ 0.30 <sup>d</sup>
Thorax mass (mg)	15.7 $\pm$ 0.32 <sup>a</sup>	16.1 $\pm$ 0.24 <sup>a</sup>	13.0 $\pm$ 0.31 <sup>b</sup>	13.4 $\pm$ 0.34 <sup>b</sup>
Thorax-abdomen ratio	2.2 $\pm$ 0.06 <sup>a</sup>	1.4 $\pm$ 0.02 <sup>b</sup>	2.0 $\pm$ 0.05 <sup>a</sup>	1.4 $\pm$ 0.03 <sup>b</sup>
Fat (mg)	0.1 $\pm$ 0.01 <sup>a</sup>	0.2 $\pm$ 0.01 <sup>b</sup>	0.1 $\pm$ 0.01 <sup>a</sup>	0.2 $\pm$ 0.01 <sup>b</sup>
Fat (%)	6.6 $\pm$ 0.46 <sup>ac</sup>	4.6 $\pm$ 0.27 <sup>b</sup>	7.5 $\pm$ 0.55 <sup>c</sup>	5.2 $\pm$ 0.33 <sup>ab</sup>
Wing length (mm)	14.4 $\pm$ 0.07 <sup>a</sup>	14.6 $\pm$ 0.06 <sup>a</sup>	13.8 $\pm$ 0.09 <sup>b</sup>	13.8 $\pm$ 0.10 <sup>b</sup>
Wing area (mm <sup>2</sup> )	83.2 $\pm$ 1.06 <sup>a</sup>	86.1 $\pm$ 0.83 <sup>a</sup>	75.4 $\pm$ 1.13 <sup>b</sup>	76.4 $\pm$ 1.11 <sup>b</sup>
Wing loading	36.1 $\pm$ 0.84 <sup>ac</sup>	40.9 $\pm$ 0.78 <sup>b</sup>	33.9 $\pm$ 0.81 <sup>c</sup>	39.1 $\pm$ 1.04 <sup>ab</sup>
Wing aspect ratio	10.1 $\pm$ 0.10 <sup>a</sup>	10.0 $\pm$ 0.08 <sup>a</sup>	10.2 $\pm$ 0.06 <sup>a</sup>	10.0 $\pm$ 0.05 <sup>a</sup>

**Table S2** Morphological and physiological traits in relation to adult feeding regime and sex in *Lycaena tityrus* (experiment 2). Given are means  $\pm$  1 SE. Group sample sizes range between 24 and 55 individuals. Different superscript letters (a, b, c, d) within a row indicate significant differences among groups (Tukey HSD after GLM). Thus, means followed by the same letter (e.g. 'a') do not differ significantly, while means followed by different letters (e.g. 'a' and 'b') do differ significantly.

<b>Trait</b>	<b>Males fed</b>	<b>Females fed</b>	<b>Males starved</b>	<b>Females starved</b>
<b>Adult body mass (mg)</b>	34.7 $\pm$ 1.1 <sup>a</sup>	41.6 $\pm$ 1.3 <sup>b</sup>	32.1 $\pm$ 0.8 <sup>a</sup>	32.8 $\pm$ 0.8 <sup>a</sup>
<b>Abdomen mass (mg)</b>	11.3 $\pm$ 0.8 <sup>ac</sup>	19.1 $\pm$ 0.9 <sup>b</sup>	8.3 $\pm$ 0.4 <sup>a</sup>	12.3 $\pm$ 0.5 <sup>c</sup>
<b>Thorax mass (mg)</b>	17.3 $\pm$ 0.5 <sup>a</sup>	16.4 $\pm$ 0.4 <sup>ab</sup>	17.0 $\pm$ 0.8 <sup>a</sup>	15.1 $\pm$ 0.3 <sup>b</sup>
<b>Thorax-abdomen ratio</b>	1.8 $\pm$ 0.11 <sup>a</sup>	0.9 $\pm$ 0.03 <sup>b</sup>	2.1 $\pm$ 0.12 <sup>c</sup>	1.3 $\pm$ 0.03 <sup>d</sup>
<b>Fat (mg)</b>	3.1 $\pm$ 0.04 <sup>a</sup>	5.7 $\pm$ 0.06 <sup>b</sup>	1.5 $\pm$ 0.03 <sup>a</sup>	2.0 $\pm$ 0.02 <sup>a</sup>
<b>Fat (%)</b>	9.8 $\pm$ 0.8 <sup>a</sup>	8.7 $\pm$ 1.0 <sup>a</sup>	7.4 $\pm$ 1.3 <sup>ab</sup>	4.7 $\pm$ 0.3 <sup>b</sup>
<b>Wing length (mm)</b>	14.8 $\pm$ 0.1 <sup>a</sup>	14.7 $\pm$ 0.1 <sup>a</sup>	15.1 $\pm$ 0.1 <sup>a</sup>	14.7 $\pm$ 0.1 <sup>a</sup>
<b>Wing area (mm<sup>2</sup>)</b>	83.5 $\pm$ 1.2 <sup>a</sup>	89.0 $\pm$ 1.1 <sup>b</sup>	85.7 $\pm$ 1.3 <sup>ab</sup>	86.5 $\pm$ 1.0 <sup>ab</sup>
<b>Wing loading</b>	41.2 $\pm$ 1.0 <sup>a</sup>	46.5 $\pm$ 1.0 <sup>b</sup>	37.5 $\pm$ 0.6 <sup>ac</sup>	37.8 $\pm$ 0.7 <sup>c</sup>
<b>Wing aspect ratio</b>	10.5 $\pm$ 0.1 <sup>a</sup>	9.7 $\pm$ 0.1 <sup>b</sup>	10.6 $\pm$ 0.1 <sup>a</sup>	10.0 $\pm$ 0.1 <sup>c</sup>

**Figure S1** Relationship between the tunnel and vortex test for experiment 1 (A) and 2 (B).




Fig. S1A




Fig. S1B



4.4 Successful despite poor flight performance: range expansion is associated with enhanced exploratory behavior and fast development

Submitted as:

Reim, E., Blesinger, S., Förster, L., Fischer, K. (2018) Successful despite poor flight performance: range expansion is associated with enhanced exploratory behavior and fast development. *Journal of Evolutionary Biology*, in press.

Accepted: 15.05.2018



1       **Successful despite poor flight performance: range**  
2       **expansion is associated with enhanced exploratory**  
3       **behaviour and fast development**

4  
5       **Elisabeth Reim\*, Simone Blesinger, Lisa Förster & Klaus Fischer**

6  
7                   Zoological Institute and Museum, Greifswald University,  
8                                   D-17489 Greifswald, Germany

9  
10  
11   **\* Corresponding author:**

12   Elisabeth Reim

13   ORCID-ID: 0000-0003-4139-2330

14   Zoological Institute and Museum

15   Greifswald University

16   Soldmannstraße 14

17   D-17489 Greifswald, Germany

18   Tel.: +49-3834-4204297

19   Fax: +49-3834-4204252

20   E-Mail: [elisabeth.reim@uni-greifswald.de](mailto:elisabeth.reim@uni-greifswald.de)

21  
22   **Running head:** Factors underlying range expansion

24 **Acknowledgements**

25 We thank Toomas Esperk and Toomas Tammaru for helpful suggestions and for  
26 sharing their knowledge about the occurrence of *L. tityrus* in Estonia with us, Jan D.  
27 Roy for assistance during field work, and Marissa K. Dowalter for assistance during  
28 the experiments. This research was funded by the DFG research training group  
29 RESPONSE (DFG GRK 2010).

30

31 **Abstract**

32 Anthropogenic interference forces species to respond to changing environmental  
33 conditions. One possible response is dispersal and concomitant range shifts, allowing  
34 individuals to escape unfavourable conditions or to track the shifting climate niche.  
35 Range expansions depend on both dispersal capacity and the ability to establish  
36 populations beyond the former range. We here compare well-established core  
37 populations with recently established edge populations in the currently northward  
38 expanding butterfly *Lycaena tityrus*. Edge populations were characterized by shorter  
39 development times and smaller size, a higher sensitivity to high temperature, and an  
40 enhanced exploratory behaviour. The differences between core and edge populations  
41 found suggest adaptation to local climates and an enhanced dispersal ability in edge  
42 populations. In particular, enhanced exploratory behaviour may be advantageous in all  
43 steps of the dispersal process and may have facilitated the current range expansion.  
44 This study describes differences associated with a current range expansion,  
45 knowledge which might be useful for a better understanding of species responses to  
46 environmental change. We further report on variation between males and females in  
47 morphology and flight behaviour, with males showing a longer flight endurance and  
48 more pronounced exploratory behaviour than females.

49

50 **Keywords:** Dispersal; global change; flight ability; life-history; local adaptation; *Lycaena*  
51 *tityrus*; range shift.

52

## 53 **Introduction**

54 Human-induced environmental change like the current increase in the Earth's mean  
55 temperature are expected to strongly affect ecosystems and biodiversity (Hansen *et*  
56 *al.*, 2012; Stanton *et al.*, 2015). Especially changes in temperature may prove to be of  
57 utmost importance, because temperature is one of the most important ecological  
58 factors affecting, for instance, species performance, survival, and distributions  
59 (Overgaard & Sørensen, 2008; Sunday *et al.*, 2011). Facing changing environmental  
60 conditions, the long-term survival of all organisms depends on their ability to respond  
61 to such changes either by (1) phenotypic plasticity, which allows species to tolerate  
62 novel conditions, (2) genetic adaptation, or (3) movement to more suitable habitats  
63 often resulting in range shifts (Davis *et al.*, 2005; Parmesan, 2006; Berg *et al.*, 2010;  
64 Hofmann & Todgham, 2010). The principal mechanism underlying range shifts is  
65 dispersal, often defined as any movements potentially leading to gene flow (Ronce,  
66 2007). It constitutes a key process in the colonization of new areas to track shifting  
67 climate niches, is central to the spatial dynamics of populations and metapopulations,  
68 and affects patterns of local adaptation (Hanski, 1999; Doebeli & Dieckmann, 2003;  
69 Hickling *et al.*, 2006; Ronce, 2007; Baguette *et al.*, 2013).

70

71 In general, dispersing individuals do not seem to comprise a random subset of a given  
72 population. Rather, dispersers often possess several correlated life-history traits  
73 forming a so called dispersal syndrome discriminating between dispersing and non-  
74 dispersing conspecifics (Bonte & Saastamoinen, 2012; Stevens *et al.*, 2013). Traits  
75 potentially affecting dispersal include morphology, physiology, and overall condition  
76 (Bowler & Benton, 2005; Bonte *et al.*, 2012; Therry *et al.*, 2014) as well as behavioural  
77 traits such as exploration, aggression, and sociability (Dingemanse *et al.*, 2003;

78 Duckworth & Badyaev, 2007; Cote *et al.*, 2010b; Ducatez *et al.*, 2012). Given that new  
79 populations are likely founded by the most dispersive individuals (cf. Hill *et al.*, 1998;  
80 Thomas *et al.*, 1998; Hanski *et al.*, 2002, 2004) and that dispersal-related traits are  
81 heritable (Roff & Fairbairn, 2001), more dispersive genotypes are expected to  
82 accumulate in recently established edge populations (spatial sorting; Shine *et al.*,  
83 2011). Life-history traits associated with dispersal ability are often affected by  
84 environmental conditions. For instance, suboptimal conditions (e.g. during  
85 development) may decrease the investment into costly dispersal traits (Bonte *et al.*,  
86 2008). On the other hand, such conditions may increase dispersal ability or at least the  
87 willingness to do so as an adaptive response for dealing with detrimental conditions  
88 (e.g. Monaghan, 2008). Environmental effects on dispersal ability / tactics have been  
89 repeatedly demonstrated through alterations in body composition or behaviour (Bonte  
90 *et al.*, 2008; Saastamoinen *et al.*, 2010; Van den Heuvel *et al.*, 2013).

91

92 However, dispersal ability is only one facet of range expansions. The other crucial  
93 component is the ability to establish populations beyond the current range once  
94 suitable habitats have been reached by dispersing individuals. For instance,  
95 temperature variation along geographical gradients may require an 'optimization' of  
96 phenotypic values and thus local adaptation (Hoffmann *et al.*, 2002; Castañeda *et al.*,  
97 2005). Hence, successful range expansions may further depend on a wide tolerance  
98 against environmental stressors and / or the evolutionary potential of a given species.  
99 Consequently, phenotypic differences between recently established edge and long-  
100 established core populations are expected in traits related to dispersal, but also in other  
101 traits such as thermal tolerance (Hanski *et al.*, 2004; Calosi *et al.*, 2008; Lindstrom *et*  
102 *al.*, 2013). This is due to differences in selection pressures but may also result from

103 differences in the genetic structure of core and edge populations (Volis *et al.*, 1998;  
104 Purves, 2009; Mägi *et al.*, 2011). Such differentiation may also include phenotypic  
105 plasticity in those traits (Otaki *et al.*, 2010; Pichancourt & van Klinken, 2012).

106

107 Depending on dispersal and establishing capacity some species may benefit while  
108 others may suffer from current climate change (Pimm, 2001; Thomas *et al.*, 2001;  
109 Klockmann *et al.*, 2016). Many temperate zone ectotherms may actually benefit from  
110 rising temperatures, because conditions at northern margins may become more  
111 suitable (Pimm, 2001; Deutsch *et al.*, 2008; Bestion *et al.*, 2015). One such species is  
112 the currently northward expanding butterfly *Lycaena tityrus* (Brunzel *et al.*, 2008;  
113 Settele *et al.*, 2008; Klockmann *et al.*, 2016), which has colonized north-eastern  
114 Estonia very recently. We here compared replicated core (Germany) and recently  
115 established edge (Estonia) populations of this temperate-zone butterfly. In order to  
116 explore variation in dispersal ability and life history indicative of local adaptation, we  
117 investigated a variety of traits ranging from development, morphology, physiology, and  
118 flight endurance through to behaviour. As differences in dispersal ability can be related  
119 to flight performance (Ducatez *et al.*, 2012; Legrand *et al.*, 2015) and / or morphology  
120 (Berwaerts & van Dyck, 2004), we focused on traits potentially associated with flight  
121 and dispersal ability, including wing size, body mass, thorax-abdomen ratio, wing  
122 aspect ratio, wing loading, and storage reserves (Betts & Wootton, 1988; Berwaerts *et*  
123 *al.*, 2002). Many of these traits might also show environmental-induced plasticity, which  
124 may in turn feedback on dispersal ability. We here tested the following hypotheses: (1)  
125 Estonian edge populations will differ from German core populations in traits indicative  
126 of local adaptation to cooler conditions and shorter vegetation period. Specifically, we  
127 predict a more rapid development in edge populations. (2) Edge populations show

128 increased dispersal ability, evidenced by a higher flight endurance and enhanced  
129 exploratory behaviour. Increased dispersal ability is associated with morphological and  
130 / or physiological differentiation among edge and core populations.

131

132

## 133 **Material and methods**

### 134 **Study organism**

135 *Lycaena tityrus* (Poda, 1761) is a widespread temperate-zone butterfly with a range  
136 from Western Europe to central Asia (Ebert & Rennwald, 1991). It is currently  
137 expanding its range towards higher latitudes and altitudes, which is assumed to be  
138 largely driven by anthropogenic climate change (Brunzel *et al.*, 2008; Settele *et al.*,  
139 2008). The species is bivoltine with two discrete generations per year in most parts of  
140 its range, although populations with one or three generations per year occur.  
141 Overwintering takes place as half-grown larvae. The principal larval host-plant is  
142 *Rumex acetosa* L. (Polygonaceae), but some congeneric *Rumex* species are utilized  
143 as well (Ebert & Rennwald, 1991; Tolman & Lewington, 1998; Settele *et al.*, 2008).  
144 Adults are nectar feeders, using a wide array of species including several Asteraceae  
145 (Ebert & Rennwald, 1991).

146

147 The butterflies used in this study were collected in three replicated populations each in  
148 north-eastern Germany and Estonia. While the German populations were located  
149 within the core of the distribution area, the Estonian ones reflect currently expanding  
150 edge populations, which were probably established within the last 5 years (T. Esperk  
151 & T. Tammaru, pers. comm.). The first records of *L. tityrus* in Estonia stem from the  
152 end of the 20<sup>th</sup> century (Õunap & Tartes, 2014). In total 69 and 56 freshly eclosed,

153 fecund females were caught in May and June 2016 in Germany and Estonia,  
154 respectively (Germany: Guest 54°02'56.4" N, 13°26'18.3" E, n = 18; Ueckermünde  
155 53°44'52.6" N, 14°15'28.3" E, n = 25; Waren 53°30'13.3" N, 12°42'31.5" E, n = 26;  
156 Estonia: Karepa 59°32'39.3" N, 26°25'34.4" E, n = 22; Kasispea 59°35'05.6" N,  
157 25°46'21.9" E, n = 12; Mahu 59°31'09.8" N, 26°42'58.5" E, n = 22). All these  
158 populations represent bivoltine lowland populations. Both regions differ substantially in  
159 annual mean temperature (Germany: 8.2°C, Estonia: 4.8°C), but less so in annual  
160 precipitation (Germany: 560 mm, Estonia: 606 mm). The temperature differences  
161 prevail throughout the time during which the first generation develops and reproduces  
162 (April: 6.6 vs 3.6°C, May: 11.5 vs 10.3°C June: 15.4 vs 14.9°C; climate-data.org). All  
163 females were transferred to a climate chamber at Greifswald University for egg laying.

164

### 165 **Experimental design**

166 Caught females were kept at 26°C, 60% relative humidity, and a L18:D6 photoperiod.  
167 For oviposition, females were placed separately in translucent 1 L plastic pots covered  
168 with gauze, and were provided with *R. acetosa* (oviposition substrate), fresh flowers  
169 (*Achillea millefolium* L., *Leucanthemum vulgare*, *Hypochaeris radicata* L., *Tanacetum*  
170 *vulgare* L.), water, and a highly concentrated sucrose solution (20 vol%) for adult  
171 feeding. Deposited eggs were collected daily and transferred, separated by female, to  
172 small plastic boxes. For each female, resulting larvae (maximum number of 20 per  
173 female) were divided into two treatment groups. The first group of larvae was reared  
174 under the same conditions used for oviposition, while the second group experienced  
175 cooler conditions (18°C, 60% relative humidity, L18:D6 photoperiod). All individuals  
176 were kept at their respective temperature treatment throughout. Larvae were  
177 individually reared in small plastic boxes (125 ml) and had access to food in ample

178 supply until pupation. Boxes contained moistened filter paper and fresh cuttings of *R.*  
179 *acetosa* and *R. acetosella*. Pupae were weighed on the day following pupation. For  
180 each individual we recorded larval and pupal development time and additionally  
181 calculated larval growth rate (pupal mass divided by larval time). Resulting butterflies  
182 were kept individually and were provided with moistened cotton for drinking. One day-  
183 old butterflies were first subjected to a behavioural experiment testing the exploratory  
184 behaviour and on the following day to a flight endurance test (see below). Exploratory  
185 behaviour, relying on spontaneous flight, was tested in a climatic chamber at 26°C and  
186 60% relative humidity. Flight endurance (forced flight), in contrast, was tested at 22°C  
187 and 60% relative humidity (note that no differences were found between 22 and 26°C  
188 in pilot experiments). All butterflies could acclimatise for 1 hour to the conditions in the  
189 experimental chambers prior to the experiments. Unfortunately, a high number of  
190 Estonian larvae raised at 18°C entered diapause which caused low sample sizes in  
191 the respective treatment groups.

192

### 193 **Exploratory behaviour**

194 This test was designed to investigate the individual's exploratory behaviour towards  
195 a food source in an unknown experimental setup. Each individual was placed singly  
196 into the back-right corner of a cage (50 x 50 x 50 cm). A food source consisting of  
197 flowering plants (see above) and two leaves of the host plant (*R. acetosa*) was placed  
198 in a small vial in the front-left corner of the cage. Butterflies were released and  
199 observed for longest 90 minutes. It was scored if the butterfly reached the food source  
200 or not. Afterwards, butterflies were kept individually at their rearing conditions until  
201 the start of the flight endurance test the next day.

202

203 **Flight endurance test**

204 Butterflies were tested for flight endurance using a well-established assay (cf. Ducatez  
205 *et al.*, 2012, 2013). This test was designed to detect individual differences in flight  
206 endurance under stressful conditions and has been shown to be correlated with other  
207 mobility measures and dispersal (Ducatez *et al.* 2012, Legrand *et al.* 2015). Each  
208 individual was placed into a plastic chamber (30 x 16 x 14 cm), which was perforated  
209 at its base and fixed to a rapid agitator (IKA Vortex 4 digital). After a habituation period  
210 of 30 seconds, the vortex was switched on to strongly shake the chamber, preventing  
211 the butterfly from holding on to the walls. This method forces the butterflies to fly during  
212 the test, as they were otherwise lying uncomfortably on the bottom of the shaking  
213 chamber. The time an individual butterfly spent flying was recorded for 60 seconds,  
214 with higher values reflecting a higher flight endurance. After the endurance test,  
215 butterflies were frozen for later analyses of potentially flight-related parameters.

216

217 **Morphological and physiological analyses**

218 To test for associations between exploratory behaviour or flight endurance and other  
219 traits, we measured a couple of morphological and physiological traits. First, adult body  
220 mass was determined to the nearest 0.01 mg (Sartorius CPA225D). Then, wings,  
221 heads and legs were removed. Thorax and abdomen were separated and afterwards  
222 weighed. Abdomen fat content was measured following Fischer *et al.* (2003), but using  
223 the less poisonous acetone instead of dichloromethane. Abdomens were first dried to  
224 constant weight for two days at 70°C. Abdomen dry mass was measured. Afterwards,  
225 fat was extracted twice for 48 h each, using 1 ml of acetone (C<sub>3</sub>H<sub>6</sub>O) for each abdomen.  
226 Solutions were exchanged between both fat extractions. Then, abdomens were again  
227 dried for two days at 70°C, after which the fat-free dry mass was measured. Absolute

228 (mg) and relative (%) abdomen fat content were determined as the mass difference  
229 between abdomen dry mass and the remaining dry mass after the two fat extractions.  
230 Forewing area and length (from basis to apex) were measured using digital images of  
231 left forewings (captured from ventrally with a digital camera mounted on a stereo  
232 microscope) and NIS elements software. Wing loading was calculated as total body  
233 mass divided by forewing area, and wing aspect ratio as  $4 \times \text{forewing length}^2$  divided  
234 by the forewing area (Berwaerts *et al.*, 2002). Thorax-abdomen ratio was calculated  
235 by dividing thorax through abdomen mass.

236

### 237 **Statistical analyses**

238 Data on developmental, morphological, and physiological traits were analysed with  
239 general linear mixed models (GLMs), using country (origin), developmental  
240 temperature and sex as fixed factors, and population (nested within country) as a  
241 random effect. Due to small sample sizes in the 18°C treatment groups for Estonian  
242 individuals, the factor family was only included in the additional models (see below).  
243 Data on abdomen mass, thorax-abdomen ratio, and absolute and relative fat content  
244 were LN transformed prior to analysis to meet GLM requirements. Data on exploratory  
245 behaviour (reaching of food source: yes / no) were analysed using generalized linear  
246 models (GzLMs) with a binomial error distribution and logit-link function, using the  
247 same factors as above.

248

249 To additionally test for the impact of continuous variables (morphology, physiology) on  
250 flight behaviour and endurance, we first performed principal component analyses  
251 (PCAs) owing to strong inter-correlations among traits. PCAs were calculated  
252 separately for males and females, as sexes differed strongly in a variety of traits. We

253 used the first four principal components (PCs) for further analyses, having Eigenvalues  
254 between 4.3 and 1.1 for males and between 5.0 and 1.0 for females (cf. Table 1). Thus,  
255 all PCs explaining  $\geq 10\%$  of the variance were included. We then constructed another  
256 set of models by including the respective PCs and the results of the behavioural or  
257 flight endurance test as covariates to the models mentioned above, followed by a  
258 stepwise backwards elimination of non-significant factors.

259

260 Due to small sample sizes in the 18°C treatment groups for Estonian individuals, we  
261 additionally calculated models (same structure as above, but additionally including  
262 family nested within population and country as a random factor) exclusively based on  
263 (1) the animals reared at 26°C, thus focusing on differences across countries  
264 (appendix A1), and (2) on the animals from Germany, thus focusing on differences  
265 across developmental temperature (18°C versus 26°C, appendix A2). As these  
266 additional analyses confirmed the results obtained from the above full-factorial models,  
267 only the latter ones are presented here (for additional models please see appendix A1  
268 and A2). All statistical tests were performed with Statistica 8.0 (Tulsa, StatSoft, OK).  
269 All means are given  $\pm 1$  SE.

270

271

## 272 **Results**

### 273 **Developmental and morphological traits**

274 Country significantly affected larval and pupal development time, pupal mass, adult  
275 body mass, wing length, wing area and aspect ratio (Table 2). German compared with  
276 Estonian individuals had longer larval and pupal development times, higher pupal and  
277 adult body masses, larger wings, and lower wing aspect ratios (Fig. 1). Differences

278 among populations were significant for all traits except for wing length, wing area and  
279 aspect ratio (cf. appendix A3). Developmental temperature significantly affected all  
280 traits except for thorax mass, relative fat content, wing length and area, indicating that  
281 higher temperatures during development led to a shorter development time, higher  
282 growth rate, higher pupal but lower abdomen and adult body mass, higher thorax-  
283 abdomen ratio, lower absolute fat content, wing loading and aspect ratio (Table 2, Fig.  
284 1). Significant sex differences were found for all traits except from pupal mass and  
285 wing length, indicating that females had on average a longer development time, lower  
286 growth rate, higher absolute fat content, wing loading, abdomen and concomitantly  
287 total body mass, lower thorax mass, thorax-abdomen ratio, relative abdomen fat  
288 content, and wing aspect ratio, and a larger wing area than males (Table 2, Fig. 1).

289  
290 Country by temperature interactions were significant for all traits except for larval time,  
291 thorax-abdomen ratio, and wing length. They show that temperature effects on pupal  
292 mass were restricted to German populations but absent in Estonian ones (Fig. 1c).  
293 Similar patterns of positive effects of the higher temperature on German but negative  
294 ones on Estonian individuals were found for thorax mass (Fig. 1g), wing area (Fig. 1l),  
295 and relative fat content (Fig. 1j). For adult body mass, the significant interaction  
296 indicates that the difference between temperatures was more pronounced in Estonian  
297 compared with German individuals (Fig. 1e). The same pattern was found for abdomen  
298 mass (Fig. 1f), absolute fat content (Fig. 1i), and wing loading (Fig. 1m). For pupal time,  
299 the difference between temperatures was more pronounced in German than in  
300 Estonian individuals (Fig. 1b), which was also the case in aspect ratio (Fig. 1n).

301

302 Country by sex interactions were significant for pupal mass, thorax mass, and wing  
303 length and area. Accordingly, sexual differences in pupal mass were restricted to  
304 German populations (Fig. 1c). A similar pattern was found for wing length (Fig. 1k). For  
305 thorax mass, the interaction shows that the difference between sexes was more  
306 pronounced in German than in Estonian individuals (Fig. 1g). For wing area, in  
307 contrast, the difference between the sexes was more pronounced in Estonian  
308 compared with German individuals (Fig. 1l).

309

310 Temperature by sex interactions were significant for pupal mass, pupal time, and wing  
311 length. For pupal mass and pupal time, the difference between the sexes was more  
312 pronounced at 26°C than at 18°C (Fig. 1b, 1c). The opposite pattern though was found  
313 for wing length (Fig. 1k). The three-way interaction, finally, was significant for wing  
314 length only. It shows that differences among countries persisted in all treatment groups  
315 except for females at 18°C (Fig. 1k).

316

### 317 **Exploratory behaviour and flight endurance**

318 Exploratory behaviour was significantly affected by the factors country, sex, and  
319 temperature (Table 3a). Estonian butterflies, males, and individuals that were raised at  
320 26°C were more successful in reaching the food source (Fig. 2a). Including principal  
321 components and the flight endurance in the vortex as covariates revealed significant  
322 effects of the country by temperature interaction, PC1 and the vortex test results for  
323 males, and of temperature for females (Table 3a). The country by temperature  
324 interaction in males shows that the temperature effect on exploratory behaviour was  
325 more pronounced in Estonian (18°C: 41.7 vs 26°C: 70.1%) than in German males  
326 (18°C: 38.0 vs 26°C: 64.5%). In males, flight behaviour was negatively correlated with

327 PC1 (beta  $-0.11 \pm 0.05$ ) and the vortex test results (beta  $-0.14 \pm 0.04$ ). PC1 was most  
328 strongly (positively) related to body mass (Table 1a), meaning that a lower body mass  
329 increased exploratory behaviour in males. Furthermore, males that were more  
330 exploratory performed worse in the vortex. The temperature effect found in females  
331 reflects the above pattern that individuals having been raised at 26°C were more  
332 exploratory.

333

334 Flight endurance was significantly affected by the factors population, sex, and the  
335 interactions between country and sex and temperature and sex (Table 3b). Males  
336 showed a higher flight endurance than females (Fig. 2b). The interaction between  
337 country and sex though shows that the higher flight endurance of males was largely  
338 restricted to German animals. The temperature by sex interaction indicates that males  
339 had a higher flight endurance when raised at 18°C only, while for individuals raised at  
340 26°C values were very similar (Fig. 2b).

341

342 Including principal components and exploratory behaviour as covariates revealed  
343 significant effects of population, temperature, PC1, PC3, and exploratory behaviour for  
344 males, and of temperature, PC1, and PC4 for females (Table 3b). Males had a higher  
345 flight endurance when having been raised at 18°C rather than 26°C. Flight endurance  
346 was positively correlated with PC1 (beta  $0.29 \pm 0.04$ ) and PC3 (beta  $0.11 \pm 0.04$ ). PC1  
347 was most strongly (positively) correlated with body mass (Table 1a), meaning that a  
348 higher body mass increased flight endurance. PC3 was most strongly (positively)  
349 correlated with thorax-abdomen ratio (Table 1a), showing that a high thorax-abdomen  
350 ratio led to a higher flight endurance in males. Males that showed reduced exploratory  
351 behaviour performed better in the vortex. Females, in contrast to males, had a higher

352 flight endurance when having been raised at 26°C rather than 18°C. Flight endurance  
353 was negatively correlated with PC1 (beta  $-0.17 \pm 0.04$ ) and PC4 (beta  $-0.11 \pm 0.04$ ).  
354 PC1 in turn was most strongly (negatively) correlated with adult body mass (Table 1b),  
355 meaning that a high body mass led to a higher flight endurance. PC4 was most strongly  
356 (negatively) correlated with aspect ratio (Table 1b), showing that a higher aspect ratio  
357 increased flight endurance in the vortex. Country also tended to affect flight endurance  
358 of females, showing that Estonian females tended to perform better than German  
359 females.

360

361

## 362 **Discussion**

363 We start our discussion with the effects of developmental temperature, country (origin),  
364 and sex on variation in developmental and morphological traits and continue with  
365 exploratory behaviour and flight endurance, as the former may causally underlie  
366 variation in the latter.

367

### 368 **Developmental and morphological traits**

369 Interestingly, Estonian individuals had shorter development times accompanied by  
370 lower body size compared with German individuals. Short development time typically  
371 evolves when a minimum size needs to be reached within a limited time period (Case,  
372 1978; Arendt, 1997). More severe time constraints in Estonia as compared with  
373 Germany certainly arise from a shorter growing season with increasing latitude  
374 (Blanckenhorn, 1997). Thus, the cooler conditions experienced in Estonia likely  
375 selected for shorter development time in order to compensate for the limitations  
376 imposed by the cooler environmental conditions. Such local adaptation is especially

377 expected in insects exhibiting more than one generation per year, driven by the need  
378 to fit in yet another generation (Fischer & Karl, 2010). Although local adaptation has  
379 been repeatedly shown in butterflies (e.g. Ayres & Scriber, 1994; Phillimore *et al.*,  
380 2012), the case reported here seems interesting as differences may have evolved  
381 within a relatively short period of time, as the sampled Estonian populations were  
382 established only recently. We also found negative effects of the higher developmental  
383 temperature on thorax mass, wing area, and relative fat content in Estonian butterflies,  
384 while German individuals benefitted from warmer conditions. This might further support  
385 the notion of local adaptation to the cooler Estonian climate or at least indicate a loss  
386 of adaptation to warmer conditions. Note furthermore the higher incidence of diapause  
387 induction in Estonian animals at the cooler temperature (Estonia: 80%; Germany: 5%),  
388 suggesting a higher threshold for direct development in Estonian individuals. Taken  
389 together, the above lines of evidence suggest that Estonian individuals are relatively  
390 better adapted to cooler and German ones to warmer environmental conditions.

391  
392 Regarding temperature effects, the higher temperature reduced development times  
393 and adult body mass (following the temperature-size rule), as expected for an  
394 ectothermic organism (Nylin, 1989; Atkinson, 1994; Blanckenhorn, 1997; Fischer &  
395 Fiedler, 2000). Shorter development times are generally achieved through enhanced  
396 growth and metabolic rates at higher temperatures (Karl & Fischer, 2008). Higher  
397 abdomen and thus adult body mass (resulting in a higher absolute fat content and wing  
398 loading, but a lower thorax-abdomen ratio) at lower temperatures typically result from  
399 behavioural and physiological mechanisms like an increased food intake and a more  
400 efficient conversion of the ingested food into body matter (Karl & Fischer, 2008; Fischer  
401 & Karl, 2010). Note though that, contrary to the temperature-size rule (Nylin, 1989;

402 Atkinson, 1994), pupal mass was higher when being reared at higher temperatures  
403 (except for the pupal mass of Estonian females), which may have been caused by the  
404 higher growth rates. As these entail increased physiological costs that lead to high  
405 weight losses during metamorphosis (Gotthard *et al.*, 1994; Fischer *et al.*, 2005;  
406 Fischer & Karl, 2010), the expected pattern nevertheless arose in the adult stage.

407

408 The sex differences in developmental, morphological, and physiological traits found in  
409 this study are in line with previous studies on (*Lycaena*) butterflies (e.g. Karl *et al.*,  
410 2008; Fischer & Karl, 2010; Saastamoinen *et al.*, 2012). Male butterflies had shorter  
411 development times and higher growth rates (and an accordingly earlier adult  
412 emergence) to enhance mating opportunities (selection for protandry; Wiklund &  
413 Fagerström, 1977; Fischer & Fiedler, 2000; Karl & Fischer 2008), but were lighter as  
414 adults compared with females (fecundity selection in females; Honek, 1993;  
415 Blanckenhorn, 2000). In German males, this pattern also applied to pupal mass,  
416 suggesting that shorter development times decreased pupal weight. Interestingly,  
417 pupal mass of Estonian individuals was almost identical in males and females, perhaps  
418 reflecting the strong time pressure to develop fast in both sexes. Regarding flight-  
419 related traits, males had a lower wing loading but a higher thorax-abdomen ratio, wing  
420 aspect ratio, and relative fat content, traits that are often associated with increased  
421 flight ability, acceleration capacity, and maneuverability (Betts & Wootton, 1988;  
422 Berwaerts *et al.*, 2002, 2006; Van Dyck & Wiklund, 2002; Saastamoinen *et al.*, 2012).

423

#### 424 **Exploratory behaviour and flight endurance**

425 Exploratory behaviour, tested here as the ability to locate a food source in an unknown  
426 experimental setup, was more pronounced in butterflies from expanding edge

427 populations (Estonia), though effects were largely restricted to individuals from the  
428 warmer temperature. However, good exploration skills may help during all steps of the  
429 dispersal process, beginning with the actual decision to emigrate via effective  
430 movement in the transition phase through to the location of suitable habitats (Fraser *et*  
431 *al.*, 2001; Ronce, 2007; Cote *et al.*, 2010a; Ducatez *et al.*, 2012). Within populations,  
432 more explorative individuals can have a higher dispersal propensity (Cote *et al.*, 2010a;  
433 Ducatez *et al.*, 2012), which may lead to an accumulation of bolder individuals in edge  
434 populations as indicated here. Interestingly, butterflies reared at 26°C were more  
435 exploratory compared with butterflies reared at 18°C. Although direct behavioural  
436 responses to the thermal conditions experienced during development are poorly  
437 documented, temperature might be a reliable indicator of optimal dispersal tactics later  
438 in life (Tautz *et al.*, 2003; Bonte *et al.*, 2008). Thus, having experienced beneficial  
439 conditions during development may indicate suitable conditions for dispersal and thus  
440 increase exploratory behaviour. This might be especially relevant in Estonia due to  
441 stronger temperature constraints, which may explain the above interaction.

442

443 An alternative explanation for the positive effect of higher temperatures could be that  
444 the butterflies were more attracted to the food source. Higher temperatures (and  
445 accordingly higher metabolic rates) during development and in the adult stage prior to  
446 the experiment could have increased their need for drinking and feeding. This would  
447 be in line with the morphological differences found, showing that butterflies reared at  
448 the higher temperature had a lower body mass and fat content. This notion is  
449 additionally supported by the negative correlation between exploratory behaviour and  
450 PC1 in males, demonstrating that lower body masses increased exploratory behaviour.  
451 Furthermore, males that were more explorative had a shorter flight endurance and vice

452 versa. The overall weak effects of morphological and physiological traits on exploratory  
453 behaviour suggest that it is not closely related to the individuals' condition, but rather  
454 a behavioural property which is modulated by environmental circumstances (e.g.  
455 developmental temperatures).

456

457 Males were in general more exploratory than females, which might be a result of  
458 differential selective pressures (Gilchrist, 1990; Van Dyck & Wiklund, 2002; Merckx &  
459 Van Dyck, 2005; Berwaerts *et al.*, 2006). After emergence, males try to quickly  
460 establish a territory to enhance mating opportunities (Davies, 1978; Fischer & Fiedler,  
461 2000; Saastamoinen *et al.*, 2012), which could explain their more exploratory  
462 behaviour evidenced by a faster location of a food source.

463

464 German males showed a higher flight endurance in the vortex test than Estonian  
465 males. Mechanistically, this difference might be linked to a higher body mass and size  
466 of German individuals. Interestingly, Estonian females tended to show the opposite  
467 pattern, viz. a higher flight endurance compared with German females. Likewise,  
468 Hanski *et al.* (2004) found in the butterfly *Melitaea cinxia* that females from newly  
469 established and isolated populations were most dispersive. Males showed a higher  
470 flight endurance than females, but only when having been raised at 18°C. As males  
471 have a lower wing loading but a higher thorax-abdomen ratio, a higher flight endurance  
472 of males was expected (Wickman, 2009; Saastamoinen *et al.*, 2012). In contrast to  
473 exploratory behaviour, morphological traits seemed to have a stronger impact on flight  
474 endurance (significant impacts of 2 PCs for each males and females), which suggests  
475 that it is more tightly linked to morphology than is the actual behaviour (Van Dyck &  
476 Wiklund, 2002; Berwaerts *et al.*, 2006). In both sexes, flight endurance was positively

477 related to body size and therefore storage reserves, which are used to fuel flight and  
478 reflect a good condition in general (Zera *et al.*, 1998; Berwaerts *et al.*, 2002; Karl &  
479 Fischer, 2008). In males, we also found a positive influence of a higher thorax-  
480 abdomen ratio, which is generally assumed to enhance flight performance and  
481 acceleration capacity (Marden, 1989; Thomas *et al.*, 1998; Berwaerts *et al.*, 2002). In  
482 females, flight endurance was additionally related to high wing aspect ratios, reflecting  
483 increased flight ability (Berwaerts *et al.*, 2002; Berwaerts & Van Dyck, 2004).  
484 Population effects were significant in most cases, indicating genetic variation among  
485 and within populations in the traits investigated (Finger *et al.*, 2009; Klockmann *et al.*,  
486 2016). This did not interfere with any of the conclusions presented here.

487

## 488 **Conclusions**

489 We here show that sexes differed substantially in developmental, morphological, and  
490 physiological traits, being arguably the result of differential selective pressures (e.g.  
491 Gilchrist, 1990; Berwaerts *et al.*, 2006). In general, males showed traits associated  
492 with increased flight ability, which resulted in a higher flight endurance in the vortex  
493 and enhanced exploratory behaviour, which might be helpful to quickly establish a  
494 territory (Fischer & Fiedler, 2000; Saastamoinen *et al.*, 2012). In the current era of  
495 climate change, many temperate-zone ectotherms might actually benefit from rising  
496 global temperatures (Deutsch *et al.*, 2008). One such example is the currently  
497 northward expanding butterfly *L. tityrus*, which has recently expanded in north-eastern  
498 direction and thereby colonized Estonia. The colonization process in turn obviously  
499 depends on dispersal ability, and may warrant local adaptation to new environmental  
500 conditions (Doebeli & Dieckmann, 2003). Despite having colonized Estonia only very  
501 recently, we detected several differences between edge and core populations

502 indicative of local adaptation. Estonian edge as compared with German core  
503 populations were characterized by shorter development times and smaller size, a  
504 higher sensitivity to high temperature, a higher diapause incidence, and enhanced  
505 exploratory behaviour. These differences seem to partly reflect adaptations to the  
506 cooler Estonian climate and the concomitantly short vegetation period (or a loss of  
507 adaptation to warmer conditions), while others, namely enhanced exploratory  
508 behaviour, may indicate increased dispersal capacity which may have partly facilitated  
509 the current range expansion. While the time frame available may not have been  
510 sufficient for *in situ* adaptation, it is currently unclear whether the differences between  
511 core and edge populations reflect clinal variation (assuming that the populations from  
512 which Estonia was colonized already differed from those in Germany) or an  
513 accumulation of cold-adapted and / or more dispersive genotypes in edge populations.  
514 Note that butterfly females mate quickly after adult eclosion (Hill *et al.*, 1999) and lay  
515 at least a proportion of their eggs after dispersal, hence enabling colonization (Stevens  
516 *et al.*, 2012). Also, we cannot rule out a contribution of parental effects, which may  
517 comprise part of adaptive responses to environmental variation (Wolf *et al.*, 1998;  
518 Mattila & Hanski, 2014). In summary, this study provides insights into the factors  
519 associated with range expansion, knowledge which might be useful for a better  
520 understanding of species responses to environmental change.

521

522

### 523 **Conflict of interests**

524 The authors have no conflict of interests to declare.

525

526

527 **References**

- 528 Arendt, J.D. 1997. Adaptive intrinsic growth rates: an integration across taxa. *Q. Rev.*  
529 *Biol.* **72**: 149–177.
- 530 Atkinson, D. 1994. Temperature and organism size – a biological law for ectotherms?  
531 *Adv. Ecol. Res.* **25**: 1–58.
- 532 Ayres, M.P. & Scriber, J.M. 1994. Local adaptation to regional climates in *Papilio*  
533 *canadensis* (Lepidoptera: Papilionidae). *Ecol. Monogr.* **64**: 465–482.
- 534 Baguette, M., Blanchet, S., Legrand, D., Stevens, V.M. & Turlure, C. 2013. Individual  
535 dispersal, landscape connectivity and ecological networks. *Biol. Rev.* **88**: 310–  
536 326.
- 537 Berg, M.P., Toby Kiers, E., Driessen, G., van der Heijden, M., Kooi, B.W., Kuenen, F.,  
538 *et al.* 2010. Adapt or disperse: understanding species persistence in a changing  
539 world. *Glob. Change Biol.* **16**: 587–598.
- 540 Berwaerts, K., Aerts, P. & Van Dyck, H. 2006. On the sex-specific mechanisms of  
541 butterfly flight: flight performance relative to flight morphology, wing kinematics,  
542 and sex in *Pararge aegeria*. *Biol. J. Linn. Soc.* **89**: 675–687.
- 543 Berwaerts, K. & Van Dyck, H. 2004. Take-off performance under optimal and  
544 suboptimal thermal conditions in the butterfly *Pararge aegeria*. *Oecologia* **141**:  
545 536–545.
- 546 Berwaerts, K., Van Dyck, H. & Aerts, P. 2002. Does flight morphology relate to flight  
547 performance? An experimental test with the butterfly *Pararge aegeria*. *Funct. Ecol.*  
548 **16**: 484–491.
- 549 Bestion, E., Clobert, J. & Cote, J. 2015. Dispersal response to climate change: scaling  
550 down to intraspecific variation. *Ecol. Lett.* **18**: 1226–1233.
- 551 Betts, C.R. & Wootton, R.J. 1988. Wing shape and flight behaviour in butterflies

- 552 (Lepidoptera: Papilionoidea and Hesperioidea): a preliminary analysis. *J. Exp.*  
553 *Biol.* **138**: 271–288.
- 554 Blanckenhorn, W.U. 1997. Altitudinal life history variation in the dung flies  
555 *Scathophaga stercoraria* and *Sepsis cynipsea*. *Oecologia* **109**: 342–352.
- 556 Blanckenhorn, W.U. 2000. The evolution of body size. What keeps organisms small?  
557 *Q. Rev. Biol.* **75**: 385–407.
- 558 Bonte, D. & Saastamoinen, M. 2012. Dispersal syndromes in butterflies and spiders.  
559 In: *Dispersal ecology and evolution* (J. Clobert *et al.*, eds), pp. 161–170. Oxford  
560 University Press, Oxford.
- 561 Bonte, D., Travis, J.M.J., De Clercq, N., Zwertvaegher, I. & Lens, L. 2008. Thermal  
562 conditions during juvenile development affect adult dispersal in a spider. *Proc.*  
563 *Natl. Acad. Sci. USA* **105**: 17000–17005.
- 564 Bonte, D., Van Dyck, H., Bullock, J.M., Coulon, A., Delgado, M., Gibbs, M., *et al.* 2012.  
565 Costs of dispersal. *Biol. Rev.* **87**: 290–312.
- 566 Bowler, D.E. & Benton, T.G. 2005. Causes and consequences of animal dispersal  
567 strategies: relating individual behaviour to spatial dynamics. *Biol. Rev.* **80**: 205–  
568 225.
- 569 Brunzel, S., Bussmann, M. & Obergruber, H. 2008. Deutliche Veränderungen von  
570 Tagfalterzönosen als Folge von Ausbreitungsprozessen. Erste Ergebnisse eines  
571 Monitorings über 17 Jahre. *Nat. Landsch.* **6**: 280–287.
- 572 Calosi, P., Bilton, D.T. & Spicer, J.I. 2008. Thermal tolerance, acclimatory capacity and  
573 vulnerability to global climate change. *Biol. Lett.* **4**: 99–102.
- 574 Case, T.J. 1978. On the evolution and adaptive significance of postnatal growth rates  
575 in the terrestrial vertebrates. *Q. Rev. Biol.* **53**: 243–282.
- 576 Castañeda, L.E., Lardies, M.A. & Bozinovic, F. 2005. Interpopulational variation in

577 recovery time from chill coma along a geographic gradient: a study in the common  
578 woodlouse, *Porcellio laevis*. *J. Insect Physiol.* **51**: 1346–1351.

579 Cote, J., Clobert, J., Brodin, T., Fogarty, S. & Sih, A. 2010a. Personality-dependent  
580 dispersal: characterization, ontogeny and consequences for spatially structured  
581 populations. *Philos. Trans. R. Soc. B* **365**: 4065–4076.

582 Cote, J., Fogarty, S., Weinersmith, K., Brodin, T. & Sih, A. 2010b. Personality traits  
583 and dispersal tendency in the invasive mosquitofish (*Gambusia affinis*). *Proc. R.*  
584 *Soc. B* **277**: 1571–1579.

585 Davies, N.B. 1978. Territorial defence in the speckled wood butterfly (*Pararge aegeria*):  
586 the resident always wins. *Anim. Behav.* **26**: 138–147.

587 Davis, M.B., Shaw, R.G. & Etterson, J.R. 2005. Evolutionary responses to changing  
588 climate. *Ecology* **86**: 1704–1714.

589 Deutsch, C.A., Tewksbury, J.J., Huey, R.B., Sheldon, K.S., Ghalambor, C.K., Haak,  
590 D.C., *et al.* 2008. Impacts of climate warming on terrestrial ectotherms across  
591 latitude. *Proc. Natl. Acad. Sci. USA* **105**: 6668–6672.

592 Dingemanse, N.J., Both, C., van Noordwijk, A.J., Rutten, A.L. & Drent, P.J. 2003. Natal  
593 dispersal and personalities in great tits (*Parus major*). *Proc. R. Soc. B* **270**: 741–  
594 747.

595 Doebeli, M. & Dieckmann, U. 2003. Speciation along environmental gradients. *Nature*  
596 **421**: 259–264.

597 Ducatez, S., Baguette, M., Trochet, A., Chaput-Bardy, A., Legrand, D., Stevens, V., *et*  
598 *al.* 2013. Flight endurance and heating rate vary with both latitude and habitat  
599 connectivity in a butterfly species. *Oikos* **122**: 601–611.

600 Ducatez, S., Legrand, D., Chaput-Bardy, A., Stevens, V.M., Fréville, H. & Baguette, M.  
601 2012. Inter-individual variation in movement: is there a mobility syndrome in the

- 602 large white butterfly *Pieris brassicae*? *Ecol. Entomol.* **37**: 377–385.
- 603 Duckworth, R.A. & Badyaev, A. V. 2007. Coupling of dispersal and aggression  
604 facilitates the rapid range expansion of a passerine bird. *Proc. Natl. Acad. Sci.*  
605 *USA* **104**: 15017–15022.
- 606 Ebert, G. & Rennwald, E. 1991. *Die Schmetterlinge Baden-Württembergs*, Vol. 2.  
607 Ulmer, Stuttgart.
- 608 Finger, A., Schmitt, T., Emmanuel Zachos, F., Meyer, M., Assmann, T. & Christian  
609 Habel, J. 2009. The genetic status of the violet copper *Lycaena helle* - a relict of  
610 the cold past in times of global warming. *Ecography* **32**: 382–390.
- 611 Fischer, K. & Fiedler, K. 2000. Sex-related differences in reaction norms in the butterfly  
612 *Lycaena tityrus* (Lepidoptera: Lycaenidae). *Oikos* **90**: 372–380.
- 613 Fischer, K. & Karl, I. 2010. Exploring plastic and genetic responses to temperature  
614 variation using copper butterflies. *Clim. Res.* **43**: 17–30.
- 615 Fischer, K., Brakefield, P.M. & Zwaan, B.J. 2003. Plasticity in butterfly egg size: Why  
616 larger offspring at lower temperatures? *Ecology* **84**: 3138–3147.
- 617 Fischer, K., Zeilstra, I., Hetz, S.K. & Fiedler, K. 2005. Physiological costs of growing  
618 fast: does accelerated growth reduce pay-off in adult fitness? *Evol. Ecol.* **18**: 343–  
619 353.
- 620 Fraser, D.F., Gilliam, J.F., Daley, M.J., Le, A.N. & Skalski, G.T. 2001. Explaining  
621 leptokurtic movement distributions: intrapopulation variation in boldness and  
622 exploration. *Am. Nat.* **158**: 124–135.
- 623 Gilchrist, G.W. 1990. The consequences of sexual dimorphism in body size for butterfly  
624 flight and thermoregulation. *Funct. Ecol.* **4**: 475–487.
- 625 Gotthard, K., Nylin, S. & Wiklund, C. 1994. Adaptive variation in growth rate: life history  
626 costs and consequences in the speckled wood butterfly, *Pararge aegeria*.

- 627 *Oecologia* **99**: 281–289.
- 628 Hansen, J., Sato, M. & Ruedy, R. 2012. Perception of climate change. *Proc. Natl. Acad.*  
629 *Sci. USA* **109**: E2415–E2423.
- 630 Hanski, I. 1999. Habitat connectivity, habitat continuity, and metapopulations in  
631 dynamic landscapes. *Oikos* **87**: 209–219.
- 632 Hanski, I., Breuker, C.J., Schops, K., Setchfield, R. & Nieminen, M. 2002. Population  
633 history and life history influence the migration rate of female Glanville fritillary  
634 butterflies. *Oikos* **98**: 87–97.
- 635 Hanski, I., Erälahti, C., Kankare, M., Ovaskainen, O. & Sirén, H. 2004. Variation in  
636 migration propensity among individuals maintained by landscape structure. *Ecol.*  
637 *Lett.* **7**: 958–966.
- 638 Hickling, R., Roy, D.B., Hill, J.K., Fox, R. & Thomas, C.D. 2006. The distributions of a  
639 wide range of taxonomic groups are expanding polewards. *Glob. Change Biol.* **12**:  
640 450–455.
- 641 Hill, J.K., Thomas, C.D. & Blakeley, D.S. 1999. Evolution of flight morphology in a  
642 butterfly that has recently expanded its geographic range. *Oecologia* **121**: 165–  
643 170.
- 644 Hill, J.K., Thomas, C.D. & Lewis, O.T. 1998. Flight morphology in fragmented  
645 populations of a rare British butterfly, *Hesperia comma*. *Biol. Conserv.* **87**: 277–  
646 283.
- 647 Hoffmann, A.A., Anderson, A. & Hallas, R. 2002. Opposing clines for high and low  
648 temperature resistance in *Drosophila melanogaster*. *Ecol. Lett.* **5**: 614–618.
- 649 Hofmann, G.E. & Todgham, A.E. 2010. Living in the now: physiological mechanisms  
650 to tolerate a rapidly changing environment. *Annu. Rev. Physiol.* **72**: 127–145.
- 651 Honek, A. 1993. Intraspecific variation in body size and fecundity in insects: a general

- 652 relationship. *Oikos* **66**: 483–492.
- 653 Karl, I. & Fischer, K. 2008. Why get big in the cold? Towards a solution to a life-history  
654 puzzle. *Oecologia* **155**: 215–225.
- 655 Karl, I., Janowitz, S.A. & Fischer, K. 2008. Altitudinal life-history variation and thermal  
656 adaptation in the copper butterfly *Lycaena tityrus*. *Oikos* **117**: 778–788.
- 657 Klockmann, M., Karajoli, F., Kuczyk, J., Reimer, S. & Fischer, K. 2016. Fitness  
658 implications of simulated climate change in three species of copper butterflies  
659 (Lepidoptera: Lycaenidae). *Biol. J. Linn. Soc.* **120**: 125–143.
- 660 Legrand, D., Trochet, A., Moulherat, S., Calvez, O., Stevens, V.M., Ducatez, S., *et al.*  
661 2015. Ranking the ecological causes of dispersal in a butterfly. *Ecography* **38**:  
662 822–831.
- 663 Lindstrom, T., Brown, G.P., Sisson, S.A., Phillips, B.L. & Shine, R. 2013. Rapid shifts  
664 in dispersal behavior on an expanding range edge. *Proc. Natl. Acad. Sci. USA*  
665 **110**: 13452–13456.
- 666 Mägi, M., Semchenko, M., Kalamees, R. & Zobel, K. 2011. Limited phenotypic  
667 plasticity in range-edge populations: a comparison of co-occurring populations of  
668 two *Agrimonia* species with different geographical distributions. *Plant Biol.* **13**:  
669 177–184.
- 670 Marden, J.H. 1989. Bodybuilding dragonflies: costs and benefits of maximizing flight  
671 muscle. *Physiol. Zool.* **62**: 505–521.
- 672 Mattila, A.L.K. & Hanski, I. 2014. Heritability of flight and resting metabolic rates in  
673 the Glanville fritillary butterfly. *J. Evol. Biol.* **27**: 1733–1743.
- 674 Merckx, T. & Van Dyck, H. 2005. Mate location behaviour of the butterfly *Pararge*  
675 *aegeria* in woodland and fragmented landscapes. *Anim. Behav.* **70**: 411–416.
- 676 Monaghan, P. 2008. Early growth conditions, phenotypic development and

677 environmental change. *Philos. Trans. R. Soc. London B* **363**: 1635–1645.

678 Nylin, S. 1989. Effects of changing photoperiods in the life cycle regulation of the  
679 comma butterfly, *Polygonia c-album* (Nymphalidae). *Ecol. Entomol.* **14**: 209–218.

680 Otaki, J.M., Hiyama, A., Iwata, M. & Kudo, T. 2010. Phenotypic plasticity in the range-  
681 margin population of the lycaenid butterfly *Zizeeria maha*. *BMC Evol. Biol.* **10**: 252.

682 Õunap, E. & Tartes, U. 2014. *Eesti päevaliblikad [Estonian butterflies]*. Varrak, Tallinn.

683 Overgaard, J. & Sørensen, J.G. 2008. Rapid thermal adaptation during field  
684 temperature variations in *Drosophila melanogaster*. *Cryobiology* **56**: 159–162.

685 Parmesan, C. 2006. Ecological and evolutionary responses to recent climate change.  
686 *Annu. Rev. Ecol. Evol. Syst.* **37**: 637–669.

687 Phillimore, A.B., Stålhandske, S., Smithers, R.J. & Bernard, R. 2012. Dissecting the  
688 contributions of plasticity and local adaptation to the phenology of a butterfly and  
689 its host plants. *Am. Nat.* **180**: 655–670.

690 Pichancourt, J.B. & van Klinken, R.D. 2012. Phenotypic plasticity influences the size,  
691 shape and dynamics of the geographic distribution of an invasive plant. *PLoS One*  
692 **7**:e32323.

693 Pimm, S.L. 2001. Entrepreneurial insects. *Nature* **411**: 531–532.

694 Purves, D.W. 2009. The demography of range boundaries versus range cores in  
695 eastern US tree species. *Proc. R. Soc. B* **276**: 1477–1484.

696 Roff, D.A. & Fairbairn, D.J. 2001. The genetic basis of dispersal and migration, and its  
697 consequences for the evolution of correlated traits. In: *Dispersal* (J. Clobert *et al.*,  
698 eds), pp. 191–202. Oxford University Press, Oxford.

699 Ronce, O. 2007. How does it feel to be like a rolling stone? Ten questions about  
700 dispersal evolution. *Annu. Rev. Ecol. Evol. Syst.* **38**: 231–253.

701 Saastamoinen, M., Brakefield, P.M. & Ovaskainen, O. 2012. Environmentally induced

702 dispersal-related life-history syndrome in the tropical butterfly, *Bicyclus anynana*.  
703 *J. Evol. Biol.* **25**: 2264–2275.

704 Saastamoinen, M., van der Sterren, D., Vastenhout, N., Zwaan, B.J. & Brakefield, P.M.  
705 2010. Predictive adaptive responses: condition-dependent impact of adult  
706 nutrition and flight in the tropical butterfly *Bicyclus anynana*. *Am. Nat.* **176**: 686–  
707 98.

708 Settele, J., Kudrna, O., Harpke, A., Kühn, I., Van Swaay, C., Verovnik, R., *et al.* 2008.  
709 *Climatic risk atlas of European butterflies. Biodiversity and ecosystem risk*  
710 *assessment 1*. Pensoft, Sofia.

711 Shine, R., Brown, G.P. & Phillips, B.L. 2011. An evolutionary process that assembles  
712 phenotypes through space rather than through time. *Proc. Natl. Acad. Sci.* **108**:  
713 5708–5711.

714 Stanton, J.C., Shoemaker, K.T., Pearson, R.G. & Akçakaya, H.R. 2015. Warning times  
715 for species extinctions due to climate change. *Glob. Change Biol.* **21**: 1066–1077.

716 Stevens, V.M., Trochet, A., Blanchet, S., Moulherat, S., Clobert, J. & Baguette, M.  
717 2013. Dispersal syndromes and the use of life-histories to predict dispersal. *Evol.*  
718 *Appl.* **6**: 630–642.

719 Stevens, V.M., Trochet, A., Van Dyck, H., Clobert, J. & Baguette, M. 2012. How is  
720 dispersal integrated in life histories: a quantitative analysis using butterflies. *Ecol.*  
721 *Lett.* **15**: 74–86.

722 Sunday, J.M., Bates, A.E. & Dulvy, N.K. 2011. Global analysis of thermal tolerance  
723 and latitude in ectotherms. *Proc. R. Soc. B* **278**: 1823–1830.

724 Tautz, J., Maier, S., Groh, C., Rössler, W. & Brockmann, A. 2003. Behavioral  
725 performance in adult honey bees is influenced by the temperature experienced  
726 during their pupal development. *Proc. Natl. Acad. Sci. USA* **100**: 7343–7347.

- 727 Therry, L., Zawal, A., Bonte, D. & Stoks, R. 2014. What factors shape female  
728 phenotypes of a poleward-moving damselfly at the edge of its range? *Biol. J. Linn.*  
729 *Soc.* **112**: 556–568.
- 730 Thomas, C.D., Bodsworth, E.J., Wilson, R.J., Simmons, A.D., Davies, Z.G., Musche,  
731 M., *et al.* 2001. Ecological and evolutionary processes at expanding range  
732 margins. *Nature* **411**: 577–581.
- 733 Thomas, C.D., Hill, J.K. & Lewis, O.T. 1998. Evolutionary consequences of habitat  
734 fragmentation in a localized butterfly. *J. Anim. Ecol.* **67**: 485–497.
- 735 Tolman, T. & Lewington, R. 1998. *Die Tagfalter Europas und Nordwestafrikas.*  
736 Franckh-Kosmos, Stuttgart.
- 737 Van den Heuvel, J., Saastamoinen, M., Brakefield, P.M., Kirkwood, T.B.L., Zwaan, B.J.  
738 & Shanley, D.P. 2013. The predictive adaptive response: modeling the life-history  
739 evolution of the butterfly *Bicyclus anynana* in seasonal environments. *Am. Nat.*  
740 **181**: E28-42.
- 741 Van Dyck, H. & Wiklund, C. 2002. Seasonal butterfly design: morphological plasticity  
742 among three developmental pathways relative to sex, flight and thermoregulation.  
743 *J. Evol. Biol.* **15**: 216–225.
- 744 Volis, S., Mendlinger, S., Olsvig-Whittaker, L., Safriel, U.N. & Orlovsky, N. 1998.  
745 Phenotypic variation and stress resistance in core and peripheral populations of  
746 *Hordeum spontaneum*. *Biodivers. Conserv.* **7**: 799–813.
- 747 Wickman, P.O. 2009. Thermoregulation and habitat use in butterflies. In: *Ecology of*  
748 *Butterflies in Europe* (J. Settele *et al.*, eds), pp. 55–61. Cambridge University  
749 Press, Cambridge.
- 750 Wiklund, C. & Fagerström, T. 1977. Why do males emerge before females? *Oecologia*  
751 **31**: 153–158.

752 Wolf, J.B., Brodie, E.D., Cheverud, J.M., Moore, A.J. & Wade, M.J. 1998.  
753 Evolutionary consequences of indirect genetic effects. *Trends Ecol. Evol.* **13**:  
754 64–69.

755 Zera, A.J., Potts, J. & Kobus, K. 1998. The physiology of life-history trade-offs:  
756 experimental analysis of a hormonally induced life-history trade-off in *Gryllus*  
757 *assimilis*. *Am. Nat.* **152**: 7–23.

758

759 **Tables**

760

761 **Table 1** Results of principal component analyses for males (a) and females (b)  
 762 including Eigen values, percentage of the variance explained, cumulative percentage  
 763 of the variance explained, and *r*-values of correlations with continuous variables (rows  
 764 4-13) for principal components 1-4. *r*-values > 0.7 in bold.

765

766 Table 1a

No	Parameter	PC1	PC2	PC3	PC4
1	Eigen value	4.3	2.0	1.8	1.1
2	Variance explained (%)	43.3	19.7	17.6	11.2
3	Cumulative variance (%)	43.3	63.0	80.5	91.8
4	Adult body mass	<b>0.989</b>	0.038	-0.042	0.065
5	Thorax mass	<b>0.795</b>	-0.070	-0.582	0.044
6	Abdomen mass	<b>0.901</b>	0.099	0.272	0.081
7	Thorax-abdomen ratio	-0.093	0.167	<b>0.903</b>	0.017
8	Fat (mg)	0.231	<b>-0.950</b>	0.183	-0.014
9	Fat (%)	0.068	<b>-0.962</b>	0.248	-0.037
10	Wing Length	<b>0.793</b>	0.206	0.357	-0.282
11	Wing area	<b>0.803</b>	0.194	0.377	0.168
12	Wing loading	<b>0.750</b>	-0.123	-0.400	-0.057
13	Wing aspect ratio	0.034	0.041	-0.018	<b>-0.998</b>

767

No	Parameter	PC1	PC2	PC3	PC4
1	Eigen value	5.1	2.0	1.3	1.0
2	Variance explained (%)	51.2	19.8	12.5	10.5
3	Cumulative variance (%)	51.2	71.0	83.5	94.0
4	Adult body mass	<b>-0.983</b>	0.118	-0.015	0.075
5	Thorax mass	<b>-0.943</b>	-0.069	-0.275	0.146
6	Abdomen mass	<b>-0.809</b>	0.363	0.304	-0.043
7	Thorax-abdomen ratio	0.559	0.398	0.605	-0.239
8	Fat (mg)	-0.505	<b>-0.790</b>	0.291	-0.179
9	Fat (%)	-0.242	<b>-0.870</b>	0.353	-0.235
10	Wing Length	<b>-0.798</b>	0.381	0.145	-0.349
11	Wing area	<b>-0.816</b>	0.345	0.352	-0.006
12	Wing loading	<b>-0.827</b>	-0.116	-0.343	0.117
13	Wing aspect ratio	0.028	0.098	-0.496	<b>-0.860</b>

770 **Table 2** General linear mixed model results for the effects of country, temperature, sex  
771 (fixed factors), and population (nested within country; random factor) on developmental  
772 and morphological traits in *Lycaena tityrus*. Models were constructed by a stepwise  
773 backwards elimination of non-significant factors. Eta-squared represents effect size.  
774 Significant *P*-values are given in bold.

775

<b>Larval time</b>	<b><i>MS</i></b>	<b><i>DF</i></b>	<b><i>Eta</i><sup>2</sup></b>	<b><i>F</i></b>	<b><i>P</i></b>
Country	769.3	1, 5	0.92	49.6	<b>0.0014</b>
Population (Country)	20.8	4, 1483	0.02	7.4	<b>&lt; 0.0001</b>
Temperature	99102.6	1, 1483	0.96	35374.3	<b>&lt; 0.0001</b>
Sex	1635.3	1, 1483	0.28	583.7	<b>&lt; 0.0001</b>
Error	2.8	1483			
<b>Pupal time</b>	<b><i>MS</i></b>	<b><i>DF</i></b>	<b><i>Eta</i><sup>2</sup></b>	<b><i>F</i></b>	<b><i>P</i></b>
Country	128.1	1, 5	0.94	97.4	<b>&lt; 0.0001</b>
Population (Country)	3.0	4, 1481	0.02	8.8	<b>&lt; 0.0001</b>
Temperature	7371.4	1, 1481	0.93	21293.6	<b>&lt; 0.0001</b>
Sex	6.8	1, 1481	0.01	19.6	<b>&lt; 0.0001</b>
Country*Temperature	34.9	1, 1481	0.06	100.7	<b>&lt; 0.0001</b>
Temperature*Sex	9.7	1, 1481	0.02	28.2	<b>&lt; 0.0001</b>
Error	0.4	1481			
<b>Pupal mass</b>	<b><i>MS</i></b>	<b><i>DF</i></b>	<b><i>Eta</i><sup>2</sup></b>	<b><i>F</i></b>	<b><i>P</i></b>
Country	9115	1, 5	0.717	13.7	<b>0.0121</b>
Population (Country)	1605	4, 1479	0.029	11.1	<b>&lt; 0.0001</b>
Temperature	1131	1, 1479	0.005	7.8	<b>0.0053</b>
Country*Temperature	2670	1, 1479	0.012	18.4	<b>&lt; 0.0001</b>
Country*Sex	2014	1, 1479	0.009	13.9	<b>0.0002</b>
Temperature*Sex	795	1, 1479	0.002	5.5	<b>0.0194</b>
3-way-interaction	476	1, 1479	0.004	3.3	0.0704
Error	145	1479			

776

<b>Growth rate</b>	<b>MS</b>	<b>DF</b>	<b>Eta<sup>2</sup></b>	<b>F</b>	<b>P</b>
Country	1.7	1, 5	0.05	0.2	0.6555
Population (Country)	10.3	4, 1482	0.84	12.3	< 0.0001
Temperature	6363.3	1, 1482	0.03	7584.7	< 0.0001
Sex	187.9	1, 1482	0.13	224.0	< 0.0001
Country*Sex	0.1	1, 1482	< 0.01	0.2	0.6880
Error	0.8	1482			
<b>Adult body mass</b>	<b>MS</b>	<b>DF</b>	<b>Eta<sup>2</sup></b>	<b>F</b>	<b>P</b>
Country	607.9	1, 5	0.637	13.4	0.0070
Population (Country)	87.8	4, 1360	0.013	4.4	0.0016
Temperature	571.5	1, 1360	0.167	28.5	< 0.0001
Sex	5459.9	1, 1360	0.007	272.7	< 0.0001
Country*Temperature	189.5	1, 1360	0.021	9.5	0.0021
Error	20.0	1360			
<b>Abdomen mass</b>	<b>MS</b>	<b>DF</b>	<b>Eta<sup>2</sup></b>	<b>F</b>	<b>P</b>
Country	0.50	1, 5	0.46	5.6	0.0520
Population (Country)	0.19	4, 1360	0.02	5.8	0.0001
Temperature	3.41	1, 1360	0.07	106.5	< 0.0001
Sex	109.29	1, 1360	0.72	3415.5	< 0.0001
Country*Temperature	0.31	1, 1360	0.01	9.5	0.0021
Error	0.03	1360			
<b>Thorax mass</b>	<b>MS</b>	<b>DF</b>	<b>Eta<sup>2</sup></b>	<b>F</b>	<b>P</b>
Country	17.4	1, 5	0.117	0.7	0.4406
Population (Country)	58.3	4, 1359	0.033	11.4	< 0.0001
Sex	749.0	1, 1359	0.098	147.1	< 0.0001
Country*Temperature	54.8	1, 1359	0.016	10.8	< 0.0001
Country*Sex	118.9	1, 1359	0.017	23.3	< 0.0001
Error	5.1	1359			

<b>Thorax-abdomen ratio</b>	<b><i>MS</i></b>	<b><i>DF</i></b>	<b><i>Eta</i><sup>2</sup></b>	<b><i>F</i></b>	<b><i>P</i></b>
Country	0.86	1, 5	0.36	2.6	0.1741
Population (Country)	0.45	4, 1360	0.02	6.9	<b>&lt; 0.0001</b>
Temperature	20.30	1, 1360	0.19	311.0	<b>&lt; 0.0001</b>
Sex	401.29	1, 1360	0.82	6149.0	<b>&lt; 0.0001</b>
Country*Sex	0.16	1, 1360	< 0.01	2.4	0.1179
Error	0.07	1360			
<b>Fat (mg)</b>	<b><i>MS</i></b>	<b><i>DF</i></b>	<b><i>Eta</i><sup>2</sup></b>	<b><i>F</i></b>	<b><i>P</i></b>
Country	1.9	1, 5	0.18	0.3	0.5854
Population (Country)	15.2	4, 1313	0.11	27.1	<b>&lt; 0.0001</b>
Temperature	9.6	1, 1313	0.01	17.0	<b>&lt; 0.0001</b>
Sex	62.4	1, 1313	0.07	110.9	<b>&lt; 0.0001</b>
Country*Temperature	10.0	1, 1313	< 0.01	17.7	<b>&lt; 0.0001</b>
Error	0.6	1313			
<b>Fat (%)</b>	<b><i>MS</i></b>	<b><i>DF</i></b>	<b><i>Eta</i><sup>2</sup></b>	<b><i>F</i></b>	<b><i>P</i></b>
Country	239.6	1, 5	0.080	0.4	0.5610
Population (Country)	1711.7	4, 1314	0.090	32.4	<b>&lt; 0.0001</b>
Temperature	154.5	1, 1314	0.002	2.9	0.0875
Sex	7459.5	1, 1314	0.097	141.2	<b>&lt; 0.0001</b>
Country*Temperature	229.3	1, 1314	0.003	4.3	<b>0.0374</b>
Error	52.8	1314			
<b>Wing length</b>	<b><i>MS</i></b>	<b><i>DF</i></b>	<b><i>Eta</i><sup>2</sup></b>	<b><i>F</i></b>	<b><i>P</i></b>
Country	65.2	1, 1364	0.118	181.7	<b>&lt; 0.0001</b>
Country*Sex	12.6	1, 1364	0.025	35.2	<b>&lt; 0.0001</b>
Temperature*Sex	2.3	1, 1364	0.005	6.3	<b>0.0121</b>
3-way Interaction	1.8	1, 1364	0.004	4.9	<b>0.0267</b>
Error	0.4	1364			

<b>Wing area</b>	<b><i>MS</i></b>	<b><i>DF</i></b>	<b><i>Eta</i><sup>2</sup></b>	<b><i>F</i></b>	<b><i>P</i></b>
Country	7818	1, 1364	0.11	170.5	< <b>0.0001</b>
Sex	12738	1, 1364	0.17	277.7	< <b>0.0001</b>
Country*Sex	340	1, 1364	0.01	7.4	<b>0.0065</b>
Country*Temperature	1511	1, 1364	0.02	33.0	< <b>0.0001</b>
Error	46	1364			
<b>Wing loading</b>	<b><i>MS</i></b>	<b><i>DF</i></b>	<b><i>Eta</i><sup>2</sup></b>	<b><i>F</i></b>	<b><i>P</i></b>
Country	17.1	1, 5	0.043	0.2	0.6671
Population (Country)	200.5	4, 1360	0.057	20.6	< <b>0.0001</b>
Temperature	957.5	1, 1360	0.067	98.3	< <b>0.0001</b>
Sex	854.7	1, 1360	0.061	87.7	< <b>0.0001</b>
Country*Temperature	63.0	1, 1360	0.005	6.5	<b>0.0111</b>
Error	9.7	1360			
<b>Wing aspect ratio</b>	<b><i>MS</i></b>	<b><i>DF</i></b>	<b><i>Eta</i><sup>2</sup></b>	<b><i>F</i></b>	<b><i>P</i></b>
Country	1.5	1, 5	0.010	13.7	<b>0.0002</b>
Temperature	2.5	1, 1364	0.016	22.6	< <b>0.0001</b>
Sex	262.8	1, 1364	0.632	2342.0	< <b>0.0001</b>
Country*Temperature	1.2	1, 1364	0.008	11.0	<b>0.0010</b>
Error	0.1	1364			

783 **Table 3** Results of a generalized linear model (exploratory behaviour, a) and a general  
784 linear mixed model (flight endurance in a vortex, b) for the effects of country,  
785 temperature, sex (all fixed), and population (nested within country; random) in the  
786 butterfly *Lycaena tityrus* (models 1). Second models were constructed by additionally  
787 including PCs 1-4 and the results of the exploratory behaviour or flight endurance test  
788 as covariates. All models were constructed by a stepwise backwards elimination of  
789 non-significant factors. The latter models were constructed separately for males and  
790 females, owing to strong variation in morphological traits among the sexes. Odds ratio  
791 (a) and Eta-squared (b) represent effect sizes. Significant *P*-values are given in bold.

792

793 Table 3a

<b>Model 1</b>	<b><i>DF</i></b>	<b><i>Odds ratio</i></b>	<b><i>Wald Chi<sup>2</sup></i></b>	<b><i>P</i></b>
Country	1	1.3	4.1	<b>0.0440</b>
Temperature	1	2.7	55.8	<b>&lt; 0.0001</b>
Sex	1	0.4	43.8	<b>&lt; 0.0001</b>
<b>Model 2 Males</b>	<b><i>DF</i></b>	<b><i>Odds ratio</i></b>	<b><i>Wald Chi<sup>2</sup></i></b>	<b><i>P</i></b>
Country*Temperature	3		23.9	<b>&lt; 0.0001</b>
PC1	1	1.2	5.3	<b>0.0218</b>
Flight endurance	1	1.0	10.7	<b>0.0011</b>
<b>Model 2 Females</b>	<b><i>DF</i></b>	<b><i>Odds ratio</i></b>	<b><i>Wald Chi<sup>2</sup></i></b>	<b><i>P</i></b>
Temperature	1	2.8	28.6	<b>&lt; 0.0001</b>

794

795 Table 3b

<b>Model 1</b>	<b>MS</b>	<b>DF</b>	<b>Eta<sup>2</sup></b>	<b>F</b>	<b>P</b>
Country	882	1, 5	0.149	0.8	0.4086
Population (Country)	1450	4, 1354	0.016	5.6	<b>0.0002</b>
Sex	5708	1, 1354	0.016	22.1	<b>&lt; 0.0001</b>
Country*Sex	3816	1, 1354	0.011	14.8	<b>0.0001</b>
Temperature*Sex	3953	1, 1354	0.022	15.3	<b>&lt; 0.0001</b>
Error	258	1354			
<b>Model 2 Males</b>	<b>MS</b>	<b>DF</b>	<b>Eta<sup>2</sup></b>	<b>F</b>	<b>P</b>
Country	52	1, 5	0.013	0.1	0.7543
Population (Country)	910	4, 551	0.027	3.8	<b>0.0043</b>
Temperature	4583	1, 551	0.034	19.4	<b>&lt; 0.0001</b>
PC1	6470	1, 551	0.047	27.3	<b>&lt; 0.0001</b>
PC3	5623	1, 551	0.041	23.8	<b>&lt; 0.0001</b>
Exploratory behaviour	3093	1, 551	0.023	13.1	<b>0.0003</b>
Error	237	551			
<b>Model 2 Females</b>	<b>MS</b>	<b>DF</b>	<b>Eta<sup>2</sup></b>	<b>F</b>	<b>P</b>
Country	927	1, 5	0.006	3.8	0.0511
Temperature	3815	1, 614	0.025	15.7	<b>&lt; 0.0001</b>
PC1	5129	1, 614	0.033	21.1	<b>&lt; 0.0001</b>
PC4	1506	1, 614	0.010	6.2	<b>0.0130</b>
Error	243	614			

796

797 **Figure legends**


798

799 **Fig. 1** Larval time (a), pupal time (b), pupal mass (c), larval growth rate (d), adult body  
800 mass (e), abdomen mass (f), thorax mass (g), thorax-abdomen ratio (h), absolute fat  
801 content (i), relative fat content (j), wing length (k), wing area (l), wing loading (m), and  
802 aspect ratio (n) in *Lycaena tityrus* males (filled bars) and females (open bars) in relation  
803 to origin (country) and developmental temperature. Given are means  $\pm$  1 SE. In some  
804 cases the error bars are not visible. Group sample sizes range between 38 and 540  
805 individuals. Different superscript letters above bars indicate significant differences  
806 among country by temperature groups (Tukey HSD after GLM).


807

808 **Fig. 2** Exploratory behaviour (animals finding a food source within 90 minutes in %, a)  
809 and flight endurance in a vortex (b) in *Lycaena tityrus* males (filled bars) and females  
810 (open bars) in relation to origin (country) and developmental temperature. Given are  
811 means  $\pm$  1 SE. In some cases the error bars are not visible.

812




815 Fig. 1a



818 Fig. 1b


820



821

822 Fig. 1c

823




824

825 Fig. 1d

826


827



828

829 Fig. 1e

830




831

832 Fig. 1f

833


834



835

836 Fig. 1g

837




838

839 Fig. 1h

840


841



842

843 Fig. 1i

844




845

846 Fig. 1j

847


848



849

850 Fig. 1k

851




852

853 Fig. 1l

854


855



856

857 Fig. 1m

858




859

860 Fig. 1n

861


862



863

864 Fig. 2a

865



866

867 Fig. 2b

## Appendices

**A1: Tables 1 and 2 are exclusively based on the animals reared at 26°C, thus focusing on differences across countries.**

**Table 1** General linear mixed model results for the effects of country, sex (fixed factors), population (nested within country), and family (nested within population and country, random factors) on developmental and morphological traits in *Lycaena tityrus*. Significant *P*-values are given in bold. Models were constructed by a stepwise backwards elimination of non-significant factors.

<b>Larval time</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Country	440.0	1, 5	48.7	<b>0.0021</b>
Population (Country)	9.4	4, 936	6.0	<b>0.0021</b>
Sex	506.0	1, 936	325.1	<b>&lt; 0.0001</b>
Error	1.6	936		
<b>Pupal time</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Country	57.2	1, 5	242.1	<b>&lt; 0.0001</b>
Sex	22.4	1, 940	94.8	<b>&lt; 0.0001</b>
Error	0.2	940		
<b>Pupal mass</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Country	32071	1, 5	19.3	<b>0.0113</b>
Population (Country)	1798	4, 829	6.1	<b>0.0002</b>
Family (Country*Pop)	317	106, 829	2.6	<b>&lt; 0.0001</b>
Sex	1423	1, 829	11.6	<b>0.0007</b>
Country*Sex	2120	1, 829	17.4	<b>&lt; 0.0001</b>
Error	122	829		

Table 1 continued

<b>Growth rate</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Country	2.2	1, 5	0.2	0.6775
Population (Country)	11.9	4, 829	6.2	<b>0.0002</b>
Family (Country*Pop)	2.1	106, 829	2.0	<b>&lt; 0.0001</b>
Sex	186.6	1, 829	179.9	<b>&lt; 0.0001</b>
Country*Sex	10.5	1, 829	10.2	<b>0.0015</b>
Error	1.0	829		
<b>Adult body mass</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Country	2075.8	1, 5	20.2	<b>0.0098</b>
Population (Country)	111.9	4, 729	3.8	<b>0.0059</b>
Family (Country*Pop)	31.6	106, 729	1.9	<b>&lt; 0.0001</b>
Sex	2669.0	1, 729	162.4	<b>&lt; 0.0001</b>
Error	16.4	729		
<b>Abdomen mass</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Country	245	1, 5	14.1	<b>0.0180</b>
Population (Country)	18.8	4, 729	3.4	<b>0.0119</b>
Family (Country*Pop)	5.9	106, 729	1.6	<b>0.0004</b>
Sex	5494.3	1, 729	1468.7	<b>&lt; 0.0001</b>
Error	3.7	729		
<b>Thorax mass</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Country	237.1	1, 5	3.7	0.1246
Population (Country)	70.4	4, 728	8.9	<b>&lt; 0.0001</b>
Family (Country*Pop)	8.5	106, 728	2.1	<b>&lt; 0.0001</b>
Sex	455.1	1, 728	112.2	<b>&lt; 0.0001</b>
Country*Sex	98.2	1, 728	24.2	<b>&lt; 0.0001</b>
Error	4.1	728		

Table 1 continued

<b>Thorax-abdomen ratio</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Country	0.96	1, 5	2.9	0.1603
Population (Country)	0.36	4, 728	4.1	<b>0.0039</b>
Family (Country*Pop)	0.09	106, 728	1.4	<b>0.0069</b>
Sex	270.33	1, 728	4095.5	<b>&lt; 0.0001</b>
Country*Sex	0.78	1, 728	11.8	<b>0.0006</b>
Error	0.07	728		
<b>Fat (mg)</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Country	2.22	1, 5	5.9	0.0694
Population (Country)	0.44	4, 693	5.0	<b>0.0009</b>
Family (Country*Pop)	0.10	106, 693	3.1	<b>&lt; 0.0001</b>
Sex	2.78	1, 693	85.2	<b>&lt; 0.0001</b>
Error	0.03	693		
<b>Fat (%)</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Country	682.9	1, 5	1.4	0.3006
Population (Country)	575.1	4, 694	5.6	<b>0.0004</b>
Family (Country*Pop)	117.8	106, 694	3.7	<b>&lt; 0.0001</b>
Sex	3754.2	1, 694	118.4	<b>&lt; 0.0001</b>
Error	31.7	694		
<b>Wing length</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Country	49.7	1, 5	45.4	<b>0.0019</b>
Population (Country)	1.2	4, 728	1.7	0.1451
Family (Country*Pop)	0.7	106, 728	2.9	<b>&lt; 0.0001</b>
Sex	1.7	1, 728	6.5	<b>0.0110</b>
Country*Sex	5.5	1, 728	21.3	<b>&lt; 0.0001</b>
Error	0.3	728		

Table 1 continued

<b>Wing area</b>	<b><i>MS</i></b>	<b><i>DF</i></b>	<b><i>F</i></b>	<b><i>P</i></b>
Country	11646	1, 5	68.7	<b>0.0009</b>
Population (Country)	183	4, 728	2.0	0.0925
Family (Country*Pop)	99	106, 728	2.9	<b>&lt; 0.0001</b>
Sex	7958	1, 728	230.1	<b>&lt; 0.0001</b>
Country*Sex	464	1, 728	13.4	<b>0.0003</b>
Error	35	728		
<b>Wing loading</b>	<b><i>MS</i></b>	<b><i>DF</i></b>	<b><i>F</i></b>	<b><i>P</i></b>
Country	14.0	1, 5	0.1	0.7497
Population (Country)	132.4	4, 729	8.4	<b>&lt; 0.0001</b>
Family (Country*Pop)	17.0	106, 729	2.0	<b>&lt; 0.0001</b>
Sex	342.2	1, 729	40.5	<b>&lt; 0.0001</b>
Error	8.4	729		
<b>Wing aspect ratio</b>	<b><i>MS</i></b>	<b><i>DF</i></b>	<b><i>F</i></b>	<b><i>P</i></b>
Country	6.88	1, 5	80.9	<b>0.0001</b>
Population (Country)	0.08	4, 729	0.3	0.8515
Family (Country*Pop)	0.26	106, 729	3.1	<b>&lt; 0.0001</b>
Sex	141.4	1, 729	1676.1	<b>&lt; 0.0001</b>
Error	0.08	729		

**Table 2** Results of a generalized linear model (exploratory behaviour, a) and of a general linear mixed model (flight performance in a vortex, b) for the effects of country, sex (both fixed), population (nested within country; random), and family (nested within population and country, random) in the butterfly *Lycaena tityrus* (models 1). Second models were constructed by additionally including PCs 1-4 and the results of the exploratory behaviour or performance test as covariates. All models were constructed by a stepwise backwards elimination of non-significant factors. The latter models were constructed separately for males and females, owing to strong variation in morphological traits among the sexes. Significant *P*-values are given in bold.

Table 2a

<b>Model 1</b>	<b><i>DF</i></b>	<b><i>Wald Chi<sup>2</sup></i></b>	<b><i>P</i></b>
Country	1	4.7	<b>0.0295</b>
Sex	1	32.5	<b>&lt; 0.0001</b>
<b>Model 2 Males</b>	<b><i>DF</i></b>	<b><i>Wald Chi<sup>2</sup></i></b>	<b><i>P</i></b>
PC2	1	4.4	<b>0.0367</b>
Flight performance	1	10.0	<b>0.0016</b>
<b>Model 2 Females</b>	<b><i>DF</i></b>	<b><i>Wald Chi<sup>2</sup></i></b>	<b><i>P</i></b>
PC2	1	8.8	<b>0.0030</b>

Table 2b

<b>Model 1</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Country	311	1, 5	0.3	0.6282
Population (Country)	1230	4, 723	2.7	<b>0.0353</b>
Family (Country*Pop)	495	106, 723	1.9	<b>&lt; 0.0001</b>
Country*Sex	1322	1, 723	5.2	<b>0.0058</b>
Error	254	723		
<b>Model 2 Males</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Country	1659	1, 5	4.5	0.0623
Population (Country)	447	4, 222	1.2	0.3074
Family (Country*Pop)	431	106, 222	1.8	<b>0.0003</b>
PC1	7139	1, 222	29.5	<b>&lt; 0.0001</b>
PC3	4593.8	1, 222	19.0	<b>&lt; 0.0001</b>
Exploratory behaviour	1768	1, 222	7.3	<b>0.0074</b>
Error	242	222		
<b>Model 2 Females</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Country	923	1, 373	4.2	<b>0.0407</b>
PC1	3519	1, 373	16.1	<b>&lt; 0.0001</b>
PC4	1153	1, 373	5.3	<b>0.0223</b>
Error	219	373		

**A2: Tables 3 and 4 are exclusively based on the animals from Germany, thus focusing on differences across developmental temperature (18°C versus 26°C).**

**Table 3** General linear mixed model results for the effects of temperature, sex (fixed factors), population, and family (nested within population, both random factors) on developmental and morphological traits in *Lycaena tityrus*. Significant *P*-values are given in bold. Models were constructed by a stepwise backwards elimination of non-significant factors.

<b>Larval time</b>	<b><i>MS</i></b>	<b><i>DF</i></b>	<b><i>F</i></b>	<b><i>P</i></b>
Population	12.0	2, 981	1.4	0.2544
Family (Population)	10.4	59, 981	4.2	<b>&lt; 0.0001</b>
Temperature	87024.0	1, 981	35069.0	<b>&lt; 0.0001</b>
Sex	1440.0	1, 981	580.3	<b>&lt; 0.0001</b>
Error	2.5	981		
<b>Pupal time</b>	<b><i>MS</i></b>	<b><i>DF</i></b>	<b><i>F</i></b>	<b><i>P</i></b>
Population	1.9	2, 980	2.7	<b>0.0765</b>
Family (Population)	0.8	59, 980	2.4	<b>&lt; 0.0001</b>
Temperature	15805.2	1, 980	46050.4	<b>&lt; 0.0001</b>
Sex	3.4	1, 980	9.8	<b>0.0018</b>
Temperature*Sex	8.9	1, 980	25.9	<b>&lt; 0.0001</b>
Error	0.3	980		
<b>Pupal mass</b>	<b><i>MS</i></b>	<b><i>DF</i></b>	<b><i>F</i></b>	<b><i>P</i></b>
Population	2598	2, 981	6.4	<b>0.0027</b>
Family (Population)	484	59, 981	3.7	<b>&lt; 0.0001</b>
Sex	5188	1, 981	39.6	<b>&lt; 0.0001</b>
Temperature	15526	1, 981	118.4	<b>&lt; 0.0001</b>
Error	131	981		

Table 3 continued

<b>Growth rate</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Population	14.1	2, 981	10.4	<b>0.0001</b>
Family (Population)	1.6	59, 981	2.4	<b>&lt; 0.0001</b>
Sex	167.6	1, 981	258.5	<b>&lt; 0.0001</b>
Temperature	5577.7	1, 981	8602.9	<b>&lt; 0.0001</b>
Error	0.7	981		
<b>Adult body mass</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Population	97.5	2, 914	2.0	<b>0.1402</b>
Family (Population)	56.3	59, 914	2.9	<b>&lt; 0.0001</b>
Sex	3116.2	1, 914	161.9	<b>&lt; 0.0001</b>
Temperature	129.1	1, 914	6.7	<b>0.0098</b>
Error	19.3	914		
<b>Abdomen mass</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Population	0.11	2, 914	2.1	0.1361
Family (Population)	0.06	59, 914	1.9	<b>&lt; 0.0001</b>
Sex	73.51	1, 914	2359.6	<b>&lt; 0.0001</b>
Temperature	2.84	1, 914	91.2	<b>&lt; 0.0001</b>
Error	0.03	914		
<b>Thorax mass</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Population	101.9	2, 914	7.7	<b>0.0009</b>
Family (Population)	15.5	59, 914	3.2	<b>&lt; 0.0001</b>
Sex	1287.6	1, 914	264.0	<b>&lt; 0.0001</b>
Temperature	96.3	1, 914	19.7	<b>&lt; 0.0001</b>
Error	4.9	914		
<b>Thorax-abdomen ratio</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Population	0.28	2, 972	14.0	<b>&lt; 0.0001</b>
Sex	119.6	1, 972	6023.0	<b>&lt; 0.0001</b>
Temperature	5.2	1, 972	263.7	<b>&lt; 0.0001</b>
Sex*Temperature	0.03	1, 972	1.6	0.2059
Error	0.02	972		

Table 3 continued

<b>Fat (mg)</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Population	23.1	2, 874	18.0	< 0.0001
Family (Population)	1.5	59, 874	3.1	< 0.0001
Sex	46.0	1, 874	93.9	< 0.0001
Error	0.5	874		
<b>Fat (%)</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Population	21.9	2, 874	18.7	< 0.0001
Family (Population)	1.4	59, 874	3.3	< 0.0001
Sex	49.2	1, 874	115.6	< 0.0001
Error	0.4	874		
<b>Wing length</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Population	1.4	2, 915	1.1	0.3416
Family (Population)	1.6	59, 915	5.3	< 0.0001
Sex	12.1	1, 915	40.0	< 0.0001
Error	0.3	915		
<b>Wing area</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Population	252	2, 914	1.5	0.2280
Family (Population)	202	59, 914	5.3	< 0.0001
Sex	6600	1, 914	171.4	< 0.0001
Temperature	1802	1, 914	46.8	< 0.0001
Error	39	914		
<b>Wing loading</b>	<b>MS</b>	<b>DF</b>	<b>F</b>	<b>P</b>
Population	150	2, 914	6.0	0.0039
Family (Population)	29	59, 914	3.2	< 0.0001
Sex	586	1, 914	64.6	< 0.0001
Temperature	910	1, 914	100.3	< 0.0001
Error	9	914		

Table 3 continued

<b>Aspect ratio</b>	<b><i>MS</i></b>	<b><i>DF</i></b>	<b><i>F</i></b>	<b><i>P</i></b>
Population	0.03	2, 914	0	0.9311
Family (Population)	0.48	59, 914	5	<b>&lt; 0.0001</b>
Sex	176.41	1, 914	1877	<b>&lt; 0.0001</b>
Temperature	12.76	1, 914	136	<b>&lt; 0.0001</b>
Error	0.09	914		

**Table 4** Results of a generalized linear model (exploratory behaviour, a) and of a general linear mixed model (flight performance in a vortex, b) for the effects of temperature, sex (both fixed), population (random), and family (nested within population, random) in the butterfly *Lycaena tityrus* (models 1). Second models were constructed by additionally including PCs 1-4 and the results of the exploratory behaviour or performance test as covariates. All models were constructed by a stepwise backwards elimination of non-significant factors. The latter models were constructed separately for males and females, owing to strong variation in morphological traits among the sexes. Significant *P*-values are given in bold.

Table 4a

<b>Model 1</b>	<b><i>DF</i></b>	<b><i>Wald Chi<sup>2</sup></i></b>	<b><i>P</i></b>
Population	2	6.1	<b>0.0465</b>
Temperature	1	43.2	<b>&lt; 0.0001</b>
Sex	1	32.0	<b>&lt; 0.0001</b>
<b>Model 2 Males</b>	<b><i>DF</i></b>	<b><i>Wald Chi<sup>2</sup></i></b>	<b><i>P</i></b>
Temperature	1	24.1	<b>&lt; 0.0001</b>
PC1	1	10.2	<b>0.0014</b>
PC4	1	4.3	<b>0.0391</b>
Flight performance	1	7.4	<b>0.0066</b>
<b>Model 2 Females</b>	<b><i>DF</i></b>	<b><i>Wald Chi<sup>2</sup></i></b>	<b><i>P</i></b>
Temperature	1	13.2	<b>0.0003</b>

Table 4b

<b>Model 1</b>	<b><i>MS</i></b>	<b><i>DF</i></b>	<b><i>F</i></b>	<b><i>P</i></b>
Population	1005	2, 909	2.6	0.0781
Family (Population)	426	59, 909	1.9	< <b>0.0001</b>
Sex*Temperature	2951	1, 909	13.3	< <b>0.0001</b>
Sex	18770	1, 909	84.5	< <b>0.0001</b>
Error	222	909		
<b>Model 2 Males</b>	<b><i>MS</i></b>	<b><i>DF</i></b>	<b><i>F</i></b>	<b><i>P</i></b>
Population	1222	2, 410	6.5	<b>0.0017</b>
Temperature	4766	1, 410	25.1	< <b>0.0001</b>
PC1	1941	1, 410	10.2	<b>0.0015</b>
PC3	7442	1, 410	39.3	< <b>0.0001</b>
Flower test	1585	1, 410	8.4	<b>0.0040</b>
Error	190	410		
<b>Model 2 Females</b>	<b><i>MS</i></b>	<b><i>DF</i></b>	<b><i>F</i></b>	<b><i>P</i></b>
Population	408	2, 373	1.0	0.3773
Family (Population)	473	59, 373	2.2	< <b>0.0001</b>
PC1	2914	1, 373	13.5	<b>0.0003</b>
PC3	4356	1, 373	20.1	< <b>0.0001</b>
PC4	942	1, 373	4.4	<b>0.0376</b>
Error	216	373		

**A3: Table 5 shows trait variation across replicate populations.**

**Table 5** Developmental, morphological, and physiological traits in relation to population (country) in *Lycaena tityrus*. Given are means  $\pm$  1 SE. Group sample sizes range between 60 and 399 individuals. Different superscript letters within a row indicate significant differences among groups (Tukey HSD).

Trait	Germany			Estonia		
	Guest	Ueckermünde	Waren	Mahu	Kasispea	Karepa
Larval time (days)	23.9 ± 0.56 <sup>a</sup>	24.6 ± 0.53 <sup>a</sup>	24.3 ± 0.46 <sup>a</sup>	15.2 ± 0.37 <sup>b</sup>	15.4 ± 0.42 <sup>b</sup>	15.9 ± 0.47 <sup>b</sup>
Pupal time (days)	10.9 ± 0.22 <sup>a</sup>	11.2 ± 0.22 <sup>a</sup>	11.1 ± 0.20 <sup>a</sup>	7.5 ± 0.15 <sup>b</sup>	7.4 ± 0.18 <sup>b</sup>	7.5 ± 0.20 <sup>b</sup>
Pupal mass (mg)	119.7 ± 0.77 <sup>ab</sup>	122.4 ± 0.81 <sup>a</sup>	117.5 ± 0.59 <sup>bd</sup>	109.5 ± 0.81 <sup>c</sup>	114.2 ± 1.25 <sup>cd</sup>	109.7 ± 1.01 <sup>c</sup>
Growth rate (mg/day)	5.9 ± 0.16 <sup>a</sup>	6.0 ± 0.15 <sup>a</sup>	5.7 ± 0.11 <sup>a</sup>	7.7 ± 0.13 <sup>b</sup>	7.7 ± 0.15 <sup>b</sup>	7.3 ± 0.14 <sup>b</sup>
Adult body mass (mg)	37.4 ± 0.33 <sup>a</sup>	37.1 ± 0.29 <sup>ab</sup>	36.1 ± 0.24 <sup>bc</sup>	33.0 ± 0.34 <sup>d</sup>	36.1 ± 0.56 <sup>ac</sup>	33.0 ± 0.46 <sup>d</sup>
Abdomen mass (mg)	11.5 ± 0.27 <sup>a</sup>	10.7 ± 0.21 <sup>ab</sup>	10.4 ± 0.18 <sup>bce</sup>	9.1 ± 0.24 <sup>d</sup>	11.6 ± 0.41 <sup>ac</sup>	9.4 ± 0.30 <sup>de</sup>
Thorax mass (mg)	17.7 ± 0.15 <sup>a</sup>	18.5 ± 0.16 <sup>b</sup>	17.5 ± 0.13 <sup>ac</sup>	16.9 ± 0.16 <sup>c</sup>	17.4 ± 0.22 <sup>ac</sup>	16.8 ± 0.20 <sup>c</sup>
Thorax-abdomen ratio	1.8 ± 0.04 <sup>ac</sup>	1.9 ± 0.04 <sup>b</sup>	1.9 ± 0.03 <sup>abc</sup>	2.0 ± 0.05 <sup>b</sup>	1.6 ± 0.07 <sup>c</sup>	2.0 ± 0.06 <sup>ab</sup>
Fat (mg)	0.5 ± 0.03 <sup>a</sup>	0.3 ± 0.01 <sup>b</sup>	0.3 ± 0.01 <sup>c</sup>	0.3 ± 0.02 <sup>b</sup>	0.2 ± 0.02 <sup>bc</sup>	0.2 ± 0.01 <sup>b</sup>
Fat (%)	14.0 ± 0.76 <sup>a</sup>	7.4 ± 0.30 <sup>bd</sup>	10.0 ± 0.35 <sup>c</sup>	9.2 ± 0.52 <sup>bc</sup>	5.3 ± 0.46 <sup>d</sup>	7.7 ± 0.46 <sup>bd</sup>
Wing length (mm)	15.2 ± 0.03 <sup>a</sup>	15.4 ± 0.04 <sup>b</sup>	15.3 ± 0.03 <sup>ab</sup>	14.9 ± 0.04 <sup>c</sup>	14.9 ± 0.06 <sup>c</sup>	14.8 ± 0.05 <sup>c</sup>
Wing area (mm <sup>2</sup> )	91.7 ± 0.46 <sup>ad</sup>	93.7 ± 0.46 <sup>b</sup>	92.8 ± 0.36 <sup>ab</sup>	87.1 ± 0.52 <sup>cd</sup>	89.7 ± 0.79 <sup>d</sup>	85.9 ± 0.66 <sup>c</sup>
Wing loading (mg/cm <sup>2</sup> )	40.7 ± 0.23 <sup>a</sup>	39.4 ± 0.20 <sup>b</sup>	38.9 ± 0.17 <sup>bd</sup>	37.8 ± 0.24 <sup>c</sup>	40.2 ± 0.37 <sup>ab</sup>	38.3 ± 0.31 <sup>cd</sup>
Wing aspect ratio	10.1 ± 0.04 <sup>ab</sup>	10.1 ± 0.03 <sup>ab</sup>	10.1 ± 0.03 <sup>ab</sup>	10.2 ± 0.04 <sup>a</sup>	9.9 ± 0.05 <sup>b</sup>	10.2 ± 0.05 <sup>a</sup>



# **EIGENSTÄNDIGKEITSERKLÄRUNG**



## **Eigenständigkeitserklärung**

Hiermit erkläre ich, dass diese Arbeit bisher von mir weder an der Mathematisch-Naturwissenschaftlichen Fakultät der Ernst-Moritz-Arndt-Universität Greifswald noch einer anderen wissenschaftlichen Einrichtung zum Zwecke der Promotion eingereicht wurde.

Ferner erkläre ich, dass ich diese Arbeit selbständig verfasst und keine anderen als die darin angegebenen Hilfsmittel und Hilfen benutzt und keine Textabschnitte eines Dritten ohne Kennzeichnung übernommen habe.

Unterschrift des Promovenden