

RESEARCH ARTICLE

A low-threshold intervention to increase physical activity and reduce physical inactivity in a group of healthy elderly people in Germany: Results of the randomized controlled MOVING study

Fabian Kleinke^{1,2*}, Sabina Ulbricht^{2,3}, Marcus Dörr^{2,4}, Peter Penndorf^{1,2}, Wolfgang Hoffmann^{1,2}, Neeltje van den Berg^{1,2}

1 Institute for Community Medicine, Section Epidemiology of Health Care and Community Health, University Medicine Greifswald, Greifswald, Germany, **2** DZHK (German Centre for Cardiovascular Research), Partner Site Greifswald, Greifswald, Germany, **3** Institute for Community Medicine, Section Prevention Research and Social Medicine, University Medicine Greifswald, Greifswald, Germany, **4** Department of Internal Medicine B, University Medicine Greifswald, Greifswald, Germany

* fabian.kleinke@uni-greifswald.de



OPEN ACCESS

Citation: Kleinke F, Ulbricht S, Dörr M, Penndorf P, Hoffmann W, van den Berg N (2021) A low-threshold intervention to increase physical activity and reduce physical inactivity in a group of healthy elderly people in Germany: Results of the randomized controlled MOVING study. *PLoS ONE* 16(9): e0257326. <https://doi.org/10.1371/journal.pone.0257326>

Editor: Thalia Fernandez, Universidad Nacional Autonoma de Mexico, MEXICO

Received: February 10, 2021

Accepted: August 23, 2021

Published: September 16, 2021

Peer Review History: PLOS recognizes the benefits of transparency in the peer review process; therefore, we enable the publication of all of the content of peer review and author responses alongside final, published articles. The editorial history of this article is available here: <https://doi.org/10.1371/journal.pone.0257326>

Copyright: © 2021 Kleinke et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Background

Lack of physical activity (PA) and a high level of physical inactivity (PI) are associated with a higher risk for mortality and responsible for several non-communicable diseases including cardiovascular disease. Higher age is associated with a decrease of PA and an increasing level of PI. Studies have shown that interventions in the elderly have the potential to increase the amount of PA and to decrease the level of PI. However, most interventions are complex, elaborated, time- and resource-consuming. Here, we examined the effect of individual feedback-letters reporting the measured PA and PI in a sample of elderly people in Germany. Primary outcomes of the study were overall PA and PI after 6 months in the intervention group compared to a control group.

Methods

We examined data from the MOVING intervention study (RCT) for people aged ≥ 65 years living in the northeast of Germany. At baseline, 3 and 6-months follow-up, all study participants wore a 3-axis accelerometer over a period of seven consecutive days. After the baseline measurement, the participants were randomized into intervention and control group. Participants in the intervention group received automatically generated, individualized feedback letters reporting their PA and PI by mail after the baseline measurement and after the 3-months follow-up. A Two-Way Mixed ANOVA with repeated measures was calculated with light, moderate and overall PA as well as PI as dependent variables, and group (between subject) and time (inner subject) as factors. The analysis based on retrospective data from the MOVING study (2016–2018).

Data Availability Statement: All relevant data are within the manuscript and its [Supporting Information](#) files.

Funding: The study was funded by the Federal Ministry of Education and Research (BMBF) as a site project of the German Centre for Cardiovascular Research (DZHK) (funding sign: 81Z7400174). The funders have had no influence on the conceptualization and conduct of the study and will not have any role in the data analysis and publication of the results.

Competing interests: The authors have declared that no competing interests exist.

Results

N = 258 patients were recruited. N = 166 participants could be included in the analysis, thereof N = 97 women (58.4%). The mean age was 70.8 years (SD 4.8). At baseline, the participants had a mean wearing time of 5,934.5 minutes (SD = 789.5) per week, which corresponds to about 14 hours daily on average. The overall PA in the intervention group at the 6-months follow up was 2488.8 (95% CI 2358.9–2618.2) minutes and 2408.2 (95% CI 2263.0–2553.4) minutes in the control group. There was no statistically significant interaction effect (time*group) between the intervention and control group for the depending variables. Sensitivity analyses showed significant small positive effects of the interaction time*partnership, $F(2, 300) = 3.020$, $p = 0.05$, partial $\eta^2 = 0.020$.

Discussion

On average, study participants had high levels of PA at baseline and showed a good adherence in wearing the accelerometer. Both is likely due to selection in the convenience study sample. Thus, some ceiling effect reduced the overall intervention effect somewhat. At baseline, the weekly average of PI was 3436.7 minutes, which corresponds to about 8.2 hours per day and about 57% of participants' daily waking time. The average level of PI could be slightly decreased in both study groups.

Trial registration number

[DRKS00010410](#), 17 May 2017.

Introduction

There is strong evidence, that physical activity (PA) promotes healthy aging and that both lack of PA and a high level of physical inactivity (PI) are crucial risk factors for global mortality and several non-communicable diseases (NCD) [1–7].

PI is defined as “any waking behavior characterized by an energy expenditure ≤ 1.5 METs (metabolic equivalent of task) while in a sitting or reclining posture” [8, 9]. 31.1% of the worldwide adult population has a sedentary lifestyle, is insufficiently physically active and does not meet the recommendations for PA from the World Health Organization (WHO) [10]. Physical inactivity causes 3.2 million deaths per year worldwide. In 2010 PI was responsible for 69.3 million DALYs (disability-adjusted life years) [11]. PI causes a significant economic burden, in 2013, PI caused 53.8 billion US-Dollar healthcare costs worldwide [12]. As a consequence, the reduction of insufficient PA is a global target of the WHO [11, 13].

Depending on the definition, the proportion of PI across European countries ranges from 43.3% in Sweden up to 87.8% in Portugal. In Germany, 70.2% of men and 71.8% of women show a sedentary lifestyle [14]. High levels of moderate to vigorous physical activity (MVPA) are associated with a reduced risk for all-cause mortality [6, 15, 16]. In general, epidemiological evidence about positive effects of regular PA is strong and has accumulated over several decades [4, 17–19].

Several studies have shown that a physically active lifestyle (including promoting PA and minimizing PI) is associated with improved overall health status [4, 20] and a reduced risk for overall mortality by 22–34%, and CVD mortality by 27–35% [1]. In addition, regular PA is a

well-established method to prevent diabetes, hypertension and several types of cancer (i.e. breast and colon cancer) also for older people. Furthermore, regular PA reduces the risk of developing stroke about 25–30% [21]. Even people who are only 1–2 times per week physically active, have a reduced risk for mortality [22] compared to those with less PA.

There is strong evidence, that moderate-to-vigorous PA (MVPA) has positive effects on important health outcomes and is associated with a reduced risk of all-cause mortality [16]. Several international physical activity guidelines from the United States [23], United Kingdom [24], Australia [25], and from the WHO [26] promote the benefits of both moderate and vigorous PA. Recommendations for light PA are lacking in the guidelines [24–27]. However, increasing light PA also seems to be positively associated with relevant health outcomes, i.e. obesity and glucose metabolism, and is associated with a reduced risk for all-cause mortality [19, 28, 29]. In addition, light PA has beneficial effects on the health situation of elderly people, e.g. a higher cognitive performance [30], and it is positively associated with physical health and life satisfaction [31]. Hence, the impact of light PA on public health may be underestimated to some extent.

Increasing age is associated with decreasing regular PA, a decreasing number of daily steps [32] and an increase of PI which contributes to higher risks for chronic diseases (e.g. stroke) and multimorbidity [10, 33–36]. In the context of demographic change, particularly older people should be an important target group for extensive and practicable prevention strategies.

Interventions in the elderly have the potential to increase PA [37–39] and to reduce PI [40–42]. Several pedometer-based intervention studies have shown positive effects on the level of PA. Harris et al. showed a positive effect of an intervention consisting of four consultations by a primary care nurse on daily steps and weekly MVPA in ≥ 10 min bouts.

A meta-analysis on pedometer-based PA interventions pointed out that usage of a pedometer has a moderate positive effect on PA levels (average increase of 2,000 daily steps) [38]. These results correspondent to a systematic literature review in which participants in the intervention group increased their PA by 2,491 daily steps by using a pedometer-based intervention compared to the control group [39].

Interventions also have the potential to decrease PI [43]. Mutrie et al. have shown that a nurse-delivered intervention in the elderly (N = 41), based on social cognitive theory, reduced daily PI by 68 minutes in the intervention group (measured by pedometer and accelerometer) compared to the controls [44]. Another study of older adults (N = 478) showed that a home-based intervention based on social cognitive theory led to a decrease of 57 minutes of sitting time per day in the intervention group compared to the control group [42].

Most of the successful studies are based on complex interventions (i.e. personal consultations) [37]. These are associated with high costs and a large demand of personal resources. Dissemination of such interventions to the general older population may be hardly feasible, especially against the background of the demographic change. In conclusion, a translation of such interventions into daily life of a major proportion of the elderly has been difficult. Therefore, we designed a low-threshold and practicable intervention to increase overall PA and decrease PI.

Research objectives

Aim of the study was the evaluation of a low-threshold intervention to improve overall PA and simultaneously reduce PI. The objective of this analysis was to examine the effect of a low-threshold intervention, consisting of automatically generated, individualized feedback-letters reporting the measured physical activity on PA and PI in a sample of apparently healthy elderly people in the Northeast of Germany (MOVING-study (Motivation-Oriented intervention

study for the elderly in Greifswald). Primary outcomes of the analysis were overall PA and PI after 6 months in the intervention group compared to the control group.

Materials and methods

Study sample

A detailed study protocol [45], as well as levels and determinants for PA and PI of the study sample of the MOVING study at baseline are published elsewhere [46]. In brief, the study population consists of people sampled from the general population who meet the following inclusion criteria:

- Age \geq 65 years
- The possibility of being physically active in daily life

Exclusion criteria were:

- Permanent use of a wheelchair (no ability to walk independently)
- Simultaneous participation in other studies addressing PA and PI
- Not accessible by telephone or cell phone
- Fulfillment of the WHO recommendations for PA (self-report) for people aged \geq 65 at baseline (\geq 150 minutes of moderate PA or 75 minutes of vigorous PA)

All eligible patients were informed in detail about the study and had to give their written informed consent prior to study inclusion. The project data was assessed using eCRFs in an IT-supported documentation system [47]. At Baseline, all participants received a study ID-card with individual 1D-barcode for reliable re-identification.

Study procedure

Potential participants and interested persons contacted the study staff by telephone. A short telephone interview was performed to check the inclusion and exclusion criteria. All persons who fulfilled the inclusion criteria were invited to the examination center of the German Centre for Cardiovascular Research (DZHK) of the University Medicine Greifswald. Here, all participants received a 3-axis accelerometer (ActiGraph GT3x-BT, Pensacola, FL, USA) and specific instructions on how to wear the accelerometer device.

After the baseline examination, participants were individually randomized 1:1 into the control and the intervention group. Follow-up examinations were conducted in both groups at 3 and 6 months after the baseline examination with the same program as in the baseline examination.

Measures

All study participants received a physical examination consisting of blood pressure measurement and measurement of body weight, waist and hip circumference. Trained and certified study staff carried out all measurements according to standard operating procedures (SOP). At baseline, participants were asked for sociodemographic data (sex, age, education (number of years participants went to school)). In addition, participants received standardized paper-pencil questionnaires on PA and PI (IPAQ [48], German PAQ 50+ [49], MOST [50], SF8 [51] and SSA-scale [52]) that had to be completed after the 7-days of wearing the accelerometer device.

After the baseline and at the two follow-up examinations, all study participants wore the accelerometer device at daytime on the right hip. The participants were instructed to remove the device only for water-based activities. The three-axis accelerometer recorded PA, PI and sedentary breaks (a period of non-sedentary activity (e.g. walking) in between two sedentary conditions (e.g. sitting) continuously at a sampling frequency of 30 Hz over a period of seven consecutive days, starting at midnight after the examination day.

The software ActiLife 6.13.2–6.13.3 (ActiGraph Cop. ©, Pensacola, FL, USA) was used for data download and data processing. Raw data were calculated into 60-second epochs and saved as GT3X files.

We used specific cut points based on Freedson [53] to categorize the intensity of participants' PA. For adults, this cut-point model is considered as established for the determination of different physical activity categories [54]. Intensity of PA is categorized as sedentary (0–99 counts), light (100–1951 counts), moderate (1952–5724 counts), and vigorous (5725–9498 counts) PA. Non-wear time was defined as ≥ 60 min of continual zero counts.

Intervention

The intervention based on individual accelerometer data and consisted of automatically generated, personalized feedback letters sent by mail. The participants in the intervention group received the letters after the baseline examination and the 3-month follow-up shortly after wearing the accelerometer. The participants in the control group received no feedback during the study period. According to the Declaration of Helsinki, we provided participants in the control group a detailed feedback about PA and PI after the end of the study within a thank you letter. This procedure ensured that all participants (both groups) received in sum the same information at the end of the study.

The feedback letters contained three graphs, reporting on the measured PA and PI, with provision of the number of steps per day, daily and weekly moderate and vigorous PA in minutes and daily and weekly sedentary time in minutes, as well as a paragraph about the importance of sedentary breaks. The participants received age appropriate recommendations for PA based on the WHO-guidelines [26]. The feedback letters were automatically generated in R software (version 3.3.2, Lucent Technologies, Murray Hill, NJ, USA).

Data analysis

To examine the effects of the intervention, we analyzed overall, light and moderate PA as well as PI as dependent variables with a Two-Way Mixed ANOVA with repeated measures with group (between subject) and time (inner subject) as factors. For baseline data, we analyzed the equality of variables between both study groups. Due to significant difference between both groups in partnership ($p = 0.006$), we included partnership as a covariable in the Two-Way Mixed ANOVA. First, we analyzed the assumptions and requirements of mixed ANOVA i.e. distribution and homoscedasticity and detected no violation of the assumptions". Bonferroni adjustment was made because several analyses were performed.

All statistical analyses were performed using pseudonymized data. In addition, we performed a sub-group analysis and excluded participants with ≥ 150 minutes of weekly moderate PA or ≥ 75 minutes of weekly vigorous PA at baseline (according to the WHO recommendation).

A valid measurement day was defined as a record of at least 10 hours of total wearing time during the day. Additionally, a record time of at least 4 valid days was required for all data analyses.

Statistical significance was assumed for p-values <0.05. Data processing and statistical calculations were performed with IBM SPSS Statistics 26 or later (1989–2020 by IBM Corp. ©, Armonk, New York, USA.).

Ethical consideration

In 2016, the clinical ethics committee of the University Medicine Greifswald (protocol number BB071/16) approved the study.

Results

A number of 258 potential participants were contacted personally or by telephone, thereof 206 were included in the study and randomized (Fig 1).

At baseline, the participants had a mean wearing time of the accelerometer of 5,934.5 minutes (SD = 789.5) per week, which corresponds to 14.1 hours daily. N = 166 participants could be included in the analysis, thereof n = 97 women (58.4%) and n = 69 men (41.6%). At baseline, the mean age was 70.8 years (SD = 4.8) (Table 1).

At baseline, the average time of overall PA was 2473.7 (95% CI 2387.0–2560.4) minutes per week, to which light PA contributed the largest part with 2260.9 (95% CI 2181.1–2340.7) minutes. In addition, the average of weekly PI (baseline) was 3460.8 (95%-CI 3337.5–3584.0) minutes per week which corresponds to about 8.2 hours per day and about 57% of participants' waking time.

Participants in the intervention group slightly increased their weekly time of light PA from baseline to 6-months follow-up on average by + 54.0 minutes (control group—59.2 minutes). The average overall PA increased in the intervention group by 25.0 minutes per week (control group—75.9 minutes) between baseline and 6 months follow-up. At 6-months follow-up (primary endpoint), the weekly difference of overall PA between intervention group (2488.8 minutes, 95% CI 2358.9–2618.2) and control group (2408.2 minutes, 95% CI 2263.0–2553.4) was 80.6 minutes which corresponds to about 1.3 hours of additional overall PA per week in the intervention group (Table 2).

At baseline, both study groups started almost on the same value of overall PA. At 6-months follow-up, the amount of overall PA decreased in the control group (red line) the baseline value while participants in the intervention group (blue line) could slightly increase their weekly overall PA (Fig 2).

At baseline, n = 97 (58.4%) of the study participants (both study groups) achieved the international recommendations for moderate PA (≥ 150 minutes moderate or ≥ 75 minutes vigorous PA). The proportion of participants who fulfilled the recommendations slightly decreased over study time (Table 3).

There was no statistically significant interaction between time and group (interaction effect: time*group) for the depending variables PI, light, moderate and overall PA (sphericity assumed). P-values were ranging between 0.317 to 0.546. *Physical inactivity* (n = 146)

The results of the Two-Way Mixed ANOVA showed that there was a significant main effect of group, $F(1, 143) = 5.335$, $p = 0.022$, partial $\eta^2 = .036$. In addition, there was no significant main effect of time, sphericity assumed $F(2, 286) = 0.763$, $p = 0.467$, partial $\eta^2 = .005$.

Descriptive statistics showed that both study groups decreased the amount of PI over study time. At 6-months follow-up, participants included in the analyses had an amount of 3310.4 minutes (SD = 536.1) of PI (mean) in the intervention group, and 3161.1 minutes (SD = 581.8) of PI (mean) in the controls. Besides that, the differences between both groups were about the same amount of minutes in PI at each measurement point (baseline, 3- and 6-months follow-up).




Fig 1. Consort diagram: Recruited participants and participants included in the analysis (N = 166).

<https://doi.org/10.1371/journal.pone.0257326.g001>

Light PA (n = 155)

The results of the Two-Way Mixed ANOVA showed that there was no significant main effect of group, $F(1, 153) = 0.017$, $p = 0.895$, partial $\eta^2 = .000$, and time, sphericity assumed $F(2, 304) = 2.196$, $p = 0.113$, partial $\eta^2 = .014$.

Descriptive statistics showed almost the same baseline value for light PA between the intervention (2235.8 minutes (SD = 495.7) and control group (2232.1 minutes, SD = 536.4). After an increase of light PA in both study groups to 3-months follow-up, participants in the control group decreased minutes of light PA even lower than the baseline value (2187.4 minutes, SD = 604.5) at 6-months follow-up. Besides that, participants in the intervention group remained almost at the same amount of light PA at 6-months follow-up (2298.8 minutes, SD = 557.8). Thus, participants in the intervention group showed a higher amount of light PA

Table 1. Descriptive characteristics of the study sample (N = 166), complete and valid data sets (baseline, 3 and 6 months follow-up).

Characteristics	Control group		Intervention group		Both groups	
	n	Mean (SD) or n (%)	n	Mean (SD) or n (%)	n	Mean (SD) or n (%)
Sex (women)	81	47 (58.0%)	85	50 (58.8%)	166	97 (58.4%)
Age (yr)	81	71.2 (SD 5.0)	85	70.4 (SD 4.6)	166	70.8 (SD 4.8)
Number of participants currently living in a partnership (yes)*	77	47 (61.0%)	83	67 (80.7%)	160	114 (71.3%)
Education (yr)	74		82		156	
< 10 years		14 (18.9%)		18 (22.0%)		32 (20.5%)
= 10 years		25 (33.8%)		17 (20.7%)		42 (26.9%)
> 10 years		30 (40.5%)		45 (54.9%)		75 (48.1%)
Other		5 (6.8%)		2 (2.4%)		7 (4.5%)
Body mass index (kg/m ²)	81		85		166	
< 25		22 (27.2%)		23 (27.1%)		45 (27.1%)
≥ 25 and < 30		33 (40.7%)		30 (35.3%)		63 (38.0%)
≥ 30		26 (32.1%)		32 (37.6%)		58 (34.9%)
Waist circumference (cm)	81	96.7 (SD 13.1)	85	94.8 (SD 14.6)	166	95.8 (SD 13.9)
Hip circumference (cm)	81	104.1 (SD 11.8)	85	102.7 (SD 10.4)	166	103.4 (SD 11.1)
Blood pressure (sys/dia mmHg) (Pulse /min)	81	133/74 (70)	84	132/72 (73)	165	133/73 (71)
Wearing time of the accelerometer						
Baseline	81	5,922.7 (SD 926.8)	85	5,945.7 (SD 637.2)	166	5,934.5 (SD 789.5)
3-month follow-up	81	5,833.6 (SD 839.1)	84	5,826.4 (SD 683.9)	165	5,829.9 (SD 761.7)
6-month follow-up	79	5,636.4 (SD 719.7)	82	5,840.7 (SD 719.0)	161	5,740.5 (SD 724.4)

Notes: n number of subjects, SD standard deviation * significant difference between intervention and control group (p = 0.006).

<https://doi.org/10.1371/journal.pone.0257326.t001>

at 6-months follow-up (primary endpoint) compared to the controls (+111.4 minutes of light PA) in the descriptive statistics.

Moderate PA (n = 147)

The results of the Two-Way Mixed ANOVA showed that there was no significant main effect of group, $F(1, 144) = 1.426$, $p = 0.234$, partial $\eta^2 = .010$, and time, sphericity assumed $F(2, 288) = 0.715$, $p = 0.490$, partial $\eta^2 = .005$.

Descriptive statistics showed a constant decline of moderate PA in the control intervention group starting from baseline (172.4 minutes, SD = 124.1) to 6-month follow-up (145.9 minutes, SD = 115.1). Participants in the intervention group started with an average amount of moderate PA about 201.8 minutes (SD = 149.6) and could increase the amount after a decline at 3-months follow-up (163.9 minutes, SD = 127.4) at 6-months follow-up to 170.9 minutes (SD = 126.3).

Overall PA (n = 155)

The results of the Two-Way Mixed ANOVA showed that there was no significant main effect of group, $F(1, 152) = 0.005$, $p = 0.945$, partial $\eta^2 = .000$, and time, sphericity assumed $F(2, 304) = 1.638$, $p = 0.196$, partial $\eta^2 = .011$.

Descriptive statistics showed slightly increase (+36.1 minutes) of the amount of overall PA in the intervention group starting at 2459.6 minutes (SD = 558) to 3-months follow-up (2495.7, SD = 564.4). Participants in the intervention group stayed almost on this level at 6-months follow-up (2494.6, SD = 590.8). The controls started almost on the same baseline value (2459.0, SD = 581.1). After an increase to 3-month follow-up (2504.7, SD = 657.2), the

Table 2. Physical activity and inactivity (mean number of minutes during the wearing-week), by intensity of physical activity and overall PA (N = 166).

Study phase	Study Group	Physical inactivity	Sedentary breaks	Light PA	Moderate PA	Vigorous PA	Overall PA	Steps Counts
		Mean (CI 95%)	Mean (CI 95%)	Mean (CI 95%)	Mean (CI 95%)	Mean (CI 95%)	Mean (CI 95%)	Mean (CI 95%)
Baseline	Intervention group (n = 85)	3482.0 (3336.1–3627.8)	97.6 (92.7–102.4)	2246.4 (2139.9–2352.8)	213.6 (180.6–246.6)	3.8 (0.6–7.0)	2463.8 (2342.8–2584.8)	99,278.0 (92,889.8–105,666.2)
	Control group (n = 81)	3438.5 (3234.3–3642.7)	95.9 (89.3–101.9)	2276.1 (2154.4–2397.8)	201.1 (166.5–235.6)	6.6 (1.2–14.3)	2484.1 (2357.2–2611.1)	97,763.2 (90,840.4–104,685.9)
	Total (n = 166)	3460.8 (3337.5–3584.0)	96.8 (93.0–100.6)	2260.9 (2181.1–2340.7)	207.5 (183.8–231.1)	5.1 (1.1–9.2)	2473.7 (2387.0–2560.4)	98,538.8 (93,884.7–103,193.0)
3-months follow-up	Intervention group (n = 84)	3351.1 (3204.8–3497.3)	92.4 (87.6–97.0)	2292.3 (2174.7–2409.8)	180.3 (148.3–212.3)	2.7 (0.2–5.6)	2475.3 (2348.7–2602.0)	99,554.3 (93,268.3–105,840.2)
	Control group (n = 81)	3318.8 (3140.0–3497.7)	93.6 (87.8–99.4)	2327.5 (2188.9–2466.2)	181.5 (151.0–212.0)	5.3 (1.0–9.7)	2514.8 (2369.9–2659.6)	97,567.4 (90,803.8–104,330.9)
	Total (n = 165)	3335.3 (3221.3–3449.2)	93.0 (89.4–96.6)	2309.6 (2219.9–)	180.9 (159.0–202.8)	4.0 (1.4–6.6)	2494.7 (2399.7–2589.7)	98,578.9 (94,013.7–103,144.1)
6-months follow-up	Intervention group (n = 83)	3354.8 (3210.0–3499.6)	92.5 (88.1–96.9)	2300.4 (2180.1–2420.8)	184.9 (154.2–215.6)	3.3 (0.7–5.8)	2488.8 (2358.9–2618.2)	99,613.6 (93,526.0–105,701.2)
	Control group (n = 79)	3228.2 (3087.0–3369.4)	90.2 (85.6–94.7)	2216.9 (2077.8–2356.0)	187.7 (150.3–225.1)	3.5 (1.6–8.5)	2408.2 (2263.0–2553.4)	94,301.4 (87,290.4–101,312.5)
	Total (n = 162)	3293.1 (3192.5–3393.7)	91.4 (88.2–94.5)	2259.7 (2168.8–2350.6)	186.3 (162.5–210.1)	3.4 (0.6–6.2)	2449.5 (2353.2–2545.8)	97,023.1 (92,426.6–101,619.7)

Notes: *n* number of subjects, *CI* 95% confidence interval.

<https://doi.org/10.1371/journal.pone.0257326.t002>

amount of overall PA declined to 6-months follow-up below the intervention group (2399.4, SD = 661.4). At 6-months follow-up, the group difference was 95.2 minutes of weekly overall PA.

Sensitivity analyses

In a sensitivity analysis, we excluded participants with ≥ 150 minutes of weekly moderate PA or with ≥ 75 minutes of vigorous PA at baseline (according to the WHO recommendation). Here ($n = 153$), we found a statistically significant interaction effect for time*partner (covariable) for light PA, sphericity assumed $F(2, 300) = 3.020$, $p = 0.05$, partial $\eta^2 = 0.020$. This result indicate a small positive effect on light PA over study time in participants who are in a partnership.

Discussion

The results of this analysis show that the levels of light, moderate and overall PA in our sample were exceptionally high for this age group [55–57]. Both groups started on almost the same values of overall PA at baseline (intervention group = 2463.8 (95% CI 2342.8–2584.8) minutes,




Fig 2. Mean minutes of overall PA (95% CI, intervention group and control group) at baseline, 3 and 6 months follow-up (N = 166), control group (red line), intervention group (blue line).

<https://doi.org/10.1371/journal.pone.0257326.g002>

control group = 2484.1 (95% CI 2357.2–2611.1) minutes. The level of overall PA over both groups remained almost on the same level at the 3-months follow-up (intervention group = 2475.3 (95% CI 2348.7–2602.0) minutes, control group = 2514.8 (95% CI 2369.9–2659.6) minutes), and 6-months follow-up (intervention group = 2488.8 (95% CI 2358.9–2618.2) minutes, control group = 2408.2 (95% CI 2263.0–2553.4) minutes). At 6-months follow-up, participants in the intervention group showed more overall PA compared to the participants in the control group (difference = 80.6 minutes per week which corresponds to about 1.3 hours of additional overall PA per week in the intervention group).

In previous randomized controlled intervention trials promoting PA increasing PA levels have often been observed also in the control groups [58, 59]. Therefore, several possible

Table 3. Number and proportion of study participants who fulfilled the WHO recommendations PA for people aged over 65 years, separately for intervention and control group (N = 166).

WHO recommendation	Study group	Baseline (n = 166, IG = 85, CG = 81)	3-month follow-up (n = 165, IG = 84, CG = 81)	6-month follow-up (n = 162, IG = 83, CG = 79)
moderate PA ≥150 min. per week or	Intervention group	49 (57.6%)	43 (51.2%)	41 (49.4%)
	Control group	48 (59.3%)	39 (48.1%)	38 (48.1%)
vigorous PA ≥75 min. per week	Total	97 (58.4%)	82 (49.7%)	79 (48.8%)

n number of subjects, fulfillment of WHO recommendations for moderate PA for people aged 65 and older: ≥150 minutes moderate PA or ≥75 minutes vigorous PA or an equivalent combination per week (2 minutes moderate PA correspond to 1 minute vigorous PA).

<https://doi.org/10.1371/journal.pone.0257326.t003>

reasons have been proposed, e.g. the “Hawthorne effect”, which may have changed participants’ behavior as a result of participating in the study—irrespective of their randomisation. In our study, we observed this effect according to the amount of PI, both study groups (including controls) decreased PI over study time.

However, in a longitudinal analysis at the 6-month follow-up, the levels of light and overall PA had decreased in the control group (below the baseline value), while PA in the intervention group remained almost at the same level. Seasonal influences, (weather) cannot be the cause for the difference, because the intervention and the control group were simultaneously recruited and included between November 2016 and December 2018.

Also the amount of PI was high and with 3460.8 (95% CI 3337.5–3584.0) minutes per week at baseline, which corresponds to 8.2 hours/day and is comparable with other studies examining the same age group [14, 60]. Thus, participants in both study groups spent on average about 57% of their waking time in PI at baseline. The high amount of PI combined with a parallel high amount of PA indicates that PI and PA can be seen as independent factors.

The intervention showed no significant effect regarding the primary outcomes. However, a large part of the participants was already physically active on a high level: 58.4% (n = 97) of the participants already met the recommendations for people aged over 65 years of the WHO at baseline. This limits the potential to increase the participant’s PA. Sensitivity analyses excluding very active participants, showed significant effects of the intervention on overall PA among the subgroup of the less active. Thus, further research should focus more on people with low PA levels. However, as a consequence of some self-selection of the more physically active elderly, recruitment of people with a low level of PA may be challenging.

The layout of interventions to increase PA and reduce PI is crucial. As previous research has shown, interventions should focus on the target group and especially on individual factors (e.g. age, sex) [46]. In addition, an inclusion of environmental and interpersonal factors can be helpful to change participants’ behavior [46].

The study participants showed a good adherence to wearing the accelerometer. At baseline, the mean wearing time was on average 5934.5 minutes (SD = 789.5), which corresponds to 14.1 hours per day. Only n = 15 participants had to be excluded from the analysis because of too little wearing time over the six months follow-up time. In conclusion, our data support that the use of an accelerometer is feasible and practical in this age group.

Previous research has shown that intervention studies have the potential to increase PA and decrease PI [37–42]. Most of the previous studies are characterized by rather complex and time-consuming interventions, e.g. personal consultations or telephone calls, which limits the transfer of successful concepts to the general population. Our intervention used individualized feedback-letters based on measured PA and PI and was developed to examine a low-threshold intervention with high practicability and a high potential for translation into the general population. Due to some ceiling effect, the overall intervention effect reduced somewhat.

Especially elderly people are a relevant target group for promoting light PA, because an increase of moderate and vigorous PA might be harder to achieve for this age group [61]. Presently, quantitative recommendations towards PI are lacking and the public health message comes down to “*move more at any intensity*” [6, 15]. Thus, light PA should be included in global recommendations [62].

Several studies have shown that effective interventions to increase PA consider behavioral-change aspects and address participants’ needs and beliefs. Thus, targeting and tailoring are two important aspects of successful interventions [63].

The number of intervention studies promoting PA and reducing PI increased continuously over the last decade. While there is international consensus regarding PA recommendations, it is too early to implement quantified recommendations regarding PI, although some countries

(e.g. Australia) introduced general guidelines to reduce sitting time in general and break up long periods of sitting regularly [25]. In addition, there is an ongoing discussion, whether PI is an independent risk factor for mortality [64], and there is still no global unified definition for PI. Thus, further research regarding PI and specific and quantitative recommendation are urgently needed.

Limitations and strengths

We recruited study participants in a convenience sample using a variety of recruitment methods. Some of these methods have likely increased the proportion of participants with above average PA compared to PA in the general population—so the composition of our study sample somewhat limits the generalizability of our results. Due to this fact, a selection bias can not be excluded. Prior to recruitment, we tried to exclude very active people from the study (based on self-assessment) to address a possible ceiling effect. As the results have shown, many participants underestimate their own amount of PA. Therefore, we calculated a sensitivity analysis excluding very active participants with an amount of ≥ 150 minutes of moderate PA. Some ceiling effect reduced the overall intervention effect somewhat.

In general, accelerometry is an internationally established method for the objective assessment of physical activity and allows a valid and reliable record of PA intensity, frequency, and duration [54, 65]. Although, currently no single specific device can be recommended for all purposes, since the choice of the device depends on the research question, design and target group [54]. However, data from the accelerometer device can potentially differ from the real levels of PA and PI especially in the elderly, because several activities are carried out in standing (i.e. gardening) or sedentary positions (i.e. gymnastic on stools) which cannot be assessed accurately. Only few missings were obtained for the dependent variables ($n = 4$).

Conclusion

Global demographic changes are associated with a large increase of the number of older people and a simultaneous growth of non-communicable-chronic diseases (i.e. cardiovascular disease). Therefore, there is a high need for effective prevention strategies. Although we detected no significant differences in the primary outcome in the interaction effect (time*group) between the study groups, descriptive statistics give a hint of a small positive effect of the intervention. Sensitivity analysis has shown, that partnership is a significant positive factor for light PA (time*partnership). Thus, further research should focus on the recruitment of the adequate target groups (especially people without partner and low levels of PA and high levels of PI). Therefore, local stakeholders should be consulted and setting related factors should be considered.

Most of the successful studies are based on complex and/or resource consuming interventions (i.e. personal consultations), which is accompanied by high costs and a large amount of personal resources. In the future, the usage of internet-based communications and e-health applications (e.g. video-consultation) can support the provision of information and interventions, which makes a translation of (complex) interventions into daily life easier.

Results of our study confirm that a low-threshold intervention with low costs and little amount of personal resources can be implemented and is able to hold up adherence of elderly people. Our analyses shows limited and not significant, but seems to have a small positive effect of a low-threshold intervention after 6 months follow-up.

Supporting information

S1 Table. Descriptive characteristics of the study sample (N = 166), complete and valid data sets (baseline, 3 and 6 months follow-up). Notes: *n* number of subjects, *SD* standard

deviation.
(PDF)

S2 Table. Physical activity and inactivity (mean number of minutes during the wearing-week), by intensity of physical activity and overall PA (N = 166). Notes: n number of subjects, CI 95% confidence interval, M mean.

(PDF)

S3 Table. Number and proportion of study participants who fulfilled the WHO recommendations PA for people aged over 65 years, separately for intervention and control group (N = 166).

(PDF)

S1 File. Dataset.

(XLSX)

Author Contributions

Conceptualization: Fabian Kleinke, Sabina Ulbricht, Marcus Dörr, Wolfgang Hoffmann, Neeltje van den Berg.

Formal analysis: Fabian Kleinke.

Project administration: Fabian Kleinke.

Software: Peter Penndorf.

Supervision: Wolfgang Hoffmann, Neeltje van den Berg.

Writing – original draft: Fabian Kleinke.

Writing – review & editing: Sabina Ulbricht, Marcus Dörr, Wolfgang Hoffmann, Neeltje van den Berg.

References

1. Lollgen H, Lollgen D. [Risk reduction in cardiovascular diseases by physical activity]. *Internist (Berl)*. 2012 Jan; 53(1):20–9. <https://doi.org/10.1007/s00108-011-2889-1> PMID: 22086105
2. American Heart Association. Supporting the national physical activity plan [Internet]. 2016 [cited 2018 Dec 12]. Available from: <https://apha.org/policies-and-advocacy/public-health-policy-statements/policy-database/2014/07/11/16/36/supporting-the-national-physical-activity-plan>
3. World Health Organization. Physical activity for health More active people for a healthier world: draft global action plan on physical activity 2018–2030 [Internet]. 2018. Available from: <http://www.who.int/ncds/governance/who-discussion-paper-gamma-9april2018.pdf?ua=1>
4. Bauman A, Merom D, Bull FC, Buchner DM, Fiatarone Singh MA. Updating the Evidence for Physical Activity: Summative Reviews of the Epidemiological Evidence, Prevalence, and Interventions to Promote “Active Aging.” *Gerontologist* [Internet]. 2016 Apr 1; 56(Suppl_2):S268–80. Available from: <http://dx.doi.org/10.1093/geront/gnw031>
5. Katzmarzyk PT, Lee I-M. Sedentary behaviour and life expectancy in the USA: a cause-deleted life table analysis. *BMJ Open* [Internet]. 2012 Jan 1; 2(4):e000828. Available from: <https://doi.org/10.1136/bmjopen-2012-000828> PMID: 22777603
6. Stamatakis E, Ekelund U, Ding D, Hamer M, Bauman AE, Lee I-M. Is the time right for quantitative public health guidelines on sitting? A narrative review of sedentary behaviour research paradigms and findings. *Br J Sports Med* [Internet]. 2018; Available from: <https://bjsm.bmj.com/content/early/2018/06/08/bjsports-2018-099131> <https://doi.org/10.1136/bjsports-2018-099131> PMID: 29891615
7. Lee I-M, Shiroma EJ, Lobelo F, Puska P, Blair SN, Katzmarzyk PT. Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *Lancet* [Internet]. 2012 Jul 21; 380(9838):219–29. Available from: [https://doi.org/10.1016/S0140-6736\(12\)61031-9](https://doi.org/10.1016/S0140-6736(12)61031-9) PMID: 22818936

8. Sedentary Behaviour Research Network. Letter to the Editor: Standardized use of the terms “sedentary” and “sedentary behaviours.” *Appl Physiol Nutr Metab* [Internet]. 2012 Apr 27; 37(3):540–2. Available from: <https://doi.org/10.1139/h2012-024> PMID: 22540258
9. Tremblay MS, Aubert S, Barnes JD, Saunders TJ, Carson V, Latimer-Cheung AE, et al. Sedentary Behavior Research Network (SBRN)—Terminology Consensus Project process and outcome. *Int J Behav Nutr Phys Act* [Internet]. 2017 Jun 10; 14(1):75. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28599680> <https://doi.org/10.1186/s12966-017-0525-8> PMID: 28599680
10. Hallal PC, Andersen LB, Bull FC, Guthold R, Haskell W, Ekelund U. Global physical activity levels: surveillance progress, pitfalls, and prospects. *Lancet* [Internet]. 2012; 380(9838):247–57. Available from: <http://www.sciencedirect.com/science/article/pii/S0140673612606461> [https://doi.org/10.1016/S0140-6736\(12\)60646-1](https://doi.org/10.1016/S0140-6736(12)60646-1) PMID: 22818937
11. Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* [Internet]. 2012 Dec 15; 380(9859):2224–60. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4156511/>
12. Ding D, Lawson KD, Kolbe-Alexander TL, Finkelstein EA, Katzmarzyk PT, van Mechelen W, et al. The economic burden of physical inactivity: a global analysis of major non-communicable diseases. *Lancet* [Internet]. 2016; 388(10051):1311–24. Available from: <http://www.sciencedirect.com/science/article/pii/S014067361630383X> [https://doi.org/10.1016/S0140-6736\(16\)30383-X](https://doi.org/10.1016/S0140-6736(16)30383-X) PMID: 27475266
13. World Health Organization. Global Status Report on noncommunicable diseases 2014. Geneva; 2014.
14. Varo JJ, Martinez-Gonzalez MA, De Irala-Esteviz J, Kearney J, Gibney M, Martinez JA. Distribution and determinants of sedentary lifestyles in the European Union. *Int J Epidemiol*. 2003 Feb; 32(1):138–46. <https://doi.org/10.1093/ije/dyg116> PMID: 12690026
15. Stamatakis E, Gill JMR. Sitting behaviour and physical activity: two sides of the same cardiovascular health coin? *Br J Sports Med* [Internet]. 2018 Jul 9; bjsports-2018-099640. Available from: <http://bjsm.bmj.com/content/early/2018/07/10/bjsports-2018-099640.abstract> <https://doi.org/10.1136/bjsports-2018-099640> PMID: 29991572
16. Gebel K, Ding D, Chey T, Stamatakis E, Brown WJ, Bauman AE. Effect of Moderate to Vigorous Physical Activity on All-Cause Mortality in Middle-aged and Older Australians Physical Activity and Mortality-Physical Activity and Mortality. *JAMA Intern Med* [Internet]. 2015 Jun 1; 175(6):970–7. Available from: <https://doi.org/10.1001/jamainternmed.2015.0541> PMID: 25844882
17. World Health Organization. Cardiovascular diseases (CVDs) [Internet]. 2017 [cited 2018 Oct 17]. Available from: [http://www.who.int/en/news-room/fact-sheets/detail/cardiovascular-diseases-\(cvds\)](http://www.who.int/en/news-room/fact-sheets/detail/cardiovascular-diseases-(cvds))
18. Arem H, Moore SC, Patel A, Hartge P, Berrington de Gonzalez A, Viswanathan K, et al. Leisure time physical activity and mortality: a detailed pooled analysis of the dose-response relationship. *JAMA Intern Med*. 2015 Jun; 175(6):959–67. <https://doi.org/10.1001/jamainternmed.2015.0533> PMID: 25844730
19. Ekelund U, Tarp J, Steene-Johannessen J, Hansen BH, Jefferis B, Fagerland MW, et al. Dose-response associations between accelerometry measured physical activity and sedentary time and all cause mortality: systematic review and harmonised meta-analysis. *BMJ* [Internet]. 2019; 366. Available from: <https://www.bmj.com/content/366/bmj.l4570> <https://doi.org/10.1136/bmj.l4570> PMID: 31434697
20. Miles L. Physical activity and health. *Br Nutr Found Nutr Bull*. 2007; 32:314–363.
21. Gallanagh S, Quinn TJ, Alexander J, Walters MR. Physical activity in the prevention and treatment of stroke. *ISRN Neurol* [Internet]. 2011/10/01. 2011; 2011:953818. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/22389836> <https://doi.org/10.5402/2011/953818> PMID: 22389836
22. Shiroma EJ, Lee I-M, Schepps MA, Kamada M, Harris TB. Physical Activity Patterns and Mortality: The Weekend Warrior and Activity Bouts. *Med Sci Sports Exerc*. 2019 Jan; 51(1):35–40. <https://doi.org/10.1249/MSS.0000000000001762> PMID: 30138219
23. United States Department of Health and Human Services. Physical Activity Guidelines [Internet]. 2018 [cited 2018 Nov 6]. Available from: <https://health.gov/paguidelines/guidelines/older-adults.aspx>
24. Bull FC; Expert Working Groups. Physical Activity Guidelines in the U.K.: Review and Recommendations. Loughborough, England; 2010.
25. Department of Health. Australia’s Physical Activity and Sedentary Behaviour Guidelines. Canberra, Australia; 2014.
26. World Health Organization. Global Recommendations on Physical Activity for Health. 65 years and above [Internet]. 2011 [cited 2018 Jul 18]. Available from: <http://www.who.int/dietphysicalactivity/physical-activity-recommendations-65years.pdf?ua=1>
27. US Department of Health and Human Services. 2008 Physical Activity Guidelines for Americans. Washington, DC; 2008.

28. Loprinzi PD. Light-Intensity Physical Activity and All-Cause Mortality. *Am J Health Promot.* 2017 Jul; 31(4):340–2. <https://doi.org/10.4278/ajhp.150515-ARB-882> PMID: 26730555
29. Füzéki E, Engeroff T, Banzer W. Health Benefits of Light-Intensity Physical Activity: A Systematic Review of Accelerometer Data of the National Health and Nutrition Examination Survey (NHANES). Vol. 47, *Sports Medicine*. Springer International Publishing; 2017. p. 1769–93.
30. Johnson LG, Butson ML, Polman RC, Raj IS, Borkoles E, Scott D, et al. Light physical activity is positively associated with cognitive performance in older community dwelling adults. *J Sci Med Sport.* 2016 Nov; 19(11):877–82. <https://doi.org/10.1016/j.jsams.2016.02.002> PMID: 26922133
31. Bae W, Ik Suh Y, Ryu J, Heo J. Physical Activity Levels and Well-Being in Older Adults. *Psychol Rep.* 2017 Apr; 120(2):192–205. <https://doi.org/10.1177/0033294116688892> PMID: 28558621
32. Althoff T, Sosič R, Hicks JL, King AC, Delp SL, Leskovec J. Large-scale physical activity data reveal worldwide activity inequality. *Nature [Internet].* 2017/07/10. 2017 Jul 20; 547(7663):336–9. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28693034> <https://doi.org/10.1038/nature23018> PMID: 28693034
33. Salive ME. Multimorbidity in older adults. *Epidemiol Rev.* 2013; 35:75–83. <https://doi.org/10.1093/epirev/mxs009> PMID: 23372025
34. Barnett K, Mercer SW, Norbury M, Watt G, Wyke S, Guthrie B. Epidemiology of multimorbidity and implications for health care, research, and medical education: a cross-sectional study. *Lancet (London, England).* 2012 Jul; 380(9836):37–43.
35. Kirchberger I, Meisinger C, Heier M, Zimmermann A-K, Thorand B, Autenrieth CS, et al. Patterns of multimorbidity in the aged population. Results from the KORA-Age study. *PLoS One.* 2012; 7(1): e30556. <https://doi.org/10.1371/journal.pone.0030556> PMID: 22291986
36. Gosswald A, Schienkiewitz A, Nowossadeck E, Busch MA. [Prevalence of myocardial infarction and coronary heart disease in adults aged 40–79 years in Germany: results of the German Health Interview and Examination Survey for Adults (DEGS1)]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz.* 2013 May; 56(5–6):650–5. <https://doi.org/10.1007/s00103-013-1666-9> PMID: 23703482
37. Harris T, Kerry SM, Victor CR, Ekelund U, Woodcock A, Iliffe S, et al. A Primary Care Nurse-Delivered Walking Intervention in Older Adults: PACE (Pedometer Accelerometer Consultation Evaluation)-Lift Cluster Randomised Controlled Trial. *PLOS Med [Internet].* 2015 Feb 17; 12(2):e1001783. Available from: <https://doi.org/10.1371/journal.pmed.1001783> PMID: 25689364
38. Kang M, Marshall SJ, Barreira T V, Lee J-O. Effect of pedometer-based physical activity interventions: a meta-analysis. *Res Q Exerc Sport.* 2009 Sep; 80(3):648–55. <https://doi.org/10.1080/02701367.2009.10599604> PMID: 19791652
39. Bravata DM, Smith-Spangler C, Sundaram V, Gienger AL, Lin N, Lewis R, et al. Using pedometers to increase physical activity and improve health: a systematic review. *JAMA.* 2007 Nov; 298(19):2296–304. <https://doi.org/10.1001/jama.298.19.2296> PMID: 18029834
40. Rosenberg DE, Gell NM, Jones SMW, Renz A, Kerr J, Gardiner PA, et al. The Feasibility of Reducing Sitting Time in Overweight and Obese Older Adults. *Heal Educ Behav Off Publ Soc Public Heal Educ.* 2015 Oct; 42(5):669–76. <https://doi.org/10.1177/1090198115577378> PMID: 25794518
41. Gardiner PA, Eakin EG, Healy GN, Owen N. Feasibility of reducing older adults' sedentary time. *Am J Prev Med.* 2011 Aug; 41(2):174–7. <https://doi.org/10.1016/j.amepre.2011.03.020> PMID: 21767725
42. Burke L, Lee AH, Jancey J, Xiang L, Kerr DA, Howat PA, et al. Physical activity and nutrition behavioural outcomes of a home-based intervention program for seniors: a randomized controlled trial. *Int J Behav Nutr Phys Act.* 2013 Jan; 10:14. <https://doi.org/10.1186/1479-5868-10-14> PMID: 23363616
43. Prince SA, Saunders TJ, Gresty K, Reid RD. A comparison of the effectiveness of physical activity and sedentary behaviour interventions in reducing sedentary time in adults: a systematic review and meta-analysis of controlled trials. *Obes Rev [Internet].* 2014 Aug 11; 15(11):905–19. Available from: <https://doi.org/10.1111/obr.12215> PMID: 25112481
44. Mutrie N, Doolin O, Fitzsimons CF, Grant PM, Granat M, Grealay M, et al. Increasing older adults' walking through primary care: results of a pilot randomized controlled trial. *Fam Pract.* 2012 Dec; 29(6):633–42. <https://doi.org/10.1093/fampra/cms038> PMID: 22843637
45. Kleinke F, Schwaneberg T, Weymar F, Penndorf P, Ulbricht S, Lehnert K, et al. MOVING: Motivation-Oriented interVention study for the elderly IN Greifswald: study protocol for a randomized controlled trial. *Trials.* 2018 Jan; 19(1):57. <https://doi.org/10.1186/s13063-017-2425-2> PMID: 29357943
46. Kleinke F, Penndorf P, Ulbricht S, Dörr M, Hoffmann W, van den Berg N. Levels of and determinants for physical activity and physical inactivity in a group of healthy elderly people in Germany: Baseline results of the MOVING-study. *PLoS One [Internet].* 2020 Aug 13; 15(8):e0237495. Available from: <https://doi.org/10.1371/journal.pone.0237495> PMID: 32790711

47. Meyer J, Fredrich D, Piegsa J, Habes M, van den Berg N, Hoffmann W. A mobile and asynchronous electronic data capture system for epidemiologic studies. *Comput Methods Programs Biomed.* 2013 Jun; 110(3):369–79. <https://doi.org/10.1016/j.cmpb.2012.10.015> PMID: 23195493
48. The International Physical Activity Questionnaire group. INTERNATIONAL PHYSICAL ACTIVITY QUESTIONNAIRE SHORT LAST 7 DAYS SELF-ADMINISTERED FORMAT- German Version [Internet]. 2018 [cited 2018 Oct 30]. Available from: <https://docs.google.com/viewer?a=v&pid=sites&srcid=ZGVmYXVsdGRvbWFPbnx0aGVpcGFfGd4OjNiMGRjYjU3NzE5Nzc0ZGY>
49. Leibnitz-Zentrum für Psychologische Information und Dokumentation (ZPID). German—PAQ—50+: Fragebogen zur Erfassung der körperlichen Aktivität [Internet]. 2015 [cited 2018 Oct 30]. Available from: https://www.psycharchives.org/bitstream/20.500.12034/407/1/PT_9006067_German-PAQ-50%2B_Fragebogen.pdf
50. Gardiner PA, Clark BK, Healy GN, Eakin EG, Winkler EAH, Owen N. Measuring older adults' sedentary time: reliability, validity, and responsiveness. *Med Sci Sports Exerc.* 2011 Nov; 43(11):2127–33. <https://doi.org/10.1249/MSS.0b013e31821b94f7> PMID: 21448077
51. Beierlein V, Morfeld M, Bergelt C, Bullinger M, Brähler E. Messung der gesundheitsbezogenen Lebensqualität mit dem SF-8. *Diagnostica* [Internet]. 2012; 58(3):145–53. Available from: <https://doi.org/10.1026/0012-1924/a000068>
52. Fuchs R, Schwarzer R. Selbstwirksamkeit zur sportlichen Aktivität: Reliabilität und Validität eines neuen Meßinstruments. *Z Differ Diagn Psychol.* 1994; 15:141–54.
53. Freedson PS, Melanson E, Sirard J. Calibration of the Computer Science and Applications, Inc. accelerometer. *Med Sci Sports Exerc.* 1998 May; 30(5):777–81. <https://doi.org/10.1097/00005768-199805000-00021> PMID: 9588623
54. Gabrys L, Thiel C, Tallner A, Wilms B, Müller C, Kahler D, et al. Akzelerometrie zur Erfassung körperlicher Aktivität. *Sportwissenschaft* [Internet]. 2014; 45(1):1–9. Available from: <https://doi.org/10.1007/s12662-014-0349-5>
55. Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical activity in the United States measured by accelerometer. *Med Sci Sports Exerc.* 2008 Jan; 40(1):181–8. <https://doi.org/10.1249/mss.0b013e31815a51b3> PMID: 18091006
56. Krug S, Jordan S, Mensink GBM, Muters S, Finger J, Lampert T. [Physical activity: results of the German Health Interview and Examination Survey for Adults (DEGS1)]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitschutz.* 2013 May; 56(5–6):765–71. <https://doi.org/10.1007/s00103-012-1661-6> PMID: 23703496
57. Lohne-Seiler H, Hansen BH, Kolle E, Anderssen SA. Accelerometer-determined physical activity and self-reported health in a population of older adults (65–85 years): a cross-sectional study. *BMC Public Health* [Internet]. 2014; 14(1):284. Available from: <https://doi.org/10.1186/1471-2458-14-284>
58. Harland J, White M, Drinkwater C, Chinn D, Farr L, Howel D. The Newcastle exercise project: a randomised controlled trial of methods to promote physical activity in primary care. *BMJ.* 1999 Sep; 319(7213):828–32. <https://doi.org/10.1136/bmj.319.7213.828> PMID: 10496829
59. Marcus BH, Bock BC, Pinto BM, Forsyth LH, Roberts MB, Traficante RM. Efficacy of an individualized, motivationally-tailored physical activity intervention. *Ann Behav Med.* 1998; 20(3):174–80. <https://doi.org/10.1007/BF02884958> PMID: 9989324
60. Matthews CE, Chen KY, Freedson PS, Buchowski MS, Beech BM, Pate RR, et al. Amount of time spent in sedentary behaviors in the United States, 2003–2004. *Am J Epidemiol.* 2008 Apr; 167(7):875–81. <https://doi.org/10.1093/aje/kwm390> PMID: 18303006
61. Westerterp KR. Physical activity as determinant of daily energy expenditure. *Physiol Behav.* 2008 Mar; 93(4–5):1039–43. <https://doi.org/10.1016/j.physbeh.2008.01.021> PMID: 18308349
62. Füzéki E, Engeroff T, Banzer W. Health Benefits of Light-Intensity Physical Activity: A Systematic Review of Accelerometer Data of the National Health and Nutrition Examination Survey (NHANES). *Sport Med* [Internet]. 2017; 47(9):1769–93. Available from: <https://doi.org/10.1007/s40279-017-0724-0>
63. Ogilvie D, Foster CE, Rothnie H, Cavill N, Hamilton V, Fitzsimons CF, et al. Interventions to promote walking: systematic review. *BMJ.* 2007 Jun; 334(7605):1204. <https://doi.org/10.1136/bmj.39198.722720.BE> PMID: 17540909
64. Ekelund U, Steene-Johannessen J, Brown WJ, Fagerland MW, Owen N, Powell KE, et al. Does physical activity attenuate, or even eliminate, the detrimental association of sitting time with mortality? A harmonised meta-analysis of data from more than 1 million men and women. *Lancet* [Internet]. 2016; 388(10051):1302–10. Available from: <http://www.sciencedirect.com/science/article/pii/S0140673616303701> [https://doi.org/10.1016/S0140-6736\(16\)30370-1](https://doi.org/10.1016/S0140-6736(16)30370-1) PMID: 27475271
65. Schrack JA, Cooper R, Koster A, Shiroma EJ, Murabito JM, Rejeski WJ, et al. Assessing Daily Physical Activity in Older Adults: Unraveling the Complexity of Monitors, Measures, and Methods. *J Gerontol A Biol Sci Med Sci.* 2016 Aug; 71(8):1039–48. <https://doi.org/10.1093/gerona/glw026> PMID: 26957472