

5. Literaturverzeichnis

- 01 Albrecht T, Hoffmann CW, Schettler S, Overberg A, Ilg M, Wolf K (1999).
Improved detection of liver metastases with phase inversion imaging during the
specific phase of the ultrasound contrast agent Levovist.
Eur. Radiol(9), S. 388.
- 02 Bauer A, Blomley M, Leen E, Cosgrove D, Schlieff R (1999).
Liver-specific imaging with SHU 563A: Diagnostic potential of new class of ultrasound
contrast media.
Eur. Radiol(9), S. 349-352.
- 03 Bluemke DA, Weber TM, Rubin D, de Lange EE, Semelka R, Redvanly RD, Chezmar
J, Outwater E, Carlos R, Saini S, Holland GA, Mammone JF, Brown JJ, Milestone B,
Javitt MC, Jacobs P (2003).
Hepatic MR imaging with ferumoxides: multicenter study of safety and effectiveness
of direct injection protocol.
Radiology(228), S. 457-464.
- 04 Boutkan H, Luth W, Meyer S, Cuesta E, van Heuzen W, Prevoo W (1992).
The impact of intraoperative ultrasonography of the liver on the surgical strategy of
patients with gastrointestinal malignancies and hepatic metastases.
Eur J Surg Oncol(18), S. 342-346.
- 05 Brandl H, Gritzky A, Haizinger M (1999).
3D ultrasound: a dedicated system.
Eur Radiol(9), S. 331-333.
- 06 Bundesärztekammer (Geschäftsführung) (2000).
Leitlinien der Bundesärztekammer zur Qualitätsicherung in der
Magnetresonanztomographie (Bechluß 29.11.1999).
Deutsches Ärzteblatt 97(39), S. A 2557-2568.
- 07 Carlos RC, Branam JD, Dong Q, Hussain HK, Francis IR (2002).
Biliary imaging with Gd-EOB-DTPA: is a 20 minutes delay sufficient?
Acad Radiol(9), S. 1322-1325.

- 08 Clement O, Muhler A, Vexler V, Berthezene Y, Brasch RC (1992).
Gadolinium – ethoxybenzyl – DTPA, a new liver specific magnetic resonance contrast agent. Kinetic and enhancement patterns in normal and cholestatic rats.
Invest Radiol(27), S. 612-619.
- 09 Coche EE, Bret PM, Reinhold C (1999).
Dynamic enhancement of upper abdominal organs in normal volunteers with MRI and effects of contrast dose reduction.
Abd Imag(24), S. 604-609.
- 10 Eggebo TM, Sorvang S, Dalaker K (1990).
Ultrasonic diagnosis of the upper abdomen performed in general practice.
Tidsskr Nor Laegeforen(110), S. 1096-1108.
- 11 Fuhrmann GM, Charnsangavej C, Abbruzzese JL et al. (1994).
Thin section contrast enhanced computed tomography accurately predicts the resectability of malignant pancreas neoplasm.
Am J Surg(167), S. 104-113.
- 12 Giovagnoni A, Paci E (1996).
Liver III: Gadolinium – based hepatobiliary contrast agents (Gd-EOB-DTPA and Gd-BOPTA/Dimeg.).
Magn Reson Imag Clin N Am(4), S. 61-72.
- 13 Glazer GM, Axel L, Goldbert HJ, Moss AA (1981).
Dynamic CT of the normal spleen.
AJR(137), S. 343-346.
- 14 Grutzmann R, Bunk A, Kersting S, Pilarsky C, Dobrowolski F, Kuhlisch E, Ockert D, Saeger HD (2003).
Prospective evaluation of ultrasound and colour duplex imaging for the assessment of surgical resectability of pancreatic tumours.
Langenbecks Arch Surg(388), S. 392-400.
- 15 Hagspiel KD, Neidl KFW, Eichenberger AC, Weder W, Marincek B (1995).
Detection of Liver Metastases: Comparison of Superparamagnetic Iron Oxide-enhanced and enhanced MR Imaging at 1.5T with Dynamic CT, Intraoperative US, and Percutaneous US.
Radiology(196), S. 471-478.

- 16 Hamed MM, Hamm B, Ibrahim ME, Taupitz M, Mahfouz AE (1992).
Dynamic MR imaging of the abdomen with gadopentetate dimeglumine: normal enhancement patterns of liver, spleen, stomach, and pancreas.
AJR(158), S. 303-307.
- 17 Hamm B, Mahfouz AE, Taupitz M, Mitchell DG, Nelson R, Halpern E, Speidel A, Wolf KJ, Saini S (1997).
Liver metastases: improved detection with dynamic gadolinium-enhanced MR imaging?
Radiology(202), S. 667-682.
- 18 Hamm B, Staks T, Mühler A, Bollow M, Taupitz M, Frenzel T, Wolf K, Weinmann HJ, Lange L (1995).
Phase I Clinical Evaluation of Gd-EOB-DTPA as a Hepatobiliary MR Contrast Agent: Safety, Pharmacocinetics, and MR-Imaging.
Radiology(195), S. 785-792.
- 19 Helmberger T, Gregor M, Holzknecht N, Gauger J, Rau H, Reiser M (1999).
Detektion und Charakterisierung fokaler Leberläsionen: Vergleich von biphasischer, kontrastmittelverstärkter Spiral-CT, nativer und SPIO-verstärkter MRT: Fortschritte in der Bildgebung des Abdomens.
Radiologe(39), S. 678-684.
- 20 Hommeyer SC, Freeny PC, Crabo LG (1995).
Carcinoma of the head of the pancreas: evaluation of the pancreaticoduodenal veins with dynamic CT – potential for improved accuracy in staging.
Radiology(196), S. 233-238.
- 21 Huppertz A (1997).
Beschreibung von Eovist® (Gd-EOB-DTPA / SH L 569 B) Vergleich der Verträglichkeit und Kinetik MRI Leberkontrastmittel Resovist® (SH U 566 A) und Eovist® (SH L 569B).
Aachen Shaker Verlag, S. 30-31.

- 22 Huppertz A, Balzer T, Blakeborough A, Breuer J, Glovagnoni A, Heinz-Peer G, Laniado M, Manfredi RM, Mathieu DG, Mueller D, Reimer P, Robinson PJ, Strotzer M, Taupitz M, Vogl TJ (2004).
Improved Detection of Focal Liver Lesions at MR Imaging: Multicenter Comparison of Gadoteric Acid-enhanced MR Images with Intraoperative Findings (For European EOB Study Group).
Radiology(230), S. 266-275.
- 23 Kanematsu M, Semelka RC, Matsuo M, Kondo H, Enya M, Goshima S, Moriyama N, Hoshi H (2002).
Gadolinium-enhanced MR imaging of the liver: optimizing imaging delay for hepatic arterial and portal venous phases – a prospective randomized study in patient with chronic liver damage.
Radiology(225), S. 407-415.
- 24 Kanematsu M, Shiratori Y, Hoshi H, Kondo H, Matsuo M, Moriwaki H (2000).
Pancreas and Peripancreatic Vessels: effect of imaging Delay on Gadolinium Enhancement at Dynamic Gradient-Recalled-Echo MR Imaging.
Radiology(215), S. 95-102.
- 25 Knopp MV, Dorsam J, Oesingmann N et al. (1997).
Funktionelle MR-Urographie bei Patienten mit Nierentransplantaten.
Radiologe(37), S. 233-238.
- 26 Konopke R, Kersting S, Bergert H, Bloomenthal A, Gastmeier J, Saeger HD, Bunk A (2006).
Contrast-enhanced ultrasonography to detect liver metastases : A prospective trial to compare transcutaneous unenhanced and contrast-enhanced ultrasonography in patients undergoing laparotomy.
Int J Colorect Dis.
- 27 Laniado L, Kopp A (1997).
Liver-Specific Contrast-Media: A Magic Bullet or a Weapon for Dedicated Targets.
Radiology(205), S. 319-322.

- 28 Leen E, Angerson W, Wotherspoon H, Moule B, Cook T, Mc Ardle C (1995).
Detection of Colorectal Liver Metastases: Comparison of Laparotomy, CT, US, and
Doppler Perfusion Index and Evaluation of Postoperative Follow-up Results.
Radiology(195), S. 113-116.
- 29 Low RN, Martinez AG, Steinberg SM et al. (1998).
Potential renal transplant donors: evaluation with gadolinium enhanced MR
angiography and MR urography.
Radiology(207), S. 165-172.
- 30 Martin J, Sentis M, Puig J, Rue M, Falco J, Donoso J, Zidan A (1996).
Comparison of in-phase and opposed phase GRE and conventional SE MR pulse
sequences in T1-weighted imaging of liver lesions.
J Comp Assist Tomogr(20), S. 890-897.
- 31 Mirowitz SA, Brown JJ, Lee JK, et al. (1991).
Dynamic gadolinium enhanced MR imaging of the spleen: normal patterns and
evaluation of splenic lesions.
Radiology(179), S. 681-686.
- 32 Mirowitz SA, Gutierrez E, Lee JKT, Brown JJ, Heiken JP (1991).
Normal Abdominal Enhancement Patterns with Dynamic Gadolinium-enhanced MR
imaging.
Radiology(180), S. 637-640.
- 33 Muhler A, Clement O, Vexler V, Berthezene Y, Rosenau W, Brasch RC (1992).
Hepatobiliary enhancement with Gd-EOB-DTPA: comparison of spin-echo and STIR
imaging for detection of experimental liver metastases.
Radiology(184), S. 207-213.
- 34 Muhler A, Freise CE, Kuwatsuru R et al. (1993).
Acute liver rejection: evaluation with cell directed MR contrast agents in a rat
transplantation model.
Radiology(186), S. 139-146.

- 35 Ni Y, Marchal G, Lukito G, Yu J, Muhler A, Baert AL (1994).
MR imaging evaluation of liver enhancement by Gd-EOB-DTPA in selective and total bile duct obstruction in rats: correlation with serologic, mikrocholangiographic, and histological findings.
Radiology(190), S. 753-758.
- 36 Niendorf HP, Haustein J, Cornelius I, Alhassan A, Clauss W (1991).
Safety of gadolinium – DTPA: Extended clinical experience.
Magn Reson in Med(22), S. 222-228.
- 37 Paaajanen H, Brasch RC, Dean PB (1988).
Experimental acute pancreatitis: MR relaxation time studies using gadolinium DTPA.
Magn Reson Med(6), S. 63-73.
- 38 Reimer P, Jähnke N, Fiebich M, Schima W, Deckers F, Marx C, Holzknacht N, Saini S (2000).
Hepatic Lesion Detection and Characterization: Value of Nonenhanced MR imaging, Superparamagnetic Iron Oxide-Enhanced MR Imaging, and Spiral CT- ROC Analysis.
Radiology(217), S. 152-158.
- 39 Reimer P, Rummeny EJ, Shamsi K, Balzer T, Daldrup HE, Tombach B, Hesse T, Berns T, Peters PE (1996).
Phase II Clinical Evaluation of Gd-EOB-DTPA Dose, Safety Aspects, and pulse Sequence.
Radiology(199), S. 177-183.
- 40 Reimer P; Vosshenrich R (2004).
Kontrastmittel in der MRT: Substanzen, Wirkungen, Pharmakologie und Zulassung.
Radiologe(44), S. 273-283.
- 41 Ren Y, Jin ZY, Xue HD, You H, Feng F (2006).
Comparison of multi-slice spiral CT and magnetic resonance imaging in evaluation of the un-resectability of blood vessels in pancreatic tumor.
Zhongguo Yi Xue Ke Xue Yuan Xue Bao(28), S. 76-79.
- 42 Sadick M, Diehl SJ, Lehmann KJ, Gaa J, Mockel R, Georgi M (2000).
Evaluation of breath-hold contrast-enhanced 3D magnetic resonance angiography technique for imaging visceral abdominal arteries and veins.
Invest. Radiol.(35), S. 111-117.

- 43 Schuhmann-Giampieri G, Schmitt-Willich H, Press WR, Negishi C, Weinmann HJ, Speck U (1992).
Preclinical evaluation of Gd-EOB-DTPA as a contrast agent in MR imaging of the hepatobiliary system.
Radiology(183), S. 59-64.
- 44 Semelka RC, Kroeker MA, Shoenut JP, et al. (1991).
Pancreatic disease: prospective comparison of CT, ERCP and 1,5T MR imaging with dynamic gadolinium enhancement and fat suppression.
Radiology(181), S. 785-791.
- 45 Simon G, Link TM, Wortler K, Doebereiner F, Schulte-Frohlinde E, Daldrup-Link H, Settles M, Rummeny EJ (2005).
Detection of hepatocellular carcinoma: comparison of Gd-DTPA- and ferumoxides-enhanced MR imaging.
Eur Radiol(15), S. 895-903.
- 46 Sironyi S, De Cobelli F, Zebri A, Balzano G, Di Carlo V, Del Maschio A (1995).
Pancreatic carcinoma: MR assessment of tumor invasion of the peripancreatic vessels.
J Comput Assist Tomogr(19), S. 739-744.
- 47 Stroszczyński C, Gaffke G, Gnauck M, Streitparth F, Wieners G, Lopez-Häninnen E (2004).
Aktueller Stand der MRT Diagnostik mit leberspezifischen Kontrastmitteln. Gd-EOB-DTPA und Gd-BOPTA.
Radiologe(44), S. 1185-1191.
- 48 Tanimoto A, Sotah Y, Yuasa Y, Jinzaki M, Hiramatsu K (1997).
Performance of Gd-EOB-DTPA and superparamagnetic iron oxide particles in the detection of primary liver cancer: a comparative study by alternative free-response receiver operating characteristic analysis.
J Magn Reson Imag(7), S. 120-124.
- 49 v. Montfoort JE, S. B (1999).
Hepatic uptake of the magnetic resonance imaging contrast agent gadoxetate by the organic anion transporting polypeptide OATP1.
JPET(290), S. 153-157.

- 50 Vallet AD, Romano W, Bach DB, Passi RB, Taves DH, Munk PL (1992).
Adenocarcinoma of the pancreas ducts: comparative evaluation with CT and MR
imaging at 1.5T.
Radiology(183), S. 87-95.
- 51 Vogl TJ, Kummel S, Hammerstingl R, Schellenbeck M, Schumacher G, Balzer T,
Schwarz W, Muller PK, Bechstein WO, Mack MG, Sollner O, Felix R (1996).
Liver tumors: comparison of MR imaging with Gd-EOB-DTPA and Gd-DTPA.
Radiology(200), S. 59-67.
- 52 Ward J, Naik D, Guthrie A, Wilson D, Robinson PJ (1999).
Hepatic Lesion Detection: Comparison of MR Imaging after Administration of
Superparamagnetic Iron Oxide with Dual-Phase CT by Using Alternative – Free
Response Receiver Operating Characteristic Analysis.
Radiology(210), S. 459-466.
- 53 Weinmann HJ, Schuhmann-Giampieri G, Schmitt-Willich H, Vogler H, Frenzel T,
Gries H (1991).
A new lipophilic gadolinium chelate as a tissue-specific contrast medium for MRI.
Magn Reson Med(22), S. 223-237.
- 54 Whittingham TA (1999).
Tissue harmonic imaging.
Eur. Radiol(9), S. 323-326.
- 55 Zangos S, Hammerstingl R, Mack MG, Straub R, Engelmann K, Eichler K, Balzer T,
Vogl TJ (2001).
Renales Anreicherungsverhalten und Elimination des hepatobiliären Kontrastmittels
Gd-EOB-DTPA in der Magnetresonanztomographie.
ROFO(173), S. 1034-1040.